

Supplementary material has been published as submitted. It has not been copyedited or typeset by Scandinavian Journal of Urology.

## SUPPLEMENTARY TABLES

	OS from index date (95% CI)		
	Overall	T2	T3/T4a
<b>1-year OS, %</b>	52 (47-57)	53 (48-60)	41 (30-57)
<b>3-year OS, %</b>	24 (20-29)	24 (20-30)	21 (13-36)
<b>5-year OS, %</b>	18 (14-23)	18 (14-24)	16 (8-31)
<b>Median OS, months</b>	12 (11-15)	13 (11-16)	9 (6-17)

**Supplementary Table 1. Overall survival (OS) rate estimates and median OS in cystectomy-ineligible/refusal muscle-invasive bladder cancer (MIBC) patients.**

CI = confidence interval; OS = overall survival.

	BI-EFS from index date (95% CI)		
	Overall	T2	T3/T4a
<b>1-year BI-EFS, %</b>	44 (39-50)	46 (41-52)	33 (23-49)
<b>3-year BI-EFS, %</b>	21 (17-26)	21 (17-27)	17 (9-32)
<b>5-year BI-EFS, %</b>	15 (11-20)	15 (11-20)	15 (7-29)
<b>Median BI-EFS, months</b>	9 (8-12)	10 (9-13)	6 (3-12)

**Supplementary Table 2. Bladder-intact event-free survival (BI-EFS) rate estimates and median BI-EFS in cystectomy-ineligible/refusal MIBC patients.**

BI-EFS = bladder-intact, event-free survival; CI = confidence interval.

	Overall		T2		T3/T4a	
BI-EFS event	n (%)	Median time-to-event, months (95% CI)	n (%)	Median time-to-event, months (95% CI)	n (%)	Median time-to-event, months (95% CI)
Proxies for histologically proven residual/recurrent MIBC and clinical evidence of nodal/metastatic disease	43 (15)	0.89 (0.66-0.92)	33 (13)	0.92 (0.66-1.02)	10 (23)	0.66 (0.35-0.92)
Radical cystectomy	4 (1)	10.1 (6.9-13.1)	4 (2)	10.1 (6.9-13.1)	0 (0)	---
Death	245 (84)	8.8 (7.1-10.3)	211 (85)	9.14 (7.6-10.8)	34 (77)	6.2 (2.8-8.4)

**Supplementary Table 3. Types of events captured for BI-EFS analysis.**

BI-EFS = bladder-intact, event-free survival; CI = confidence interval; MIBC = muscle-invasive bladder cancer.