

Supplemental table 1

	Before SCP introduction 2011-2014 numbers (%)	After SCP introduction 2016-2019 numbers (%)
Males stratified by age^		
<60	78/721 (11)	30/809 (4)
60-69	248/1974 (13)	118/1904 (6)
70-79	345/2771 (15)	262/3607 (7)
80-89	258/1771 (15)	166/2037 (8)
90 and above	41/251 (16)	26/301 (9)
Females stratified by age^		
<60	26/283 (9)	15/314 (5)
60-69	66/632 (10)	35/665 (5)
70-79	98/806 (12)	78/1159 (7)
80-89	85/648 (13)	50/719 (7)
90 and above	22/127 (17)	23/132 (17)
Educational level*		
Low	532/3981 (13)	310/4011 (8)
Middle	476/3888 (12)	324/4868 (7)
High	236/1989 (12)	153/2600 (6)
DCI (tertiles)		
-0.75	409/3449 (12)	242/4106 (6)
0.75-2	371/3017 (12)	241/3440 (7)
2-	487/3532 (14)	320/4101 (8)
Anticoagulants at diagnosis		
No	697/5536 (13)	471/7146 (7)
Platelet aggregation-inhibitors	393/3169 (12)	226/3284 (7)
Other	177/1293 (14)	106/1217 (9)
Healthcare region		
Stockholm	358/1842 (19)	168/2286 (7)
Mid	252/2011 (13)	193/2363 (8)
Southeastern	94/1205 (8)	43/1348 (3)
Southern	251/2080 (12)	227/2451 (9)
Western	202/1906 (11)	100/2161 (5)
Northern	110/954 (12)	72/1028 (7)

Supplemental Table 1. Proportion unscheduled hospitalization before and after the introduction of standardized care pathways (SCP). DCI = Drug Comorbidity Index; ^ = missing value for age in 14 individuals; * = missing data for 140 and 168 individuals before and after SCP, respectively