

Table SII. Sensitivity analyses: multivariate analyses with and without atopic dermatitis (AD) eyelid involvement in the past year

	Multivariate analyses with eyelid involvement (<i>n</i> = 193) ^a			Multivariate analyses without eyelid involvement (<i>n</i> = 193) ^a		
	OR	95% CI	<i>p</i> -value	OR	95% CI	<i>p</i> -value
Age at baseline (years)	1.01	0.89–1.14	0.924	0.99	0.89–1.12	0.921
Age at baseline (square)	1.00	1.00–1.00	0.960	1.00	1.00–1.00	0.950
Men	1.14	0.56–2.32	0.716	1.18	0.59–2.34	0.637
No history of any eye disease and no ophthalmic drugs at baseline, reference	–	–	–	–	–	–
History of any eye disease and no ophthalmic drugs at baseline	0.71	0.17–2.97	0.642	0.70	0.17–2.80	0.609
History of any eye disease and ophthalmic drugs at baseline	3.44	0.84–14.03	0.085	3.98	1.00–15–78	0.050
Self-reported episodic acute allergic conjunctivitis	1.59	0.64–3.97	0.317	1.86	0.75–4.60	0.178
Allergic asthma	0.72	0.35–1.46	0.361	0.79	0.40–1.58	0.509
Allergic rhinitis	0.78	0.31–1.98	0.595	0.85	0.34–2.13	0.731
Food allergy	1.11	0.55–2.25	0.774	1.07	0.53–2.14	0.854
EASI score at baseline, per severity category ^b	–	–	–	–	–	–
Mild and moderate AD (EASI 0.1–22.9) reference	–	–	–	–	–	–
Severe AD (EASI 23.0–72.0)	0.94	0.45–1.99	0.870	0.859	0.41–1.78	0.684
AD eyelid involvement in the past year	4.00	1.68–9.50	0.002	–	–	–

^aThe selected patients for this sensitivity analysis (*n*=193) were patients in which the presence or absence of AD eyelid involvement in the past year had been reported at baseline. As a result of this smaller sample size, the odds ratio (OR), 95% confidence intervals (CI), and *p*-values were not comparable with those of the main analyses.

^bSince there were only few patients with mild AD in which the presence or absence of AD eyelid involvement in the past year was known at baseline, patients with mild AD and moderate AD were combined into 1 category.

EASI: Eczema Area and Severity Index. Bold values indicate statistical significance.