

Supplementary material has been published as submitted. It has not been copyedited, typeset or checked for scientific content by Acta Dermato-Venereologica

Table SI. Characteristics of study population

Variable		N (%)
Age	Age at diagnosis (years±SD, cont.)	70.8 ± 14.5
Gender	Male	503 (50.3)
	Female	497 (49.7)
Educational level	No education	90 (9.0)
	Primary school	254 (25.4)
	Secondary school	301 (30.1)
	Higher education	355 (35.5)
Employment situation	Unemployed	95 (9.5)
	Active	222 (22.2)
	Retired	683 (68.3)
Marital Status	Single	114 (11.4)
	Married or part of a couple	599 (59.9)
	Widowed or divorced	287 (28.7)
Lives alone	Yes	281 (28.1)
	No	719 (71.9)
Skin phototype	I	87 (8.7)
	II	477 (47.7)
	III	333 (33.3)
	IV	93 (9.3)
	V	10 (1)
	VI	0 (0)
Type of tumor	Melanoma	83 (8.3)
	BCC	685 (68.5)
	SCC	232 (23.2)
Detection pattern	Patient	451 (45.1)
	Family or relatives	137 (13.7)
	Healthcare professionals	412 (41.2)

Long axis diameter	Long axis diameter (mm±SD, cont)	10.9 ± 8.9
Location	Face	459 (45.9)
	Scalp and neck	146 (14.6)
	Anterior trunk	82 (8.2)
	Back	142 (14.2)
	Upper extremities	91 (9.1)
	Lower extremities	80 (8.0)
Time of disease evolution	Unknown	221 (22.1)
	< 6 months	296 (29.6)
	6-12 months	234 (23.4)
	>12 months	248 (24.8)
Associated symptoms	Recent appearance	116 (11.6)
	Growth	227 (22.7)
	Bleeding	125 (12.5)
	Itch, local discomfort	196 (19.6)
	No	336 (33.6)
Personal history of skin cancer	Yes	566 (56.6)
	No	434 (43.4)
Family history of skin cancer	Yes	410 (41)
	No	590 (59)
Sunburn	Yes	565 (56.5)
	No	435 (43.5)
Immunosuppressed	Yes	100 (10)
	No	900 (90)
Total		1000 (100)

SD: Standard deviation, BCC: Basal cell carcinoma, Cont: continuous variable, SCC: Squamous cell carcinoma