

Table SI. Treatments received by patients before the switch to FDLPPS and after the diagnosis

Treatment	Treatments used during FD period, n (%) of patients	Treatments used 3 months before the switch to FDLPPS, n (%) of patients ¹	Treatments used during FDLPPS period, n (%) of patients
Topical treatments			
Steroids	21 (67.7)	9 (50.0)	25 (80.6)
Antibiotic	13 (41.9)	6 (33.3)	10 (32.3)
Dapsone	5 (16.1)	5 (27.8)	10 (32.3)
Minoxidil	1 (3.2)	-	3 (9.7)
Ciclopirox (shampoo)	3 (9.7)	3 (9.7)	7 (22.6)
Systemic treatments			
Isotretinoin	5 (16.1)	-	4 (12.9)
HXQ	2 (6.5)	-	5 (16.1)
Doxycycline	18 (58.1)	14 (77.8)	5 (16.1)
Azithromycin	-	-	1 (3.2)
Clindamycin	2 (6.5)	-	-
Clindamycin and Rifampicin	3 (9.7)	3 (16.7)	1 (3.2)
Itraconazol	-	-	2 (6.5)
DXM	-	-	1 (3.2)
Cyclosporine	-	-	1 (3.2)
Finasteride	-	-	1 (3.2)
Minoxidil	-	-	8 (25.8)
Intralesional steroids	8 (25.8)	7 (38.9)	20 (64.5)
TOTAL	31 (100)	18 (100)	31 (100)

HXQ, Hydroxychloroquine; DXM, Dexamethasone.

¹A total of 18 patients out of the total switch to the FDLPPS during the follow-up in our department.