

Pemphigus Improving with Gluten-free Diet

Francesco Drago, Marco Cacciapuoti, Matteo Basso, Aurora Parodi and Alfredo Rebora

SEM Section of Dermatology, Viale Benedetto XV, 7, IT-16132 Genoa, Italy. E-mail: fdrago@libero.it

Accepted June 28, 2004.

Sir,

Serum IgA class anti-endothelial antibodies (EMA) are markers of coeliac disease and dermatitis herpetiformis (DH) and are useful in monitoring compliance to diet. Their overall sensitivity for the diagnosis of DH is about 80% and their specificity around 96% (1). As anti-gliadin antibodies (AGA) have already been reported in pemphigus (2, 3), the presence of AGA in another patient with pemphigus may not surprise, but the finding of IgA-EMA in yet another, and the response of both patients to a gluten-free diet treatment are worthy of attention.

CASE REPORTS

Patient 1

An 82-year-old woman presented with a 20-day history of a bullous eruption. She had not been taking any drug and had previously been in good health. She exhibited vesiculobullous lesions on the trunk and eczematous lesions on the thighs. There were no mucosal or scalp lesions.

Histopathology showed spongiosis and lymphocytic exocytosis. Oedema and a modest inflammatory infiltrate were present in the superficial dermis. The infiltrate was perivascular and interstitial, with prevalence of eosinophils.

Direct immunofluorescence disclosed a pemphigus-like deposition of IgA in the epidermal intercellular space (ICS). No other immunoglobulins or complement components were deposited.

Circulating anti-ICS IgA was also demonstrated by indirect immunofluorescence on monkey oesophagus (Alfabiotech, Rome, Italy) at a titre of 1:40 together with IgA-EMA (+). ELISA showed circulating IgG-AGA (++) and IgA-AGA (+). The liquid from bullae did not contain anti-ICS IgA, but it was strongly positive for IgA-EMA. Anti-transglutaminase antibodies were absent in the serum. Immunoblotting performed on keratinocyte extracts showed circulating antibodies directed to an aspecific band at 140-kDa, both with anti-total Ig and anti-IgA.

Full blood count, erythrocyte sedimentation rate, liver function tests, electrolytes and total IgE were within

normal limits. No monoclonal IgA was found by immunoelectrophoresis.

Even in the absence of DH features, either in histology or in direct immunofluorescence, IgA-EMA, IgG-AGA and IgA-AGA prompted us to introduce a gluten-free diet. Twenty days later the lesions had cleared. Interruption of the diet caused clinical lesions to relapse promptly.

Patient 2

An 18-year-old woman presented with a 1-month history of vesiculobullous eruption on the chest. On examination, the bullae were tense and arose on erythematous skin. Gradually, they extended centrifugally from the breast to the abdomen, neck and lumbar regions, and then to the limbs and arms. No other abnormality was observed.

Full blood count, erythrocyte sedimentation rate, liver function tests, electrolytes, antinuclear antibodies, serum levels of immunoglobulins and complement components were within normal limits.

Histological examination of lesional skin showed a sub-corneal bullous lesion. There was acantholysis and spongiosis with eosinophilic exocytosis. Oedema and a perivascular infiltrate of eosinophils and lymphocytes were seen in the upper dermis.

Direct immunofluorescence of the lesional skin revealed a deposition of IgG and C3 in ICS. Indirect immunofluorescence on monkey oesophagus revealed circulating IgG anti-ICS (1:160). ELISA showed IgG-AGA (+++) and IgA-AGA (++) in the serum. Circulating IgA-EMA and anti-transglutaminase antibodies were absent. Serum immunoblotting on keratinocyte extracts reacted at 160-kDa antigen, typical of pemphigus foliaceus. ELISA confirmed the presence of serum antibodies directed to the 160-kDa antigen.

The presence of IgG-AGA prompted us to introduce a gluten-free diet, even in the absence of any clinical, histological and further immunological evidence of coeliac disease. One month later, lesions had cleared and the IgG-AGA titre abated, in spite of IgG anti-ICS having increased up to 1:160.

DISCUSSION

The various test results of the two patients are shown in Table I. The patients had IgA pemphigus and

Table I. Summary of immunoserological results

Patient no.	DIF	Circulating antibodies				Blister fluid
		IIF	ELISA	IB	TG Ab	IIF
1	IgA in ICS	IgA anti-ICS 1:40 IgA-EMA +	IgG-AGA ++ IgA-AGA +	140 kDa	Negative	IgA-EMA+++
2	IgG and C ₃ in ICS	IgG anti-ICS 1:160 IgA-EMA	IgG-AGA +++ IgA-AGA ++	160 kDa	Negative	ND

DIF, direct immunofluorescence; IIF, indirect immunofluorescence; IB, immunoblotting; Ab, antibody; ICS, intercellular substance; AGA, anti-gliadin antibody; EMA, anti-endothelial antibody; ND, not done; TG, transglutaminase.

pemphigus foliaceus, respectively. This was confirmed by direct immunofluorescence. Both shared two interesting features, the presence of serological markers of glutensensitive enteropathy and the impressive response to a gluten-free diet.

IgA-AGA and/or IgG-AGA mark coeliac disease and DH, but are also encountered in other disorders including psoriasis (4, 5), SLE (6) and pemphigus/pemphigoid (2). In fact, their prevalence in the population at large is as high as 5.7% (7).

IgA-EMA has a higher degree of specificity for DH and coeliac disease, although their prevalence in the population at large is 1.2% (7) and their sensitivity may be disappointing (8). IgA-EMA have never been reported in patients with pemphigus or pemphigoid (9).

Coeliac disease has an extremely polymorphic nature. Atypical or subclinical cases affecting several organs, without gastrointestinal symptoms, must be common, especially as cases with normal mucosal architecture of the bowel exist (8). Unfortunately, an accurate estimation of the incidence of these cases is lacking. In Italy, a recent survey disclosed that almost half of the diagnosed patients had a subclinical/silent form. The most frequent extra-intestinal marker of the subclinical cases were iron-deficiency anaemia (28%), alopecia and DH (11%), osteoporosis (7%) and recurrent aphthous stomatitis (6%), while first-degree relatives (30%), Basedow's disease (25%) and insulin-dependent diabetes (20%) were the most frequent associations in silent forms (10).

In our patients, the presence of both markers of coeliac disease suggests an underlying subclinical coeliac disease without diarrhoea, steatorrhoea, anaemia or low weight. The response to the gluten-free diet and the relapse that followed its withdrawal both strongly support the proposition that the patients had a silent gluten sensitivity. Unfortunately, both the patients refused duodenal biopsies.

Examples of a successful gluten-free diet in other cutaneous diseases exist, however. Psoriasis patients with high levels of IgA- and IgG-AGA improved with such a regimen even in the absence of IgA-EMA and of high numbers of intra-epithelial lymphocytes in duodenal mucosa (11). Alopecia areata occurs in 1.6% of patients with DH (12), but the effect of the diet is controversial (13–15). Although dapsone is regarded as the first-line therapy in IgA pemphigus (16) (possibly just because of a concomitant subclinical coeliac disease), a gluten-free diet has never been used before to treat pemphigus.

REFERENCES

- Peters MS, McEvoy MT. IgA antiendomysial antibodies in dermatitis herpetiformis. *J Am Acad Dermatol* 1989; 21: 1225–1231.
- Kumar V, Zane H, Kaul N. Serologic markers of glutensensitive enteropathy in bullous diseases. *Arch Dermatol* 1992; 128: 1474–1478.
- Economidou J, Avgerinou G, Tsiroyianni A, Stavropoulos P, Varelzidis A, Katsambas A. Endomysium and anti gliadin antibodies in dermatitis herpetiformis and other bullous diseases. *J Eur Acad Dermatol Venereol* 2000; 11: 184–185.
- Kamaeva OI, Reznikov IP, Pimenova NS, Dobritsyna LV. Anti gliadin antibodies in the absence of celiac disease. *Klin Med (Mosk)* 2000; 76: 33–35.
- Michaëlsson G, Gerden B. How common is gluten intolerance among patients with psoriasis? *Acta Derm Venereol* 1991; 71: 90.
- Rensch MJ, Szyjowski R, Shaffer RT, Finks S, Kopecky C, Grissmer L, et al. The prevalence of celiac disease autoantibodies in patients with systemic lupus erythematosus. *Am J Gastroenterol* 2001; 96: 1113–1115.
- McMillan SA, Watson RP, McCrum EE, Evans AE. Factors associated with serum antibodies to reticulin, endomysium, gliadin in an adult population. *Gut* 1996; 39: 43–47.
- Rostami K, Kerckhaert J, Tiemessen R, von Blomberg BM, Meijer JW, Mulder CJ. Sensitivity of antiendomysium anti gliadin antibodies in untreated celiac disease: disappointing in clinical practice. *Am J Gastroenterol* 1999; 94: 888–894.
- Kumar V, Hemedinger E, Chorzelski TP, Beutner EH, Valeski JE, Kowalewski C. Reticulin and endomysial antibodies in bullous diseases. Comparison of specificity and sensitivity. *Arch Dermatol* 1987; 123: 1179–1182.
- Tursi A, Giorgetti G, Brandimarte G, Rubino E, Lombardi D, Gasbarrini G. Prevalence and clinical presentation of subclinical/silent celiac disease in adults: an analysis on a 12-year observation. *Hepatogastroenterology* 2001; 48: 462–464.
- Michaëlsson G, Gerdén B, Hagforsen E, Nilsson B, Pihl-Lundin I, Kraaz W, et al. Psoriasis patients with antibodies to gliadin can be improved by a gluten-free diet. *Br J Dermatol* 2000; 142: 44–51.
- Reunala T, Collin P. Diseases associated with dermatitis herpetiformis. *Br J Dermatol* 1997; 136: 315–318.
- Corazza GR, Andreani ML, Ventura N, Bernardi M, Tosti A, Gasbarrini G. Celiac disease and alopecia areata: report of a new association. *Gastroenterology* 1995; 109: 1333–1337.
- Bondavalli P, Quadri G, Parodi A, Rebora A. Failure of gluten-free diet in celiac disease-associated alopecia areata. *Acta Derm Venereol* 2000; 78: 319.
- Bardella MT, Marino R, Barbareschi M, Bianchi F, Faglia G, Bianchi P. Alopecia areata and coeliac disease: no effect of a gluten-free diet on hair growth. *Dermatology* 2000; 200: 108–110.
- Jeffes EW 3rd, Ahmed AR. Therapy of pemphigus. *Clin Dermatol* 1983; 1: 122–140.