

Birt-Hogg-Dubé Syndrome: Germline Mutation in the (C)₈ Mononucleotide Tract of the *BHD* Gene in a German Patient

Christof Lamberti¹, Nadine Schweiger², Wolfgang Hartschuh³, Tilman Schulz⁴, Petra Becker-Wegerich², Wolfgang Küster⁵, Arno Rütten⁶, Tilman Sauerbruch¹, Thomas Ruzicka² and Roland Kruse²

¹Department of Internal Medicine I, University Hospital Bonn, Bonn, ²Department of Dermatology, University Hospital Duesseldorf, Moorenstrasse 5, DE-4074 Duesseldorf, ³Department of Dermatology, University Hospital Heidelberg, Heidelberg, ⁴Institute of Pathology, Ansbach, ⁵Department of Dermatology, TOMESA Clinics, Bad Salzschlirf, Bad Salzschlirf and ⁶Laboratory of Dermatohistopathology, Friedrichshafen, Germany. E-mail: rkruse@uni-duesseldorf.de
Accepted August 16, 2004.

Sir,

Birt-Hogg-Dubé syndrome (BHDS) (also known as Hornstein-Knickenberg syndrome in the German literature) is an autosomal dominant inherited skin disease with coexistence of adnexal skin tumours, namely multiple fibrofolliculomas/trichodiscomas and several internal disorders, e.g. renal and colonic neoplasms, and lung cysts. Various other associations may arise incidentally (1–6). The fibrofolliculomas/trichodiscomas are disseminated symmetrically in the face, although neck, shoulder and the trunk can additionally be affected. There is no significant phenotypic overlap of

BHDS with other cancer-prone genodermatoses, e.g. Muir-Torre syndrome, Gardner syndrome or Cowden syndrome. Recently, the responsible *BHD* gene was mapped to chromosome 17p11.2 and was identified to encode a 64 kD protein, referred to as folliculin, of so far unknown function (7, 8). Two studies reported germline mutations in the *BHD* gene and suspected a mutational hot spot in a (C)₈ mononucleotide tract in exon 11, as frameshift mutations in this mononucleotide tract were found in about 44% of the unrelated patients with BHDS (8, 9). Other germline mutations – located in exons 7, 9 and 12 – were reported only in single BHDS cases (8).




Fig. 1. Germline mutation 1733insC in the (C)₈ mononucleotide tract of exon 11 of the *BHD* gene which is predicted to truncate the protein (family 2). The figure shows the antisense strand of the DNA sequence containing the altered poly C tract.

To check the presumed mutational hot spot, we evaluated the (C)₈ mononucleotide tract of the *BHD* gene for germline mutations in three unrelated German patients with clinically proven BHDS. The phenotypes of all three index patients met the definition of BHDS, i.e. the occurrence of multiple fibrofolliculomas/trichodiscomas. Clinical characteristics of these three patients and their affected relatives were published recently (2, 5, 10). In all three families the index patients and at least one first degree relative were affected with histologically verified fibrofolliculomas. In addition, colorectal adenomas were found in two index patients and two affected parents died as a consequence of colorectal cancer. For mutation analysis of the (C)₈ mononucleotide tract in exon 11 of the *BHD* gene the PCR conditions and primer sequences (SKA7: 5'GGTTCCACTTTGGGCC TGAG; SKA8: 5'GGTAGTAGAGCATGGATGGCC) described by Nickerson et al. (8) were used. Sequencing was performed on an ABI 377 DNA Sequencer (Applied Biosystems). Whereas neither of the index patients in references 2 and 10 showed a frameshift mutation in the mononucleotide tract, in the third patient (5) the frameshift mutation 1733insC could be identified (Fig. 1). This insertion is predicted to truncate the protein 26 missense amino acids downstream.

Our findings support the importance of the (C)₈ mononucleotide tract in exon 11 of the *BHD* gene in the aetiology of BHDS, as the majority of *BHD* germline mutations were C insertion or deletion mutations in this mononucleotide tract (8, 9, authors' own data). The high prevalence of these germline mutations raises the possibility of establishing a simple, rapid and cost-effective molecular screening test for patients suspected of BHDS.

Furthermore, in accordance with more recent studies performed by Schulz & Hartschuh (5) we found an increased frequency of extracutaneous manifestations, especially colonic neoplasms, in our families with BHDS. This observation points out the necessity of

screening these patients at regular intervals by colonoscopy to prevent the development of colorectal cancer. Conversely, all patients affected with colorectal adenomas and carcinomas should be carefully examined for cutaneous manifestations and asked for their family tumour history.

REFERENCES

1. Rongioletti F, Hazini R, Gianotti G, Rebora A. Fibrofolliculomas, trichodiscomas and acrochordons (Birt-Hogg-Dubé) associated with intestinal polyposis. *Clin Exp Dermatol* 1989; 14: 72–74.
2. Schachtschabel AA, Küster W, Happle R. [Perifollicular fibroma of the skin and colonic polyps: Hornstein-Knickenberg syndrome]. *Hautarzt* 1996; 47: 304–306.
3. Toro JR, Glenn G, Duray P, Darling T, Weirich G, Zbar B, et al. Birt-Hogg-Dubé syndrome: a novel marker of kidney neoplasia. *Arch Dermatol* 1999; 135: 1195–1202.
4. Walter P, Kirchhof B, Korge B, Heimann K. Flecked chorioretinopathy associated with Birt-Hogg-Dubé syndrome. *Graefes Arch Clin Exp Ophthalmol* 1997; 235: 359–361.
5. Schulz T, Hartschuh W. Birt-Hogg-Dubé syndrome and Hornstein-Knickenberg syndrome are the same. Different sectioning technique as the cause of different histology. *J Cutan Pathol* 1999; 26: 55–61.
6. Schulz T, Hartschuh W. Characteristics of the Birt-Hogg-Dubé/Hornstein-Knickenberg syndrome. *Am J Dermatopathol* 2000; 22: 293–294.
7. Schmidt LS, Warren MB, Nickerson ML, Weirich G, Matrosova V, Toro JR. Birt-Hogg-Dubé syndrome, a genodermatosis associated with spontaneous pneumothorax and kidney neoplasia, maps to chromosome 17p11.2. *Am J Hum Genet* 2001; 69: 876–882.
8. Nickerson ML, Warren MB, Toro JR, Matrosova V, Glenn G, Turner ML, et al. Mutations in a novel gene lead to kidney tumors, lung wall defects, and benign tumors of the hair follicle in patients with the Birt-Hogg-Dubé syndrome. *Cancer Cell* 2002; 2: 157–164.
9. Khoo SK, Giraud S, Kahnoski K, Chen J, Motorna O, Nickolov R, et al. Clinical and genetic studies of Birt-Hogg-Dubé syndrome. *J Med Genet* 2002; 39: 906–912.
10. Becker-Wegerich P, Kruse R, Bruch-Gerharz D, Fritsch C, Assmann T, Ruzicka T. Hornstein-Knickenberg-Syndrom. *Z Hautkr* 2002; 77: 199–215.