

Table SII. Associations between localization and other parameters

	Nose <i>n</i> (%)	Ear <i>n</i> (%)	Periorbital <i>n</i> (%)	Scalp <i>n</i> (%)	Cheek, forehead, perioral and temple <i>n</i> (%)	Trunk (including neck) <i>n</i> (%)	Upper extremities <i>n</i> (%)	Lower extremities <i>n</i> (%)	<i>p</i> -value
Physician specialty									
General surgeon	0 (0)	0 (0)	0 (0)	0 (0)	2 (3.6)	36 (64.3)	10 (17.9)	8 (14.3)	<0.0001
General practitioner	0 (0)	1 (1.4)	1 (1.4)	4 (5.6)	8 (11.1)	40 (55.6)	10 (13.9)	8 (11.1)	
Dermatologist	13 (5.1)	4 (1.6)	4 (1.6)	2 (0.8)	153 (60.5)	48 (19.0)	16 (6.3)	13 (5.1)	
Plastic surgeon	65 (23.0)	23 (8.2)	14 (5.0)	35 (12.4)	101 (35.8)	26 (9.2)	5 (1.8)	13 (4.6)	
Otorhinolaryngologist	38 (12.7)	44 (14.7)	7 (2.3)	9 (3.0)	187 (62.5)	14 (4.7)	0 (0)	0 (0)	
Other	0 (0)	0 (0)	7 (58.3)	0 (0)	0 (0)	1 (8.3)	4 (33.3)	0 (0)	
Physician experience									
Specialist	71 (11.4)	49 (7.8)	15 (2.4)	26 (4.2)	309 (49.4)	103 (16.5)	25 (4.0)	27 (4.3)	0.099
Resident	17 (12.7)	6 (4.5)	4 (3.0)	8 (6.0)	51 (38.1)	30 (22.4)	9 (6.7)	9 (6.7)	

Significant *p*-values in bold.