

## CLINICAL REPORT

# Nickel Sensitization in Adolescents and Association with Ear Piercing, Use of Dental Braces and Hand Eczema.

*The Odense Adolescence Cohort Study on Atopic Diseases and Dermatitis (TOACS)*

CHARLOTTE GOTTHARD MORTZ, JENS MARTIN LAURITSEN, CARSTEN BINDSLEV-JENSEN and KLAUS EJNER ANDERSEN

*Department of Dermatology, Odense University Hospital, Odense, Denmark*

**The prevalence of nickel allergy (sensitization) and the associations with ear piercing, use of dental braces and hand eczema were assessed in a cohort of 1,501 8th grade schoolchildren (aged 12–16 years) in Odense, Denmark. Nickel allergy was found in 8.6% and was clinically relevant in 69% of cases. Nickel allergy was found most frequently in girls and the association with ear piercing was confirmed. Application of dental braces (oral nickel exposure) prior to ear piercing (cutaneous nickel exposure) was associated with a significantly reduced prevalence of nickel allergy. In adolescents a significant association was found between hand eczema and nickel allergy. A follow-up study of this population is planned in order to assess the course and development of contact dermatitis, hand eczema and atopic diseases in adulthood and after choice of occupation. Key words: schoolchildren; atopic dermatitis; inhalant allergy; hand eczema; multivariate graphical analysis.**

(Accepted May 8, 2002.)

Acta Derm Venereol 2002; 82: 359–364.

Charlotte Gotthard Mørtz, Department of Dermatology, Odense University Hospital, DK-5000 Odense C, Denmark. E-mail: mortz@imbmed.ou.dk

Studies in children and adolescents visiting dermatological clinics as well as population-based studies have repeatedly shown that nickel allergy is the most common contact allergy (1–13). The prevalence of nickel sensitization in children and adolescents in the general population varies from 0.9 to 14.9% (2, 14, 15). The most common cause of sensitization appears to be ear piercing (16–19), and the risk for nickel allergy is increased with the number of piercings (16, 17). Other sources of sensitization include jewellery, belt buckles, metal fasteners on clothing and eyeglass frames.

Few reports have suggested that treatment with nickel-containing metallic orthodontic appliances (oral nickel exposure) before ear piercing (cutaneous nickel exposure) may reduce the frequency of nickel allergy (20, 21). However, more clinical studies are needed to confirm this hypothesis.

Nickel allergy in adults has been shown to increase

the risk of development of hand eczema (22). However it is not known whether this relationship exists already in childhood and adolescence.

The aims of this study were 1) to estimate prevalence measures of nickel sensitization in an unselected population of adolescents, 2) to identify groups of nickel sensitized persons with a weak, moderate or strong degree of nickel allergy in order to evaluate in future studies if the degree of nickel allergy is correlated with later development of dermatitis, 3) to confirm the relationship between nickel allergy and ear piercing and the use of dental braces, and 4) to study the relationship between nickel allergy and atopic diseases and nickel allergy and hand eczema.

## METHODS

### *Population and study design*

The Odense Adolescence Cohort Study (TOACS) is an epidemiological follow-up study. Phase one (1995–1996) was conducted as a cross-sectional study among 1,501 8th grade schoolchildren in 40 schools in the Municipality of Odense. Phase one included questionnaires, interviews and clinical examinations, blood samples for IgE measurement and patch tests. Phase two (1996–1997) was conducted as a case-control study in selected groups of schoolchildren. The population and study design has been described already (23).

### *Definitions and description of terms*

*Atopic dermatitis, inhalant allergy and allergic contact dermatitis.* The criteria for the diagnoses have already been reported in detail (23, 24).

*Hand eczema.* The lifetime prevalence of hand eczema was evaluated from the questionnaire, as already reported (23, 24). At the clinical examination, hand eczema was defined as eczema localized to the fingers or finger webs, backs of hands or palms, characterized by itching, erythema, vesicles and/or papules and scaling. More chronic erythematous, scaling, fissuring and/or lichenified types of dermatitis were also included. The eczema should have a duration of at least 2 days.

*Allergic contact dermatitis caused by metal contact.* The lifetime prevalence of self-reported eczema caused by metal skin contact was evaluated from the question: “Do you get eczema (rash) from jeans buttons, metal fasteners, metal costume jewellery (e.g. earrings) or other metal parts of clothes next to your skin (excluding under the ring)?” (25).

*Ear piercing and use of dental braces.* The questions were: (i)

Have you ever had your ears pierced or had any other piercing of your skin?", (ii) "Have you ever had dental braces?" The time for both events should be given.

*Contact allergy/Type IV sensitization.* This was defined as at least one positive patch test reaction to allergens in the TRUE Test® including a nickel dilution series.

#### Patch tests

The TRUE Test® panels 1 and 2 (Pharmacia & Upjohn, Hillerød, Denmark) was used together with a TRUE Test patch dilution series consisting of nickel sulphate (NiSO<sub>4</sub>, 6H<sub>2</sub>O) in 3 concentrations (200, 10, 1 µg/cm<sup>2</sup>) and one placebo (26, 27). The patch test method has been described previously (24). Briefly, the patch tests were applied to the upper back for 2 days, removed by the investigators and scored after 3 days (28). The relevance of a positive test result was evaluated in relation to exposure history, dermatitis history and present dermatitis pattern.

#### Ethics

The ethics committee for Vejle and Funen County (proj. no. 95/22) approved the study. Informed consent was obtained both from the school children and from their parents.

#### Data handling and statistics

All data were entered twice in the databases. When differences were found, a comparison with raw input forms was made and corrections done accordingly. Statistical analyses were performed using Stata 5.0 (Stata corporation, TX, USA), with the exception of in graphical models.

The prevalence proportion was defined by the number of positive answers divided by the total number of schoolchildren questioned. The 95% confidence intervals are shown in parentheses (95% CI). The prevalence proportions for boys and girls are given and if a significant sex difference is found, the *p*-value is given or significance is indicated. Comparisons were made by  $\chi^2$ -based table analysis.

Odds ratio (OR) is given as the Mantel Haenzel odds ratio stratified by sex, with associated confidence intervals in parentheses (95% CI). Differences by sex are noted, when the stratum-specific estimates indicate significant "effect modification". Statistical significance was defined as *p* < 0.05.

Because of the close association between the investigated diseases, a multivariate analysis was performed, which at the same time could account for the interdependence and possible association with external factors (control for all associations at the same time). The analysis was performed using specialized software (Digram®) following the procedure as described by Klein et al. (29). The results are expressed in the form of a

graph on which non-random associations between variables are represented by a line (further description in (24)). Because the variables are binary or ordinal, the strength and degree of statistical significance of an association can be measured by Kruskal and Goodman's gamma coefficient in the form of a conditional or partial gamma (30). Gamma coefficients numerically less than 0.15 indicate weak associations, those between 0.15 and 0.30 indicate moderate associations, and more than 0.30 strong associations. Because of the many statistical tests in these analyses, a significance level of 0.01 was used to compensate for false associations (Type I error).

## RESULTS

### Prevalence of nickel sensitization

The pattern of participation, patch test characteristics and frequencies of individual patch test reactions have been described previously (23, 24). Patch testing was performed in 76.3% (1,146/1,501) of the schoolchildren.

Nickel sensitization was found in 8.6% using results from both patches with nickel sulphate 200 µg/cm<sup>2</sup> (Table I). In the TRUE Test patch dilution series 2.8% reacted to nickel sulphate 10 µg/cm<sup>2</sup> and 0.3% to nickel sulphate 1 µg/cm<sup>2</sup>. Significantly more girls than boys had nickel allergy (Table I). No reactions to nickel sulphate 10 µg/cm<sup>2</sup> and 1 µg/cm<sup>2</sup> were obtained without a reaction to 200 µg/cm<sup>2</sup> and no reactions to the placebo were seen.

### Nickel allergy in relation to atopic dermatitis, inhalant allergy and hand eczema

Nickel allergy was significantly associated with hand eczema (OR 2.36, 95% CI 1.39–4.01, *p* < 0.002 stratified for sex). The association was still significant when those with concomitant atopic dermatitis and inhalant allergy were excluded from the hand eczema group (OR 4.01, 95% CI 1.82–8.84, *p* < 0.001 stratified for sex).

Of those with present or past hand eczema, 18.2% had nickel allergy (girls 23.5%, boys 7.5%, *p* < 0.04 for sex difference) compared with 7.4% of those without hand eczema (girls 12.2%, boys 2.1%, *p* < 0.001 for sex difference). On the other hand, 22.4% of nickel allergic schoolchildren had present or past hand eczema (girls 22.4%, boys 23.1%) compared with only 9.4% of those without nickel allergy (girls 11.6%, boys 7.2%, *p* < 0.02

Table I. Prevalence of nickel sensitization, ear piercing, use of dental brace and hand eczema (lifetime prevalence measures)

	Total population	Girls	Boys
Nickel sensitization			
200 µg/cm <sup>2</sup>	8.6% (98/1,146)	13.7% (85/620)*	2.5% (13/526)
10 µg/cm <sup>2</sup>	2.8% (32/1,146)	5.0% (31/620)*	0.2% (1/526)
1 µg/cm <sup>2</sup>	0.3% (4/1,146)	0.6% (4/620)	0% (0/526)
Ear piercing	50.8% (731/1,438)	81.5% (581/713)*	20.7% (150/725)
Dental brace	31.7% (456/1,438)	39.3% (280/713)*	24.3% (176/725)
Hand eczema	9.2% (133/1,438)	12.2% (87/713)*	6.3% (46/725)

\**p* < 0.001 for sex difference.

for sex difference). Furthermore, we also found an association between clinically relevant nickel allergy and hand eczema by analysing self-reported metal-contact-related eczema verified by a positive patch test to nickel (OR 2.39, 95% CI 1.27–4.50,  $p < 0.006$ , stratified for sex). There was no association between contact allergy to other compounds than nickel and hand eczema.

Among the adolescents, 1.6% (13 girls, 9 boys) had current hand eczema. Twenty-one of the 22 were patch-tested and 2 girls, both with pierced ears, had positive reactions to nickel. One of the girls had nickel allergy of current relevance and a strong reactivity down to  $10 \mu\text{g}/\text{cm}^2$ . The other girl had a reaction only to  $200 \mu\text{g}/\text{cm}^2$  and had no history of dermatitis from metal skin contact.

There were no significant associations between nickel allergy and atopic dermatitis or inhalant allergy (data not shown).

A multivariate analysis with nickel sensitization as the outcome is illustrated in Fig. 1 and Table II. All those with other contact allergies were excluded from the analysis. The 3 different boxes in Fig. 1 indicate outcome variable (nickel sensitization), investigated diseases and background variables. The figure should be read from the right to the left. Arrows indicate associations between different levels (boxes) and lines indicate associations within the same level (box). As expected, the analysis shows a strong association between atopic dermatitis and inhalant allergy, between atopic dermatitis and hand eczema, and also an association between nickel allergy and hand eczema. However, conflicting gamma and  $\chi^2$  values were found for the relationship between nickel allergy and hand eczema




Fig. 1. Summary of results from a multivariate graphical analysis with nickel sensitization as outcome ( $n = 1,070$ ). In this analysis all those with other contact allergies have been excluded. See also Table II. Ni: Nickel sensitization – positive patch test for nickel; AD: present or past atopic dermatitis; IA: present or past inhalation allergy (allergic asthma and/or allergic rhinitis); HE: present or past hand eczema; Eczema: present or past eczema caused by metal contact; Piercing: present or past ear piercing.

(gamma 0.36 vs.  $\chi^2$  7.7). This analysis shows no association between nickel allergy and atopic dermatitis or inhalant allergy.

In a multivariate analysis for contact allergy excluding nickel sensitization, no association was found between contact allergy and hand eczema, atopic dermatitis or inhalant allergy (data not shown).

#### Nickel allergy by patch testing in relation to clinical relevance

The association between reported eczema caused by metal skin contact and nickel allergy by patch testing adjusted for sex was highly significant (OR 9.33, 95% CI 5.62–15.47,  $p < 0.001$ ). In the questionnaire a total of 15.5% (223/1,438) of schoolchildren reported eczema caused by metal skin contact (girls 23.4%, boys 7.7%,  $p < 0.001$  for sex difference), and among those participating in patch testing 17.9% (205/1,146) reported metal-contact-related eczema (girls 24.8%, boys 9.7%,  $p < 0.001$  for sex difference). Of those with a history of eczema from metal skin contact, 30.7% (girls 37.7%, boys 9.8%,  $p < 0.001$  for sex difference) showed a positive nickel patch test, compared with 3.7% (girls 5.8%, boys 1.7%,  $p < 0.002$  for sex difference) in those with a negative history of eczema from metal skin contact.

The clinical relevance of a positive patch test to nickel sulphate was assessed by exposure history, dermatitis history and present dermatitis pattern immediately after patch testing. 69.4% (68/98) of the nickel allergic cases had a clinical relevance (girls 74.1%, boys 38.5%,  $p < 0.01$  for sex difference) and only 5 of these 68 cases were not identified from the questionnaire.

The clinical relevance of nickel allergy was found in 87.5% of those with reactions to nickel sulphate  $10 \mu\text{g}/\text{cm}^2$  and in all 4 reacting to nickel sulphate  $1 \mu\text{g}/\text{cm}^2$ .

#### Nickel allergy in relation to ear piercing and wearing of dental braces

In this study 50.8% of the adolescents had their ears pierced, and 31.7% had used dental braces (Table I).

As expected, nickel allergy was significantly associated with ear piercing adjusted for sex (OR 3.18, 95% CI 1.43–7.08,  $p < 0.004$ ). However, there was a significant effect modification by sex. A significant association was found between nickel allergy and ear piercing in girls (OR 5.12, 95% CI 1.82–14.44,  $p < 0.002$ ). Of the girls with pierced ears, 15.9% had nickel allergy compared with 3.6% of girls without ear piercing. Among girls with nickel allergy, 95.3% had ear piercing, while 79.8% of the girls without nickel allergy had ear piercing. In boys, nickel allergy was not associated with ear piercing (OR 0.88, 95% CI 0.24–3.23,  $p = 0.841$ ).

The occurrence of nickel allergy in relation to different time sequence combinations of use of dental braces and ear piercing is presented in Table III. None of the boys

Table II. Summary of results from the multivariate graphical analysis with nickel sensitization as outcome ( $n=1070$ ). In this analysis all those with other contact allergies have been excluded. See also Fig. 1

Hypothesis	$\chi^2$	Df	Exact $p$ -value	Gamma	Exact $p$ -value (one-sided)
Ni vs. HE $\perp$ sex, eczema	7.7	4	0.094	0.36	0.006
Ni vs. sex $\perp$ HE, eczema	23.5	4	0.000	-0.61	0.000
Ni vs. eczema $\perp$ HE, sex	103.2	4	0.002	0.81	0.002
AD vs. IA	94.3	1	0.000	0.62	0.000
AD vs. HE $\perp$ sex, eczema	81.5	4	0.000	0.68	0.000
AD vs. sex $\perp$ HE, eczema	8.5	4	0.068	-0.18	0.010
AD vs. eczema $\perp$ HE, sex	12.5	4	0.018	0.27	0.002
HE vs. sex $\perp$ AD, eczema, piercing	15.8	8	0.042	-0.26	0.060
HE vs. eczema $\perp$ AD, sex, piercing	27.6	8	0.004	0.41	0.002
HE vs. piercing $\perp$ sex, eczema	12.1	4	0.020	-0.19	0.082
Sex vs. eczema $\perp$ piercing	14.5	2	0.000	-0.38	0.000
Sex vs. piercing $\perp$ eczema	433.9	2	0.000	-0.87	0.000
Eczema vs. piercing $\perp$ sex	22.3	2	0.000	0.48	0.000

The Table contains the statistical- or user-specified non-random associations.

Ni vs. HE  $\perp$  sex, eczema, e.g. Ni is associated with hand eczema conditional ( $\perp$ ) on sex and eczema.

Conditional in this context means "sex and eczema", plus all other information in the analysis.

Ni: Nickel sensitization – positive patch test for nickel; AD: present or past atopic dermatitis; IA: present or past inhalation allergy (allergic asthma and/or allergic rhinitis); HE: present or past hand eczema; Eczema: present or past eczema caused by metal contact; Piercing: present or past ear piercing.

Table III. Relations between use of dental brace, ear piercing and nickel allergy

Dental brace	Pierced ears	Group size			Percentage of nickel patch test positive		
		Total ( $n=1,438$ )	Girls ( $n=713$ )	Boys ( $n=725$ )	Total ( $n=1,146$ )	Girls ( $n=620$ )	Boys ( $n=526$ )
+*	+	266	227	39	15.7% (36/230)	18.0% (36/200)	0% (0/30)
+ first	+	67	51	16	1.7% (1/60)	2.1% (1/48)	0% (0/12)
+	+ first	177	160	17	20.4% (31/152)	22.5% (31/138)	0% (0/14)
+ same year	+ same year	22	16	6	22.2% (4/18)	28.6% (4/14)	0% (0/4)
-	-	517	79	438	2.9% (11/382)	4.5% (3/66)	2.5% (8/316)
+	-	190	53	137	2.7% (4/150)	2.2% (1/46)	2.9% (3/104)
-	+	465	354	111	12.2% (47/384)	14.6% (45/308)	2.6% (2/76)

\*Those with both dental brace and ear piercing were separated into 3 groups, depending on whether the dental brace was the first event, the ear piercing was the first event or the two events occurred during the same year.

with both dental braces and ear piercing had nickel allergy, while 18.0% of the girls with both conditions had nickel allergy. The application of dental braces prior to ear piercing was significantly associated with a reduced frequency of nickel allergy in girls compared to the use of dental braces after ear piercing (OR 0.07, 95% CI 0.01–0.59,  $p < 0.002$ ). Only 2.1% of the girls who acquired dental braces before their ears were pierced had nickel allergy, compared with 22.5% of the girls who had ear piercing before acquiring the dental braces.

Considering the question about self-reported metal-related eczema, 17.4% of the girls who had ear piercing prior to using dental braces reported eczema caused by metal skin contact and had a positive patch test to nickel, while none of the girls with dental braces before piercing reported metal-related eczema and had a positive patch test to nickel ( $p < 0.003$ ).

## DISCUSSION

The study confirmed that the prevalence of nickel allergy was high in adolescents (8.6%) in accordance with other population-based studies in this age group (0.9% to 14.9%) (2, 14, 15), and, as expected, nickel allergy was significantly more frequent in girls than boys (girls 13.7% vs. boys 2.5%) (5, 6, 11, 12, 31).

The European Union (EU) regulation on nickel exposure (94/27/EEC) is now in effect in 2001 whereas it was implemented in Denmark in 1991 (32). The reduction of nickel release from jewellery seems to reduce the frequency of nickel allergy in young people (33).

A possible relationship between the level of skin reactivity to nickel sulphate and the presence of hand dermatitis has not been studied in a prospective protocol. Two retrospective studies revealed no relationship (27,

34). Furthermore, an inverse correlation between skin reactivity to nickel and the effect of a low nickel diet on activity of dermatitis has been published (35). In our study, patch testing was performed with 3 different concentrations of nickel sulphate to identify groups of people with varying degrees of nickel sensitization. Reactions to nickel sulphate concentrations of  $10 \mu\text{g}/\text{cm}^2$  were seen in 2.8% and to  $1 \mu\text{g}/\text{cm}^2$  in 0.3%. The concentrations were selected from a previous dose-response study in selected nickel-sensitive patients (27). The 3 groups of nickel-sensitized persons will be followed prospectively to determine whether those with moderate or strong sensitization are more likely to develop hand eczema.

Nickel allergy was considered clinically relevant in 69.4% of the nickel sensitized, and those reacting to the 2 lower concentrations were almost all of clinical relevance. Recall bias could explain some of the reactions without a positive history. However, the frequency of relevance of a positive nickel patch test was higher in this study than in two other studies including schoolchildren (16, 17), and was in agreement with a Danish study among veterinary students (36).

Of those reporting metal-related eczema, only 30.7% had a positive nickel patch test, as also reported by Kieffer (36). Possible explanations for the difference could be irritant skin reactions under wristwatch cases, buckles and rings or false negative patch test reactions. It has also been suggested that a positive history but a negative nickel patch test indicates atopy (37).

The most common cause of sensitization to nickel appears to be ear piercing (16, 17) and the risk for nickel allergy is increased with the number of piercings (17). However, we could not confirm the association in men as reported by others (38).

Explanations for the more frequent occurrence of nickel allergy in girls compared with boys include the fact that more girls than boys have their ears pierced, and boys may have ear piercing at a later age, have piercing only in one ear, or they use jewellery containing less nickel (28). However, it has also been suggested that women are sensitized by a lower exposure than men (39).

There is no method of desensitization. However, some studies have shown that treatment with nickel-containing metallic orthodontic appliances before ear piercing (cutaneous nickel exposure) may reduce the frequency of nickel hypersensitivity (20, 21). Studies have verified that nickel-containing alloys undergo degradation and small quantities of free metals (ions) are released in the oral cavity (40). We also found that the use of dental braces before ear piercing was associated with a lower prevalence of nickel allergy compared to ear piercing before the use of dental braces. These results support the view that oral administration of a contact allergen prior to attempted sensitization induces immune tolerance.

Several studies in adults have shown an association between hand eczema and nickel allergy (22, 39, 41, 42). In a previous report we found a significant association between contact allergy and hand eczema in adolescents (24), and it was nickel allergy that was responsible for this association. Those with nickel allergy had a higher prevalence of hand eczema (22.4%) compared with those without nickel allergy (9.4%). The risk assessment cannot be evaluated in this cross-sectional design, but will be evaluated in a follow-up study in this population.

The association between nickel allergy and atopic dermatitis, inhalant allergy and hand eczema was evaluated in a simple model using the Mantel Haenzel analysis, stratified for sex. Because of the association between the 3 diseases, analyses have also been performed for each disease alone, excluding schoolchildren with one or both of the other diseases. This is a simplification of real life, and more complicated statistical approaches were performed for verification of the results. Because of the association between all three diseases, a logistical regression model was not considered appropriate and a multivariate graphical analysis was used as an alternative. This gave similar results, as the simpler stratified analyses, for example, verified the association between nickel allergy and hand eczema.

In the future this cohort of schoolchildren will be followed with regard to the course and development of atopic diseases, hand eczema and contact dermatitis.

#### ACKNOWLEDGEMENT

We thank the schoolchildren, their parents, the schools involved and Odense Education Authority for their cooperation, and nurse Annemarie Yde-Andersen for skilful technical help. This study was supported by The Faculty of Health Sciences, University of Southern Denmark – Odense University, the Danish Asthma and Allergy Association, the Gerda and Aage Haensch Foundation and the Ingemann O. Buck Foundation. Pharmacia & Upjohn donated the TRUE Test® and TRUE Test patch dilution series with nickel sulphate.

#### REFERENCES

1. Mortz CG, Andersen KE. Allergic contact dermatitis in children and adolescents. *Contact Dermatitis* 1999; 41: 121–130.
2. Dotterud LK, Falk ES. Contact allergy in relation to hand eczema and atopic diseases in north Norwegian schoolchildren. *Acta Paediatr* 1995; 84: 402–406.
3. Ayala F, Balato N, Lembo G, Patruno C, Tosti A, Schena D, et al. A multicentre study of contact sensitization in children. Gruppo Italiano Ricerca Dermatiti da Contatto e Ambientali (GIRDCA). *Contact Dermatitis* 1992; 26: 307–310.
4. Balato N, Lembo G, Patruno C, Ayala F. Patch testing in children. *Contact Dermatitis* 1989; 20: 305–307.
5. Brasch J, Geier J. Patch test results in schoolchildren. *Contact Dermatitis* 1997; 37: 286–293.

6. Katsarou A, Koufou V, Armenaka M, Kalogeromitros D, Papanayotou G, Varelzidis A. Patch tests in children: a review of 14 years' experience. *Contact Dermatitis* 1996; 34: 70–71.
7. Pambor M, Kruger G, Winkler S. Results of patch testing in children. *Contact Dermatitis* 1992; 27: 326–328.
8. Pevny I, Brennenstuhl M, Razinskas G. Patch testing in children. (I) Collective test results; skin testability in children. *Contact Dermatitis* 1984; 11: 201–206.
9. Rademaker M, Forsyth A. Contact dermatitis in children. *Contact Dermatitis* 1989; 20: 104–107.
10. Romaguera C, Alomar A, Camarasa JM, Garcia BB, Garcia PA, Grimalt F, et al. Contact dermatitis in children. *Contact Dermatitis* 1985; 12: 283–284.
11. Rudzki E, Rebandel P. Contact dermatitis in children. *Contact Dermatitis* 1996; 34: 66–67.
12. Sevila A, Romaguera C, Vilaplana J, Botella R. Contact dermatitis in children. *Contact Dermatitis* 1994; 30: 292–294.
13. Veien NK, Hattel T, Justesen O, Norholm A. Contact dermatitis in children. *Contact Dermatitis* 1982; 8: 373–375.
14. Barros MA, Baptista A, Correia TM, Azevedo F. Patch testing in children: a study of 562 schoolchildren. *Contact Dermatitis* 1991; 25: 156–159.
15. Weston WL, Weston JA, Kinoshita J, Kloepfer S, Carreon L, Toth S, et al. Prevalence of positive epicutaneous tests among infants, children, and adolescents. *Pediatrics* 1986; 78: 1070–1074.
16. Dotterud LK, Falk ES. Metal allergy in north Norwegian schoolchildren and its relationship with ear piercing and atopy. *Contact Dermatitis* 1994; 31: 308–313.
17. Larsson-Stymne B, Widstrom L. Ear-piercing – a cause of nickel allergy in schoolgirls? *Contact Dermatitis* 1985; 13: 289–293.
18. Nielsen NH, Menne T. Nickel sensitization and ear piercing in an unselected Danish population. Glostrup Allergy Study. *Contact Dermatitis* 1993; 29: 16–21.
19. McDonagh AJ, Wright AL, Cork MJ, Gawkrödger DJ. Nickel sensitivity: the influence of ear piercing and atopy. *Br J Dermatol* 1992; 126: 16–18.
20. Van Hoogstraten I, Andersen KE, Von Blomberg B, Boden D, Bruynzeel DP, Burrows D, et al. Reduced frequency of nickel allergy upon oral nickel contact at an early age. *Clin Exp Immunol* 1991; 85: 441–445.
21. Kerosuo H, Kullaa A, Kerosuo E, Kanerva L, Hensten-Pettersen A. Nickel allergy in adolescents in relation to orthodontic treatment and piercing of ears. *Am J Orthod Dentofac Orthop* 1996; 109: 148–154.
22. Menne T, Borgan O, Green A. Nickel allergy and hand dermatitis in a stratified sample of the Danish female population: an epidemiological study including a statistic appendix. *Acta Derm Venereol* 1982; 62: 35–41.
23. Mortz CG, Lauritsen JM, Bindslev-Jensen C, Andersen KE. Prevalence of atopic dermatitis, asthma, allergic rhinitis, and hand and contact dermatitis in adolescents. The Odense Adolescence Cohort Study on Atopic Diseases and Dermatitis. *Br J Dermatol* 2001; 144: 523–532.
24. Mortz CG, Lauritsen JM, Bindslev-Jensen C, Andersen KE. Contact allergy and allergic contact dermatitis in adolescents: prevalence measures and associations. The Odense Adolescence Cohort Study on Atopic Diseases and Dermatitis. *Acta Derm Venereol* 2002; 82: 352–358.
25. Susitaival P, Kanerva L, Hannuksela M, Jolanki R, Estlander T. Tuohilampi questionnaire for epidemiological studies of contact dermatitis and atopy. *People and Work* 1996; 10: 1–26.
26. Fischer T, Maibach HI. The thin layer rapid use epicutaneous test (TRUE-test), a new patch test method with high accuracy. *Br J Dermatol* 1985; 112: 63–68.
27. Andersen KE, Liden C, Hansen J, Volund A. Dose-response testing with nickel sulphate using the TRUE test in nickel-sensitive individuals. Multiple nickel sulphate patch-test reactions do not cause an “angry back”. *Br J Dermatol* 1993; 129: 50–56.
28. Rycroft RJG, Menne T, Frosch PJ. *Textbook of contact dermatitis*. 2nd ed. Berlin, Heidelberg: Springer-Verlag, 1995.
29. Klein JP, Keiding N, Kreiner S. Graphical models for panel studies, illustrated on data from the Framingham Heart Study. *Stat Med* 1995; 14: 1265–1290.
30. Agresti A. *Analysis of ordinal categorical data*. New York: Wiley and Sons, 1984.
31. Goncalo S, Goncalo M, Azenha A, Barros MA, Bastos AS, Brandao FM, et al. Allergic contact dermatitis in children. A multicenter study of the Portuguese Contact Dermatitis Group (GPEDC). *Contact Dermatitis* 1992; 26: 112–115.
32. Menne T. Prevention of nickel allergy by regulation of specific exposures. *Ann Clin Lab Sci* 1996; 26: 133–138.
33. Johansen JD, Menne T, Christophersen J, Kaaber K, Veien NK. Changes in the pattern of sensitization to common contact allergens in Denmark between 1985–86 and 1997–98, with a special view to the effect of preventive strategies. *Br J Dermatol* 2000; 142: 490–495.
34. Wahlberg JE, Skog E. Nickel allergy and atopy. *Br J Dermatol* 1971; 85: 97–104.
35. Veien NK, Hattel T, Laurberg G. Low nickel diet: an open, prospective trial. *J Am Acad Dermatol* 1993; 29: 1002–1007.
36. Kieffer M. Nickel sensitivity: relationship between history and patch test reaction. *Contact Dermatitis* 1979; 5: 398–401.
37. Moller H, Svensson A. Metal sensitivity: positive history but negative test indicates atopy. *Contact Dermatitis* 1986; 14: 57–60.
38. Meijer C, Bredberg M, Fischer T, Widstrom L. Ear piercing, and nickel and cobalt sensitization, in 520 young Swedish men doing compulsory military service. *Contact Dermatitis* 1995; 32: 147–149.
39. Gawkrödger DJ, Vestey JP, Wong W-K, Buxton PK. Contact clinic survey of nickel-sensitive subjects. *Contact Dermatitis* 1986; 14: 165–169.
40. Brune D. Metal release from dental biomaterials. *Biomaterials* 1986; 7: 163–175.
41. Christensen OB, Moller H. Nickel allergy and hand eczema. *Contact Dermatitis* 1975; 1: 129–135.
42. Menne T, Holm NV. Hand eczema in nickel-sensitive female twins. Genetic predisposition and environmental factors. *Contact Dermatitis* 1983; 9: 289–296.