

Supplementary material has been published as submitted. It has not been copyedited, typeset or checked for scientific content by Acta Dermato-Venereologica

Table SI. Affected body regions

	Non naïve N=25	Naïve N=63	Total N=88
Zones	25	62	87
Missing	0	1	1
No zone	4 (16.0%)	7 (11.3%)	11 (12.6%)
Arms and armpits only	0 (0.0%)	1 (1.6%)	1 (1.1%)
Scalp only	1 (4.0%)	2 (3.2%)	3 (3.4%)
Buttocks and thighs only	0 (0.0%)	1 (1.6%)	1 (1.1%)
Knees, lower part of the legs, ankles only	1 (4.0%)	3 (4.8%)	4 (4.6%)
Hands, fingers and nails only	2 (8.0%)	4 (6.5%)	6 (6.9%)
Feet and toes only	0 (0.0%)	2 (3.2%)	2 (2.3%)
Several zones	15 (60.0%)	37 (59.7%)	52 (59.8%)
Face, neck and ears only	2 (8.0%)	5 (8.1%)	7 (8.0%)

Table SII. SF-12 physical and mental dimensions

	Non naïve N=25	Naïve N=63	Total N=88	P-value
SF-12 score : Physical dimension				
N	24	60	84	
Missing	1	3	4	
Mean ± SD	48.30 ± 7.97	51.55 ± 7.18	50.63 ± 7.51	
Median	49.9	52.4	52.1	0.093 [d]
Q1 ; Q3	42.0 ; 54.2	46.7 ; 56.6	45.9 ; 56.3	
Min. ; Max.	32.2 ; 60.1	22.6 ; 62.5	22.6 ; 62.5	
SF-12 score: Mental dimension				
N	24	60	84	
Missing	1	3	4	
Mean ± SD	39.27 ± 10.33	37.26 ± 10.75	37.84 ± 10.61	0.438 [c]
Median	39.2	35.8	37.1	
Q1 ; Q3	30.7 ; 45.6	28.4 ; 45.2	30.1 ; 45.6	
Min. ; Max.	22.4 ; 58.3	17.0 ; 62.4	17.0 ; 62.4	