

**Table SI. Published cases of bullous pemphigoid (BP) associated with anti-programmed cell death protein-1 (PD-1)/programmed death ligand-1 (PD-L1) immunotherapy**

Patient number	Sex/age, years	Tumour type	PD-1/PD-L1 inhibitor	Time to skin toxicity (weeks)	Time to blisters (weeks)	Systemic BP treatment	Action regarding immunotherapy because of BP	CA outcome	BP on last FU
1 (S1)	M/75	Melanoma	Pembrolizumab	9	22	p.o. CS	n/a (discont. before BP)	PD	4w; BP resolved
2 (S2)	M/68	Melanoma	Pembrolizumab	24	78	None	Cont.	PR	5w; BP resolved
3 (S2)	M/72	Melanoma	Pembrolizumab	18	18	p.o. CS; Doxycycline; MTX	Lower frequency	PD	3m; BP stable (CS+MTX ongoing)
4 (S3)	M/80	Melanoma	Nivolumab	20	24	p.o. CS; Antibiotics+NCM	Discont.	CR	4m; BP stable (antibiotics+NCM ongoing)
5 (S3)	M/85	Lung SCC	Nivolumab	12	18	p.o. CS	Initially cont., then discont.	PR	10m; intermittent flares
6 (S3)	F/78	Melanoma	Durvalumab	16	52	none	Discont.	PR	1y; intermittent flares
7 (S4)	M/66	Melanoma	Pembrolizumab	9	n/a	p.o. CS	Discont.	CR	1y; still mild symptoms
8 (S5)	F/73	Lung adeno CA	Nivolumab	6	6	p.o. CS; Doxycycline+NCM	Initially cont., then discont.	PD	42d; BP progress
9 (S6)	M/63	Melanoma	Pembrolizumab	88	88	p.o. CS	Cont.	CR	30m; BP resolved
10 (S7)	F/77	Lung adeno CA	Nivolumab	3	6	p.o. CS; Omalizumab	Withhold, then restarted	n/d	n/d
11 (S8)	F/82	Melanoma	Pembrolizumab	6	32	p.o. CS	Discont.	CR	4m; only pruritus
12 (S9)	M/38	Melanoma	Nivolumab	30	36	none	n/a (discont. before BP)	n/d	n/d
13 (S10)	M/73	Melanoma	Pembrolizumab	18	18	Doxycycline+NCM	Discont.	n/d	n/d
14 (S10)	M/90	Melanoma	Nivolumab	24	24	p.o. CS	Cont.	PR	3m; BP resolved
15 (S11)	F/56	Melanoma	Pembrolizumab	4	24	p.o. + i.v.CS; MTX	Discont.	CR	6m; intermittent flares (CS+MTX ongoing)
16 (S12)	M/58	Lung adeno CA	Atezolizumab	36	64	p.o. CS; Doxycycline	Discont.	n/d	7m; BP stable (CS ongoing)
17 (S13)	M/80	Lung adeno CA	Nivolumab	52	80	p.o. CS; Rituximab	Discont.	CR	8m; BP resolved
18 (S14)	M/60	Melanoma	Nivolumab	12	52	p.o. CS	Discont.	n/d	2w; BP resolved
19 (S15)	M/78	Melanoma	Nivolumab	47	47	None	n/a (discont. before BP)	PD	1m; BP resolved
20 (S16)	M/42	Melanoma	Pembrolizumab	32	44	p.o. CS; Doxycycline+NCM	Discont.	SD	4m; BP resolved
21 (S16)	M/65	Melanoma	Pembrolizumab	12	51	p.o. CS	Cont.	CR	n/d
22 (S17)	F/82	Melanoma	Pembrolizumab	24	24	p.o. CS	Cont.	PR	6m; mild symptoms
23 (S18)	M/77	Melanoma	Nivolumab	17	17	p.o. CS; Doxycycline	Cont.	CR	2w; BP resolved
24 (S19)	M/66	Melanoma	Nivolumab	16	28	p.o. CS	Cont.	PR	14m; BP stable (CS ongoing)
25 (S19)	M/78	Melanoma	Nivolumab	12	16	p.o. CS	Cont.	PD	1m; BP resolved
26 (S19)	F/68	NSCLC	Nivolumab	16	16	p.o. CS	Cont.	PD	3w; BP resolved
27 (S20)	M/48	Melanoma	Nivolumab	31	31	p.o. CS	n/a (discont. before BP)	PD	~9m; BP resolved
28 (S21)	M/64	Melanoma	Pembrolizumab	12	21	p.o. CS	Withhold, then restarted	n/d	~9m; BP resolved
29 (S21)	M/71	Melanoma	Pembrolizumab	27	27	p.o. CS	Discont.	PD	n/d
30 (S22)	F/72	NSCLC	Nivolumab	1	6	p.o. CS; Minocycline+NCM	Discont.	n/d	6w; BP stable
31 (S23)	M/78	RCC	Nivolumab	10	10	Rituximab	Discont.	CR	3m; BP resolved
32 (S24)	M/68	NSCLC	Nivolumab	52	52	p.o. CS	Cont.	SD	4w; BP improved
33 (S25)	M/68	Melanoma	Nivolumab	32	32	p.o. CS; plasmaexchange; Rituximab	Discont.	n/d	n/d; BP resolved
34 (S26)	M/77	NSCLC	Nivolumab	12	12	Doxycycline; p.o.CS	Discont.	SD	18m; stillpruritus
35 (S26)	F/77	NSCLC	Nivolumab	3	3	p.o. CS; Omalizumab	Discont.	PD	17m; BP resolved
36 (S26)	M/69	NSCLC	Nivolumab	4	4	p.o. CS; Doxycycline+NCM	Discont.	PD	15m; BP flares
37 (S26)	M/68	Melanoma	Pembrolizumab	40	40	p.o. CS; Doxycycline+NCM	Discont.	PD	7m; BP flares
38 (S26)	M/48	RCC	Nivolumab	28	28	p.o. CS; Doxycycline+NCM	n/a (discont. before BP)	PD	6m; BP resolved
39 (S26)	F/61	NSCLC	Pembrolizumab	36	36	p.o. CS; Doxycycline	Discont.	PD	7m; BP resolved
40 (S26)	F/77	Urothelial CA	Atezolizumab	20	20	p.o. CS; Omalizumab; MTX	Discont.	PR	22m; still pruritus
41 (S27)	F/56	Endometrial CA	Pembrolizumab	16	60	p.o. CS	Cont.	n/d	n/d; BP resolved
42 (S28)	F/86	Melanoma	Pembrolizumab	68	68	p.o. CS	Discont.	n/d	3m; BP stable (CS ongoing)
43 (S28)	M/82	Melanoma	Pembrolizumab	36	36	p.o.+i.v. CS	Discont.	n/d	4m; BP flares
44 (S29)	M/80	NSCLC	Pembrolizumab	21	21	p.o. CS	Discont.	n/d	2w; BP stable (CS ongoing)
45 (S30)	F/69	Melanoma	Nivolumab	22	22	p.o. CS; Dapsone	Discont.	PR	4w; still pruritus
46 (S31)	M/63	Lung adeno CA	Nivolumab	52	104	Doxycycline; p.o.+i.v. CS; Dapsone	Cont.	n/d	16w; BP stable (Dapsone ongoing)
47 (S32)	F/75	Melanoma	Nivolumab	44	44	p.o. CS	Discont.	n/d	10m; BP resolved
48 (S33)	M/78	Melanoma	Nivolumab	40	n/a	Doxycycline+NCM; p.o. CS; Omalizumab	Discont.	CR	15m; BP resolved
49 (S33)	M/78	Esophagealadeno CA	Pembrolizumab	2	n/a	p.o. CS; Doxycycline+NCM; Omalizumab	Cont.	PD	15m; BP resolved
50 (S33)	M/62	Lung adeno CA	Pembrolizumab	52	n/a	p.o. CS; Tetracycline+NCM; Rituximab	Cont.	PD	n/d; BP resolved
51 (S33)	M/58	Melanoma	Pembrolizumab	28	n/a	p.o. CS; Doxycycline+NCM; Rituximab	Discont.	PR	n/d; BP resolved
52 (S34)	M/62	Clear cell CA	Nivolumab	Months	Months	p.o. CS	Withhold, then restarted	CR	3m; BP resolved
53 (S35)	M/87	Urothelial CA	Atezolizumab	76	76	Doxycycline+NCM	Withhold, then restarted	n/d	~4m; BP resolved

**Table SI. Contd.**

Patient number	Sex/age, years	Tumour type	PD-1/PD-L1 inhibitor	Time to skin toxicity (weeks)	Time to blisters (weeks)	Systemic BP treatment	Action regarding immunotherapy because of BP	CA outcome	BP on last FU
54 (S36)	M/71	RCC	Nivolumab	52	72	p.o. CS; IvIG	Discont.	n/d	8m; BP stable (CS ongoing)
55 (S37)	M/60	Lung adenoCA	Pembrolizumab	18	18	p.o. CS	Discont.	n/d	n/d
56 (S38)	M/84	NSCLC	Pembrolizumab	48	48	p.o. CS	Discont.	CR	11m; BP resolved
57 (S39)	M/68	Cutaneous SCC	Cemiplimab	6	6	p.o. CS; Rituximab	Discont.	n/d	6m; BP resolved
58 (S40)	M/71	Melanoma	Pembrolizumab	36	n/a	p.o. CS; Dapsone; Doxycycline+NCM; Rituximab	Discont.	n/d	18m; still pruritus

adenoCA: adenocarcinoma; CA: carcinoma; CR: complete response; CS: corticosteroids; F: female; IvIG: intravenous immunoglobulin; MTX: methotrexate; n/a: not applicable; M: male; NCM: niacinamide; n/d: not documented; NSCLC: non-small cell lung cancer; PD: progressive disease; PR: partial response; RCC: renal cell carcinoma; SCC: squamous cell carcinoma; SD: stable disease; BP: bullous pemphigoid; w: weeks; m: months; p.o.: per os; i.v.: intravenously; cont.: continued; discont: discontinued; FU: follow up; PD-1: programmed cell death protein 1; PD-L1: programmed cell death ligand 1.

#### SUPPLEMENTARY REFERENCES

- S1. Carlos G, Anforth R, Chou S, Clements A, Fernandez-Peñas P. A case of bullous pemphigoid in a patient with metastatic melanoma treated with pembrolizumab. *Melanoma Res* 2015; 25: 265–268.
- S2. Hwang SJ, Carlos G, Chou S, Wakade D, Carlini MS, Fernandez-Peñas P. Bullous pemphigoid, an autoantibody-mediated disease, is a novel immune-related adverse event in patients treated with anti-programmed cell death 1 antibodies. *Melanoma Res* 2016; 26: 413–416.
- S3. Naidoo J, Schindler K, Querfeld C, Busam K, Page DB, Postow MA, et al. Autoimmune bullous skin disorders with immune checkpoint inhibitors targeting PD-1 and PD-L1. *Cancer Immunol Res* 2016; 4: 383–389.
- S4. Brunet-possent F, Mignot S, Deschamps L, Descamps V. Antiepidermis autoantibodies induced by anti-PD-1 therapy in metastatic melanoma. *Melanoma Res* 2016; 26: 540–543.
- S5. Jour G, Glitz IC, Ellis RM, Torres-Cabala CA, Tetzlaff MT, Li JY, et al. Autoimmune dermatologic toxicities from immune checkpoint blockade with anti-PD-1 antibody therapy: a report on bullous skin eruptions. *J Cutan Pathol* 2016; 43: 688–696.
- S6. Mochel MC, Ming ME, Imadoujemu S, Gangadhar TC, Schuchter LM, Elenitas R, et al. Cutaneous autoimmune effects in the setting of therapeutic immune checkpoint inhibition for metastatic melanoma. *J Cutan Pathol* 2016; 43: 787–791.
- S7. Damsky W, Kole L, Tomayko MM. Development of bullous pemphigoid during nivolumab therapy. *JAAD Case Rep* 2016; 2: 442–444.
- S8. Lomax AJ, Ge K, Anand S, McNeil C, Lowe P. Bullous pemphigoid-like reaction in a patient with metastatic melanoma receiving pembrolizumab and previously treated with ipilimumab. *Australas J Dermatol* 2016; 57: 333–335.
- S9. Kuwatsuka Y, Iwanaga A, Kuwatsuka S, Okubo Y, Murayama N, Ishii N, et al. Bullous pemphigoid induced by ipilimumab in a patient with metastatic malignant melanoma after unsuccessful treatment with nivolumab. *J Dermatol* 2018; 45: e21–e22.
- S10. Bandino JP, Perry DM, Clarke CE, Marchell RM, Elston DM. Two cases of anti-programmed cell death 1-associated bullous pemphigoid-like disease and eruptive keratoacanthomas featuring combined histopathology. *J Eur Acad Dermatol Venereol* 2017; 31: e378–e380.
- S11. Rofe O, Bar-Sela G, Keidar Z, Sezin T, Sadik CD, Bergman R. Severe bullous pemphigoid associated with pembrolizumab therapy for metastatic melanoma with complete regression. *Clin Exp Dermatol* 2017; 42: 309–312.
- S12. Russo I, Sacco G, Frega S, Polo V, Pasello G, Alaibac M. Immunotherapy-related skin toxicity: bullous pemphigoid in a lung adenocarcinoma patient treated with the anti-PDL1 anti-body atezolizumab. *Eur J Dermatol* 2017; 27: 205–208.
- S13. Sowerby L, Dewan AK, Granter S, Gandhi L, LeBoeuf NR. Rituximab treatment of nivolumab-induced bullous pemphigoid. *JAMA Dermatol* 2017; 153: 603–605.
- S14. Kwon CW, Land AS, Smoller BR, Scott G, Beck LA, Mercurio MG. Bullous pemphigoid associated with nivolumab, a programmed cell death 1 protein inhibitor. *J Eur Acad Dermatol Venereol* 2017; 31: e349–e350.
- S15. Hirotsu K, Chiou AS, Chiang A, Kim J, Kwong BY, Pugliese S. Localized bullous pemphigoid in a melanoma patient with dual exposure to PD-1 checkpoint inhibition and radiation therapy. *JAAD Case Rep* 2017; 3: 404–406.
- S16. Parakh S, Nguyen R, Opie JM, Andrews MC. Late presentation of generalised bullous pemphigoid-like reaction in a patient treated with pembrolizumab for metastatic melanoma. *Australas J Dermatol* 2017; 58: e109–e112.
- S17. Amber KT, Valdebran M, Lu Y, De Feraudy S, Linden KG. Localized pretibial bullous pemphigoid arising in a patient on pembrolizumab for metastatic melanoma. *J Dtsh Dermatol Ges* 2018; 16: 196–198.
- S18. Tanita K, Fujimura T, Kambayashi Y, Tsukada A, Sato Y, Hashimoto A, et al. Intensity-modulated radiotherapy triggers onset of bullous pemphigoid in a patient with advanced melanoma treated with nivolumab. *Case Rep Oncol* 2018; 11: 114–118.
- S19. Le Naour S, Peuvrel L, Saint-Jean M, Dreno B, Quereux G. Three new cases of bullous pemphigoid during anti-PD-1 antibody therapy. *J Eur Acad Dermatol Venereol* 2018; 32: e104–e106.
- S20. Anastasopoulou A, Papaxoinis G, Diamantopoulos P, Christofidou E, Benopoulou O, Stratigos A, et al. Bullous pemphigoid-like skin lesions and overt eosinophilia in a patient with melanoma treated with nivolumab: case report and review of the literature. *J Immunother* 2018; 41: 164–167.
- S21. Thomsen K, Diernaejs J, Ollegaard TH, Spaun E, Vestergaard C. Bullous pemphigoid as an adverse reaction to pembrolizumab: two case reports. *Case Rep Dermatol* 2018; 10: 154–157.
- S22. Lopez AT, Geskin L. A case of nivolumab-induced bullous pemphigoid: review of dermatologic toxicity associated with programmed cell death protein-1/programmed death ligand-1 inhibitors and recommendations for diagnosis and management. *Oncologist* 2018; 23: 1119–1126.
- S23. Grimaux X, Delva R, Jadaud E, Croue A. Nivolumab-induced bullous pemphigoid after radiotherapy and abscopal effect. *Australas J Dermatol* 2019; 60: e235–e236.
- S24. Panariello L, Fattore D, Annunziata MC, Piantedosi F, Gilli M, Fabbrocini G. Bullous pemphigoid and nivolumab: dermatologic management to support and continue oncologic therapy. *Eur J Cancer* 2018; 103: 284–286.
- S25. Ridpath A V, Rzepka P V, Shearer SM, Scrape SR, Olenki TE, Kaffenberger BH. Novel use of combination therapeutic plasma exchange and rituximab in the treatment of nivolumab-induced bullous pemphigoid. *Int J Dermatol* 2018; 57: 1372–1374.
- S26. Siegel J, Totonchy M, Damsky W, Berk-Krauss J, Castiglione F, Szoln M, et al. Bullous disorders associated with anti-PD-1 and anti-PD-L1 therapy: a retrospective analysis evaluating the clinical and histopathologic features, frequency, and impact on cancer therapy. *J Am Acad Dermatol* 2018; 79: 1081–1088.
- S27. Honigman AD, Lai F, Elakis J, Prall O, Goh M, McCormack C. Pembrolizumab-induced sarcoid granulomatous panniculitis and bullous pemphigoid in a single patient. *Clin Case Rep* 2019; 7: 773–775.
- S28. Sun CW, Grossman SK, Aphale A, Hsu S. Pembrolizumab-induced bullous pemphigoid. *JAAD Case Rep* 2019; 5: 362–364.
- S29. Adachi E, Honda T, Nonoyama S, Irie H, Yamamura K, Otsuka A, et al. Severe bullous pemphigoid in a metastatic lung cancer patient treated with pembrolizumab. *J Dermatol* 2019; 46: e232–e233.
- S30. Sadik CD, Langan EA, Grätz V, Zillikens D, Terheyden P. Checkpoint inhibition may trigger the rare variant of anti-LAD-1 IgG-negative bullous pemphigoid. *Front Immunol* 2019; 10: 1934.
- S31. Grän F, Goebeler M, Gesierich A. Bullous pemphigoid developing upon immune checkpoint inhibition with nivolumab. *Eur J Dermatol* 2019; 29: 448–449.
- S32. Aoki Y, Miyagawa F, Miyashita K, Nishimura Y, Hashimoto T, Asada H. Nivolumab-associated bullous pemphigoid reactions involving multiple epitopes. *Eur J Dermatol* 2019; 29: 552–554.
- S33. Singer S, Nelson CA, Lian CG, Dewan AK, LeBoeuf NR. Nonbullous pemphigoid secondary to PD-1 inhibition. *JAAD Case Rep* 2019; 5: 898–903.
- S34. Palla AR, Smith E, Doll D. Bullous pemphigoid associated with the 480-mg nivolumab dose in a patient with metastatic renal cell carcinoma. *Immunotherapy* 2019; 11: 1187–1192.
- S35. Kosche C, Owen JL, Sadowsky LM, Choi JN. Bullous dermatoses secondary to anti-PD-L1 agents: a case report and review of the literature. *Dermatol Online J* 2019; 25: pii: 13030/qt7gg675.
- S36. Matsui Z, Makino T, Ishii N, Hashimoto T, Shimizu T. Detection of IgG antibodies to BP180 NC16a and C-terminal domains and LAD-1 in nivolumab-associated bullous pemphigoid. *Eur J Dermatol* 2019; 29: 554–555.
- S37. Hara K, Yamasaki K, Yamada S, Yatera K. Bullous pemphigoid in a patient treated with pembrolizumab. *Intern Med* 2020; 59: 139–140.
- S38. Muto Y, Namikawa K, Fujiwara Y, Mori T, Nakano E, Takahashi A, et al. Bullous pemphigoid induced by pembrolizumab in a patient with non-small-cell lung cancer who achieved durable complete response despite treatment cessation and long-term corticosteroid administration: a case report. *J Dermatol* 2020; 47: e9–e11.
- S39. Virgen CA, Nguyen TA, Di Raimondo C, Amini A, Margolin KA, Parekh V, et al. Bullous pemphigoid associated with cemiplimab therapy in a patient with locally advanced cutaneous squamous cell carcinoma. *JAAD Case Rep* 2020; 6: 195–197.
- S40. Morris LM, Lewis HA, Cornelius LA, Chen DY, Rosman IS. Neutrophil-predominant bullous pemphigoid induced by checkpoint inhibitors: a case series. *J Cutan Pathol* 2020; 47: 742–746.