

Table SIII. Quick inventory of depressive symptomatology-self report (QIDS-SR16) total score categories at each study visit

Severity Category	Blinded treatment period		Open-label treatment period		
	PBO (n = 74)		PBO/IXE Q4W (n = 65)		
	Week 0 (Nx = 74) n (%)	Week 12 (Nx = 64) n (%)	Week 24 (Nx = 65) n (%)	Week 40 (Nx = 64) n (%)	Week 52 (Nx = 63) n (%)
None	49 (66.2)	46 (71.9)	51 (78.5)	50 (78.1)	54 (85.7)
Mild	17 (23.0)	10 (15.6)	12 (18.5)	13 (20.3)	8 (12.7)
Moderate	7 (9.5)	6 (9.4)	2 (3.1)	1 (1.6)	1 (1.6)
Severe	0	2 (3.1)	0	0	0
Very Severe	1 (1.4)	0	0	0	0

Severity Category	IXE Q2W (n = 75)		IXE Q2W/IXE Q4W (n = 74) ^a		
	Week 0 (Nx = 75) n (%)	Week 12 (Nx = 71) n (%)	Week 24 (Nx = 69) n (%)	Week 40 (Nx = 67) n (%)	Week 52 (Nx = 66) n (%)
None	55 (73.3)	66 (93.0)	64 (92.8)	55 (82.1)	54 (81.8)
Mild	15 (20.0)	4 (5.6)	3 (4.3)	10 (14.9)	10 (15.2)
Moderate	5 (6.7)	1 (1.4)	1 (1.4)	1 (1.5)	2 (3.0)
Severe	0	0	1 (1.4)	1 (1.5)	0
Very Severe	0	0	0	0	0

^aPatients initially randomized to IXE Q2W at Week 0 who received at least 1 dose of IXE Q4W during the open-label treatment period. Percentages were calculated as n/Nx where n is the number of patients in the specified category and Nx is the total number of patients with nonmissing values. QIDS-SR16 categories are determined based on total score as follows: None: 0–5, Mild: 6–11, Moderate: 11–15, Severe: 16–20, Very Severe: 21–27. IXE: ixekizumab; PBO: placebo; Q2W: every 2 weeks; Q4W: every 4 weeks; QIDS-SR16: Quick Inventory of Depressive Symptomatology-Self Report (16-items).