

Supplementary material has been published as submitted. It has not been copyedited, typeset or checked for scientific content by Acta Dermato-Venereologica

**Table SI. Follow-up data for 14 patients with mycosis fungoides palmaris et plantaris (MFPP)**

Patients	Relapse	Time to relapse (months)	Total duration of alitretinoin treatment (weeks)	Treatment course	Total follow-up duration (months)	PGA grade at last follow-up	Alitretinoin-related side-effects
1	+	13	34	Ali (R) → recur, PUVA (NR) → Ali (R)	77	almost clear	Dry mouth
2	+	4	29	Ali	19	clear	Elevated TG
3	+	11	49	Ali	25	clear	–
4	NA	NA	12	Ali (NR) → denied treatment	17	moderate	–
5	+	61	26	Ali	68	almost clear	–
6	–	–	16	Ali	13	clear	–
7	–	–	13	Ali	13	almost clear	Headache
8	NA	NA	152	Ali (PR) → UVA-1, MTX (NR) → Ali (PR)	71	mild	Headache, Elevated TG
9	–	–	63	Ali	34	clear	Headache, Elevated TG
10	–	–	51	Ali (R) → Ali, PUVA (R)*	37	clear	Dry skin, Flushing
11	+	10	78	Ali	35	almost clear	–
12	NA	NA	91	Ali (PR) → UVA-1, MTX (PR)	32	moderate	Headache, Elevated TG
13	NA	NA	37	Ali (PR) → MTX (R)	16	almost clear	Headache, Dry skin
14	NA	NA	29	Ali	7	mild	–

Ali: alitretinoin; MTX: methotrexate; NA: not applicable; NR: no response; PR: partial response; PUVA: psoralen plus ultraviolet A; R: response; TG: triglyceride; UVA-1: ultraviolet A-1.

\*The clinical response was achieved with alitretinoin monotherapy, but since some lesions persisted, PUVA was added.