

Table SI. History of use of facial and hair cosmetics, sunscreens and hair care practices in patients with frontal fibrosing alopecia and controls obtained by questionnaire

	FFA (n=20) n (%)	Controls (n=40) n (%)	p-value
Facial cosmetics			
Leave on			
< Twice weekly	3 (15)	9 (22.5)	0.494
≥ Twice weekly	17 (85)	31 (77.5)	
Wash off			
< Twice weekly	13 (65)	25 (62.5)	0.85
≥ Twice weekly	7 (35)	15 (37.5)	
Sunscreens			
No	3 (15)	9 (22.5)	0.646
Irregular (before sun exposure)	4 (20)	12 (40)	
≥ Twice weekly at summer	3 (15)	4 (10)	
≥ Twice weekly all year	10 (50)	15 (37.5)	
Hair cosmetics			
< Twice weekly	11 (55)	19 (47.5)	0.584
≥ Twice weekly	9 (45)	21 (52.5)	
Hair care practices			
Heated hair styling			
No	12 (60)	20 (50)	0.76
< Twice weekly	5 (25)	12 (30)	
≥ Twice weekly	3 (15)	5 (20)	
Dyeing			
No	2 (10)	3 (7.5)	0.652
< Once monthly	11 (55)	18 (45)	
≥ Once monthly	7 (35)	19 (47.5)	
Traction practices			
No	16 (80)	31 (77.5)	0.825
Yes	4 (20)	9 (22.5)	
Straightening/perming			
No	17 (85)	35 (87.5)	0.672
< Once a year	3 (15)	49 (10)	
≥ Once a year	-	1 (2.5)	