

Table SI. Data on all 50 patients with prurigo

Patient demographics and duration of disease	26 females, mean 64 years (range 22–89 years); mean 57 years on first visit at the centre, mean treatment period 4 years (range 0–27 years)
Comorbidities	High blood pressure 16 p, heart disease 11 p, rheumatologic disease 8 p, lung disease <i>n</i> = 6, arthritis <i>n</i> = 6, diabetes <i>n</i> = 6, gastrointestinal disease <i>n</i> = 5, cancer <i>n</i> = 3, liver disease <i>n</i> = 2, haematological disease <i>n</i> = 2, kidney disease <i>n</i> = 2, depression <i>n</i> = 8, psychosocial stress <i>n</i> = 3, psychiatric or neurologic disease <i>n</i> = 2
Skin disease	Atopic dermatitis <i>n</i> = 14, nummular/microbial/contact eczema <i>n</i> = 6, insect stings <i>n</i> = 2, bullous pemphigoid <i>n</i> = 2
Topical treatments	Emollients <i>n</i> = 50, potent steroid creams (without or with occlusion) <i>n</i> = 50, tacrolimus <i>n</i> = 7, capsaicin <i>n</i> = 1, potassium permanganate baths <i>n</i> = 2
Systemic treatments	Antihistamines <i>n</i> = 50, UVB-NB <i>n</i> = 31, psoralen plus ultraviolet A (PUVA) <i>n</i> = 5, systemic steroids <i>n</i> = 15, intralesional steroids <i>n</i> = 6, antidepressants/psychopharmacy/ psychotherapy <i>n</i> = 8, gabapentin <i>n</i> = 8, methotrexate <i>n</i> = 19, ciclosporin <i>n</i> = 4, azathioprine <i>n</i> = 2, dapsone <i>n</i> = 2, acitretin <i>n</i> = 1, omalizumab <i>n</i> = 1

UVB-NB: ultraviolet B narrow band.