

Table SI. Measures of central tendency and number of patients for neutrophil-to-lymphocyte ratios at baseline and 90-day bins under treatment (cf. Fig. 1).

Bin [days]	Baseline	0–90	90–180	180–270	270–360	360–450	450–540	540–630	630–720	720–810	810–900	900–990
All												
gmean	2.91	2.22	2.02	2.03	2.00	2.01	2.02	2.06	2.12	1.93	1.95	1.86
95% CI	2.7–3.2	2.1–2.4	1.9–2.2	1.9–2.2	1.9–2.1	1.9–2.2	1.9–2.2	1.8–2.3	1.9–2.4	1.8–2.1	1.8–2.1	1.7–2.0
Mean	3.62	2.50	2.21	2.22	2.12	2.15	2.12	2.40	2.31	1.99	1.98	1.91
95% CI	0.5	0.2	0.2	0.2	0.2	0.2	0.2	0.7	0.3	0.2	0.2	0.2
Median	2.74	2.22	1.98	2.01	2.00	2.00	2.02	1.91	2.13	1.87	1.81	1.78
25Q–75Q	1.9–4.0	1.7–2.9	1.5–2.5	1.5–2.6	1.5–2.4	1.5–2.4	1.5–2.6	1.5–2.6	1.4–2.7	1.5–2.5	1.5–2.4	1.4–2.3
n	169	183	155	134	116	102	87	77	68	68	58	50
TNF-α												
gmean	2.89	2.17	1.95	1.96	1.97	1.95	2.00	1.99	1.99	1.84	1.89	1.80
95% CI	2.6–3.2	2.0–2.3	1.8–2.1	1.8–2.1	1.8–2.1	1.8–2.1	1.8–2.2	1.8–2.2	1.8–2.2	1.7–2.0	1.7–2.1	1.6–2.0
Mean	3.63	2.45	2.07	2.13	2.08	2.06	2.10	2.40	2.11	1.88	1.91	1.83
95% CI	0.6	0.2	0.2	0.2	0.2	0.2	0.2	0.8	0.2	0.2	0.2	0.2
Median	2.71	2.12	1.94	1.90	1.96	1.95	2.00	1.84	2.09	1.82	1.79	1.74
25Q–75Q	1.9–3.9	1.7–2.8	1.5–2.4	1.5–2.4	1.5–2.3	1.5–2.4	1.5–2.5	1.4–2.4	1.3–2.6	1.3–2.4	1.4–2.4	1.4–2.2
n	148	160	133	115	100	88	75	66	59	60	52	46
IL12/23												
gmean	3.07	2.54	2.54	2.53	2.21	2.44	2.17	2.51	3.19	2.67	2.48	2.62
95% CI	2.4–4.0	2.1–3.1	2.0–3.2	2.1–3.0	1.8–2.7	2.0–3.0	1.9–2.5	2.0–3.2	2.3–4.5	2.2–3.3	1.9–3.2	1.9–3.6
Mean	3.62	2.86	3.05	2.75	2.38	2.74	2.25	2.72	3.60	2.80	2.60	2.87
95% CI	1.0	0.6	0.9	0.5	0.4	0.8	0.3	0.6	1.3	0.6	0.6	0.9
Median	3.00	2.56	2.28	2.89	2.41	2.27	2.36	2.56	2.69	2.45	2.51	2.87
25Q–75Q	2.5–4.5	2.0–3.7	1.8–3.2	1.9–3.5	1.8–3.2	1.9–3.1	1.9–2.7	1.9–3.7	2.0–5.2	2.2–3.2	1.8–3.4	2.2–3.5
n	21	23	22	19	16	14	12	11	9	8	6	4

gmean: geometric mean; C: confidence interval; 25Q–75Q: 25th–75th quartile; TNF-α: tumour necrosis factor-α antagonist; IL-12/23: interleukin-12/23 antagonist.