

Table SI. Blood analyses before treatment and at 10, 60 days after initiated treatment

Day	Inflammation (blood)						Coagulation	Other	
(RV)	CRP (<3)	SR (<20)	Procalcitonin (<0.5)	Leukocytes (3.5-8.8)	Eosinophils (0.0-0.5)		INR (<1.3)	PEth 16:0/ 18:1 (<0.05)	
0	32	104	0.24	9.8	0.2		1	0.08	
10	1	62	x	5.7	0.2		1		
60	x	6	x	5.3	x		x		
Liver-enzymes (blood)							Autoimmune hepatitis		
(RV)	ALP (0.7-1.9)	GT (1.3)	Bilirubin (<26)	ALAT (<0.76)	ASAT (<0.61)	Albumin (36-45)	ANA (<1)	mtDNA (<4)	SMA, LKM (neg)
0	12.2	8.7	23	8.57	4.33	26	<0.1	1.5	neg
10	2.9	1.8	3	0.58	0.43	26	x	x	x
60	1	0.25	x	0.3	0.34	40	x	x	x
Further hematology (blood)									
(RV)	Hemoglobin (117-153)	MCV (82-98)	MCH (27-33)	Leukocytes (3.5-8.8)	Trombocytes (165-387)	LD (3.5)	Iron (9-34)	Ferritin (13-150)	Transferrin (1.90-3.30)
0	115	86	28	9.8	435	5.3	14	190	1.89
10	111	88	29	5.7	500	x	x	x	x
60	132	90	28	5.3	337	x	x	x	x
Cerebrospinal fluid (CSF)									
(RV)	Clarity (clear)	Colour (colourless)	Leukocytes total (0-5)	Leukocytes poly (<1)	Leukocytes mono (0-5)	Erythrocytes (<1)			
0	Clear	Colourless	26	4	22	5			

Analyses at “day 0” were gathered at days -6 to 0, before initiated antibiotic treatment. Day 10 is the last day of treatment with IV penicillin G.
RV: reference value.