

**Table SI. Reports of multiple keratoacanthoma centrifugum marginatum (KCM) published in the literature in English**

Author, year	Sex/age, years	Disease duration, months	Family history	Localization of lesions	Treatment	Duration of treatment, months	Outcome	Relevant associated conditions
Cherif et al., 2002 (3)	M/63	6	NA	Lower extremities	Acitretin 1 mg/kg	5	Complete clearance	NA
Ogasawara et al., 2003 (4)	M/48	18	No	Face, upper and lower extremities	Etretinate 20–30 mg/day	12	Complete clearance	Colonic polyps
Mangas et al., 2004 (5)	M/21	12	No	Lower extremities, buttocks	Isotretinoin 1 mg/kg MTX 15 mg/week + prednisone p.o. 1 mg/kg/day followed by MTX 7.5 mg/week	2 2 + 6	No effect Complete clearance	None
Aroni et al., 2007 (7)	F/45	12	NA	Lower extremities	Surgical excision Acitretin 20 mg/day Imiquimod cream Tretinoin 0.1% cream	– NA 7 NA	Relapse Improvement; stopped due to side-effects Complete clearance Lost to follow-up	None Schizophrenia
Khaled et al., 2010 (8)	M/48	36	NA	Lower extremities	Intralesional 10% triamcinolone acetonide Tretinoin 0.1% gel	0.5 2	No effect Almost complete clearance	None
Binitha et al., 2011 (9)	F/50	24	No	Upper extremities, upper trunk left ear	Oral isotretinoin 0.5 mg/kg	10	Regression of the majority of lesions; stopped due to concerns of skeletal hyperostosis	None
Dogra et al., 2017 (11)	M/5	36	No	Neck, back, upper and lower extremities	MTX 0.3 mg/kg/week Acitretin 25 mg/day (0.3 mg/kg)	NA 1	Disease stabilization Improvement (subsequently lost to follow-up)	None Primary myelofibrosis

MTX: methotrexate; NA: not available; M: male; F: female.