

Table SI. Distribution of the main sub-epidermal autoimmune bullous diseases in different countries of the world

Author (ref)	Country	Total	BP n (%)	MMP n (%)	PG n (%)	EBA n (%)	LAD n (%)
Bernard et al. (2)	France	100	74 (74)	12 (12)	4 (4)	3 (3)	5 (5)
Zillikens et al. (3)	Germany	80	61 (76)	8 (10)	5 (6)	2 (3)	2 (3)
Wong & Chua (4)	Singapore	67	59 (88)	0	0	4 (6)	2 (3)
Nanda et al. (5)	Kuwait	68	27 (40)	1 (1)	24 (36)	3 (4)	9 (13)
Bertram et al. (6)	Germany	41	27 (66)	4 (10)	4 (10)	1 (2)	2 (5)
Baicán et al. (7)	Romania	48	40 (83)	0	1 (2)	2 (4)	3 (6)
Zaraa et al. (8)	Tunisia	73	41 (56)	2 (3)	18 (24)	1 (1)	11 (15)
Daneshpazhooh et al. (9)	Iran	195	163 (84)	10 (5)	10 (5)	7 (3)	5 (2)
Milinkovic et al. (10)	Serbia	527	471 (89)	13 (2)	7 (1)	9 (2)	27 (5)
Present study (2017)	France	642	538 (84)	58 (9)	13 (2)	7 (1)	26 (4)

BP: bullous pemphigoid; EBA: epidermolysis bullosa acquisita; LAD: linear IgA dermatosis; MMP: mucous membrane pemphigoid; PG: pemphigoid gestationis; Total: total number of included patients.