

Table SI. Clinical characteristics of the study group

	Patients with chronic plaque psoriasis <i>n</i> = 65	Controls <i>n</i> = 35
Sex, female/male, <i>n</i> (%)	28 (43.1)/37 (56.9)	16 (45.7)/19 (54.3)
Age, years, mean±SD	48.33±17.95	46.41±14.11
Disease duration, years, mean±SD [range]	14.35±13.32 [0.8–47]	
PASI, mean±SD	14.23±7.07	
BMI, mean±SD	27.89±15.94	26.76±13.74
Hypertension, <i>n</i> (%) ^a		
Total	21 (32.31)	10 (28.57)
Use of hypotensive therapy	18 (27.69)	9 (25.71)
Dyslipidaemia, <i>n</i> (%) ^b		
Total	11 (16.92)	5 (14.28)
Use of lipid-lowering therapy	9 (13.85)	5 (14.28)
Obesity, <i>n</i> (%) ^c	12 (18.46)	5 (14.28)
Smoking habit, <i>n</i> (%) ^d	18 (25.69)	6 (17.14)

^aSystolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg or use of antihypertensive medication. ^bElevated level of: total cholesterol and/or low-density lipoprotein cholesterol and/or triglycerides and/or use of lipid-lowering therapy. ^cSubjects with body mass index (BMI) ≥ 30 . ^dSmoking at least 1 cigarette per day during the last 6 months. SD: standard deviation; PASI: Psoriasis Area and Severity Index.