

Table SI. Characteristics of patients with chronic liver disease with and without pruritus

Demographic data	Pruritus (-)	Pruritus (+)	<i>p</i> -value
Sex, male/female, <i>n</i>	4/3	3/10	
Age, years, median (IQR)	71 (66–79)	62 (53–72.5)	0.101
Aetiology, <i>n</i>			
Autoimmune hepatitis (AIH)	0	3	
Viral infection (HBV, HCV)	0	3	
Alcoholic liver cirrhosis	0	1	
Primary biliary cholangitis	6	4	
Overlap syndrome (PBC+AIH)	1	0	
Others	0	2	
Comorbidity, <i>n</i>			
Rheumatoid arthritis		1	
Laboratory data, median (IQR)			
AST (IU/l)	23 (23–29)	43 (33.5–87)	0.017
ALT (IU/l)	16 (12–20)	44 (19.5–81)	0.011
ALP (IU/l)	361 (253–523)	362 (332.5–1230)	0.523
GGT (IU/l)	33 (16–84)	133 (40.5–321.5)	0.057
TBil (mg/dl)	0.5 (0.4–0.8)	0.8 (0.65–2.6)	0.037

IQR: interquartile range; HBV: hepatitis B virus; HCV: hepatitis C virus; PBC: primary biliary cholangitis; AST: aspartate transaminase; ALT: alanine transaminase; ALP: alkaline phosphatase; GGT: gamma-glutamyltransferase; TBil: total bilirubin.