

Table SI. Basal cell carcinoma (BCC) and squamous cell carcinoma (SCC) patient characteristics according to satisfaction with information provision^a

	Dissatisfied patients ^b n = 237	Satisfied patient ^b n = 422	p-value
Age at time of survey, years, median (IQR)	75 (16)	71 (15)	0.015
Years since diagnosis, median (IQR)	4 (0.6)	4 (0.5)	0.363
Sex, n (%)			0.148
Male	113 (33)	226 (67)	
Female	124 (39)	196 (61)	
Education, n (%)			0.051
Low	74 (41)	104 (58)	
Medium	142 (34)	279 (66)	
High	15 (30)	35 (70)	
Current occupation, n (%)			0.094
(Self-) Employed	48 (30)	114 (70)	
Unemployed	168 (37)	289 (63)	
Partnership, n (%)			0.002
Partner	163 (33)	339 (67)	
No partner	70 (47)	78 (53)	
Tumour type, n (%)			0.548
BCC	203 (36)	358 (64)	
SCC	18 (32)	38 (68)	
Location of tumour, n (%)			0.004
Face	197 (39)	311 (61)	
Other	40 (26)	111 (74)	
Type BCC, n (%)			0.633
Multifocal superficial	14 (23)	48 (77)	
Infiltrating	51 (50)	51 (50)	
Nodular	124 (34)	238 (66)	
Other	14 (40)	21 (60)	
Comorbidity, n (%)			< 0.001
No	100 (28)	260 (72)	
1	33 (40)	50 (60)	
≥ 2	95 (49)	100 (51)	

^aBecause of missing values numbers do not always add up to 721. ^bMeasured by the dichotomized item 'satisfaction with information provision' of the European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire INFO25 (EORTC QLQ-INFO25) questionnaire.

IQR; interquartile range.