

Supplementary Tables have been published as submitted, and have not been copyedited, typeset or checked for scientific content by Acta Dermato-Venereologica

Table SI. Patient Global Assessment of Disease Status (PGADS) questionnaire

Patients with atopic dermatitis

Patient Global Assessment of Disease Status

Considering all the ways your atopic dermatitis has affected you, how do you feel?

- Poor
- Fair
- Good
- Very Good
- Excellent

Table SII. Primary and secondary effectiveness outcomes of 16-week baricitinib treatment stratified by dupilumab non-responders (dup-NR) and dupilumab responders/naïve patients (dupilumab-R/naïve)

	Baseline	Week 4	Week 8	Week 16	P-value^a
Number of patients					-
Dup-NR	N=26	N=26	N=25	N=19	
Dup-R/naïve	N=25	N=25	N=23	N=17	
Patients who discontinued treatment, n (%)					-
Dup-NR	0 (0)	1 (3.8)	6 (23.1)	4 (15.4)	
Dup-R/naïve	0 (0)	2 (8.0)	6 (24.0)	4 (16.0)	
Primary endpoints					
EASI score, mean (95% CI)					0.920
Dup-NR	18.6 (13.4-23.9)	10.1 (7.0-13.2)	8.3 (5.4-11.2)	9.2(4.7-13.8)	
Dup-R/naïve	18.0 (12.6-23.3)	10.2 (7.1-13.4)	11.1 (8.1-14.2)	13.1 (8.4-17.8)	
Mean weekly average NRS-pruritus (95% CI)					0.914
Dup-NR	6.6 (5.8-7.5)	4.3 (3.3-5.4)	4.7 (3.8-5.6)	4.7 (3.6-5.8)	
Dup-R/naïve	6.7 (5.8-7.6)	4.9 (3.9-6.0)	5.2 (4.2-6.1)	6.0 (4.8-7.1)	
Mean weekly average NRS-pain (95% CI)					0.974
Dup-NR	4.3 (3.2-5.4)	2.9 (1.9-4.0)	2.7 (1.7-3.7)	2.8 (1.5-4.1)	
Dup-R/naïve	4.1 (3.0-5.3)	2.8 (1.7-3.9)	2.7 (1.5-3.8)	3.8 (2.4-5.2)	
DLQI score, mean (95% CI)					0.619
Dup-NR	11.9 (9.6-14.3)	6.9 (4.6-9.2)	6.1 (4.1-8.1)	4.7 (3.0-6.4)	
Dup-R/naïve	12.0 (9.6-14.3)	9.2 (6.9-11.5)	8.5 (6.3-10.7)	6.5 (4.7-8.3)	
POEM score, mean (95% CI)					0.954
Dup-NR	17.1 (15.1-19.1)	13.1 (10.4-15.8)	9.8 (7.7-11.9)	12.1 (9.3-14.9)	
Dup-R/naïve	16.7 (14.6-18.8)	14.6 (11.8-17.3)	11.6 (9.5-13.8)	13.2 (10.3-16.0)	
Secondary endpoints					
EASI score ≤7, n(%), probability % (95% CI)					0.246
Dup-NR	5 (19.2), 6.8 (1.2-30.7)	8 (30.8), 19.2 (4.7-53.1)	13 (52.0), 48.9 (16.4-82.3)	11 (57.9), 37.6 (12.2-72.4)	

Dup-R/naïve	5 (20.0), 8.3 (1.6-33.6)	11 (44.0), 34.4 (10.5-70.1)	7 (30.4), 16.3 (2.9-56.2)	8 (47.1), 20.9 (5.3-55.3)	
NRS-pruritus ≤ 4 , n(%), probability % (95% CI)					0.808
Dup-NR	4 (15.4), 6.8 (1.4-27.8)	12 (46.2), 41.4 (16.4-71.8)	10 (40.0), 30.1 (8.5-66.6)	9 (47.4), 27.1 (9.1- 58.0)	
Dup-R/naïve	3 (12.0) 5.1 (0.9-24.3)	9 (36.0), 30.5 (10.5-62.3)	7 (30.4), 18.0 (3.5-57.0)	6 (35.3), 14.8 (3.4-42.4)	
NRS-pain ≤ 4 , n(%), probability % (95% CI)					0.866
Dup-NR	12 (46.2), 45.5 (18.6-75.3)	18 (69.2), 81.9 (50.9-95.2)	18 (72.0), 54.8 (24.9-81.5)	14 (73.7), 78.7 (43.4-94.7)	
Dup-R/naïve	13 (52.0), 51.1 (21.2-80.2)	19 (76.0), 85.0 (56.3-96.1)	17 (73.9), 47.1 (19.8-76.2)	12 (70.6), 79.2 (41.0-95.5)	
DLQI score ≤ 5 , n (%), probability % (95% CI)					0.415
Dup-NR	4 (15.4), 9.1 (2.2-31.2)	11 (42.3), 38.5 (15.9-67.4)	11 (44.0), 36.9 (13.6-68.6)	11 (57.9), 35.6 (13.2-66.8)	
Dup-R/naïve	4 (16.0), 10.0 (2.4-33.2)	8 (32.0), 25.4 (8.6-55.4)	4 (17.4), 9.5 (1.2- 47.3)	8 (47.1), 27.0 (8.7- 58.9)	
POEM score ≤ 7 , n (%), probability % (95% CI)					0.240
Dup-NR	2 (7.7), 1.0 (0.1- 17.6)	3 (11.5), 2.1 (0.1-25.1)	7 (28.0), 10.9 (1.2- 56.4)	4 (21.1), 2.5 (0.2- 30.7)	
Dup-R/naïve	1 (4.0) 0.4 (0.0-13.8)	6 (24.0), 9.8 (1.3-47.3)	2 (8.7), 1.6 (0.4-38.1)	4 (23.5), 4.8 (0.4-17.6)	
PGADS score ≥ 3 , n (%), probability % (95% CI)					0.504
Dup-NR	8 (30.8), 25.1 (7.6-57.9)	15 (57.7), 57.5 (29.2-81.5)	12 (48.0), 46.6 (20.8-74.6)	15 (78.9), 60.2 (34.1-81.6)	
Dup-R/naïve	7 (28.0), 26.2 (8.1-58.6)	12 (48.0), 48.6 (22.7-75.3)	11 (47.8), 40.6 (16.6-70.1)	10 (58.8), 36.5 (15.1-64.9)	
Data after multiple imputation. CI; Confidence Interval; dup-NR: dupilumab non-responders; dup-R/naïve: dupilumab naïve/responders; EASI: Eczema Area and Severity Index; NRS: Numeric Rating Scale; DLQI: Dermatology Life Quality Index; POEM: Patient-Oriented Eczema Measure; PGADS: Patient Global Assessment of Disease Status. ^a P-values based on overall likelihood ratio tests for time.					