

Supplementary material has been published as submitted. It has not been copyedited, or typeset by Acta Dermato-Venereologica

Table SI. Laboratory features of steroid-associated and non-steroid-associated rosacea groups

	<b>HbA1c</b> (%)	<b>Glu</b> (mg/dL)	<b>BUN</b> (mg/dL)	<b>Cr</b> (mg/dL)	<b>cholesterol</b> (mg/dL)	<b>AST</b> (U/L)	<b>ALT</b> (U/L)	<b>rGT</b> (U/L)	<b>CRP</b> (mg/dL)	<b>TG</b> (mg/dL)	<b>ESR</b> (mm/hr)
<b>Non-steroid-associated rosacea (n=254)</b>	5.6	102.2	13.4	1.8	188.4	22.2	20.0	24.2	0.1	132.2	13.5
<b>All steroid-associated rosacea (n=192)</b>	6.4	105.7	13.3	0.7	185.3	23.1	20.7	24.0	0.2	135.2	14.3
<b><i>p</i>-value</b>	0.023*	0.266	0.976	0.247	0.25	0.202	0.113	0.412	0.061	0.429	0.507
<b>Topical steroid-associated rosacea (n=103)</b>	7.0	106.6	13.0	0.7	182.0	22.6	19.5	20.6	0.1	128.4	13.0
<b><i>p</i>-value</b>	0.07	0.476	0.087	0.937	0.1	0.871	0.633	0.36	0.621	0.632	0.945
<b>Systemic steroid-associated rosacea (n=46)</b>	5.7	107.1	13.7	0.8	192.3	23.1	22.7	31.8	0.3	138.2	14.5
<b><i>p</i>-value</b>	0.565	0.663	0.493	0.228	0.574	0.901	0.816	0.456	0.608	0.493	0.522

Data are presented as n (%). BMI, body mass index; SD, standard deviation. \*  $p < 0.05$ .

†Other steroid forms and mixed steroid form used patients were not included for analysis d/t low number of each form and heterogeneity in steroid usage history