

Supplementary material has been published as submitted. It has not been copyedited, or typeset by Acta Dermato-Venereologica

Table SI. OPM distribution in MF patients categorized according to the Netherlands Cancer Registration, which are based on the 10th International Classification of Diseases

OPM	Before MF diagnosis (N)	After MF diagnosis (N)	Total N (%)
Skin*	34	54	88 (23.3%)
<i>cSCC</i>	18	43	61 (16.2%)
<i>Melanoma</i>	15	11	26 (6.9%)
<i>Merkel cell carcinoma</i>	1	0	1
Hematological	43	18	61 (16.2%)
Digestive tract	22	27	49 (13.0%)
<i>Oesophagus</i>	1	2	3
<i>Stomach</i>	2	3	5
<i>Pancreas</i>	1	5	6
<i>Small intestine</i>	0	1	1
<i>Colon</i>	12	11	23
<i>Rectum</i>	6	5	11
Male genital organs	28	20	48 (12.7%)
<i>Prostate</i>	27	19	46
<i>Seminoma</i>	1	1	2
Breast	26	14	40 (10.6%)
<i>Invasive breast cancer</i>	23	13	36
<i>Ductal carcinoma in situ</i>	3	1	4
Urinary tract	10	18	28 (7.4%)
<i>Kidney</i>	1	3	4
<i>Bladder</i>	9	15	24
Respiratory tract	10	14	24 (6.4%)
<i>Small cell</i>	0	2	2
<i>Non-small cell</i>	10	12	22
Head and neck	4	8	12 (3.2%)
<i>Lip</i>	0	2	2
<i>Nasal Cavity</i>	0	1	1
<i>Floor of mouth</i>	0	1	1
<i>Maxillary sinus</i>	1	0	1
<i>Oropharynx carcinoma</i>	1	2	3
<i>Larynx</i>	1	0	1
<i>Supraglottic</i>	0	2	2
<i>Glottis carcinoma</i>	1	0	1
Female genital organs	8	3	11 (2.9%)
<i>Vulva</i>	0	1	1
<i>Corpus uteri</i>	7	1	8
<i>Epithelial ovarian</i>	1	1	1
Primary location unknown	0	7	7 (1.9%)

Bone, articular cartilage and soft tissues	4	1	5 (1.3%)
<i>Bone tumour of the extremities</i>	2	0	2
<i>Soft tissue sarcoma</i>	2	1	3
Endocrine glands	2	0	2 (0.5%)
<i>Thyroid</i>	1	0	1
<i>Endocrine glands, other</i>	1	0	1
Eye	1	0	1 (0.3%)
<i>Melanoma of the eye</i>	1	0	1
CNS	1	0	1 (0.3%)
<i>Cerebral</i>	1	0	1
Total	193	184	377

OPM, Other primary malignancies; cSCC, cutaneous squamous cell carcinoma; CNS, Central Nervous System; MF, Mycosis Fungoides