

**Table SI.** Trichoscopic activity and affected area

|                                                            | LPP<br>(n=161) | FFA<br>(n=154) | Total (n=315) | <i>P</i> value |
|------------------------------------------------------------|----------------|----------------|---------------|----------------|
| Trichoscopic activity, n (%)                               |                |                |               |                |
| First visit                                                |                |                |               |                |
| No                                                         | 14 (9.0%)      | 20 (13.3%)     | 34 (11.1%)    | .062           |
| Moderate                                                   | 131 (84.0%)    | 127 (84.7%)    | 258 (84.3%)   |                |
| Severe                                                     | 11 (7.1%)      | 3 (2.0%)       | 14 (4.6%)     |                |
| Last visit                                                 |                |                |               |                |
| No                                                         | 53 (34.6%)     | 45 (30.6%)     | 98 (32.7%)    | .758           |
| Moderate                                                   | 97 (63.4%)     | 99 (67.3%)     | 196 (65.3%)   |                |
| Severe                                                     | 3 (2.0%)       | 3 (2.0%)       | 6 (2.0%)      |                |
| Progress                                                   |                |                |               |                |
| Worsening                                                  | 1 (0.7%)       | 8 (5.4%)       | 9 (3.0%)      | .036           |
| Stable                                                     | 106 (69.7%)    | 104 (70.7%)    | 210 (70.2%)   |                |
| Improving                                                  | 45 (29.6%)     | 35 (23.8%)     | 80 (26.8%)    |                |
| Affected area (spreading), n (%)                           |                |                |               |                |
| First visit                                                |                |                |               |                |
| Small                                                      | 41 (27.3%)     | 56 (37.8%)     | 97 (32.6%)    | .073           |
| Moderate                                                   | 77 (51.3%)     | 72 (48.6%)     | 149 (50.0%)   |                |
| Large                                                      | 32 (21.3%)     | 20 (13.5%)     | 52 (17.4%)    |                |
| Last visit                                                 |                |                |               |                |
| Small                                                      | 29 (19.6%)     | 53 (36.6%)     | 82 (28.0%)    | .004           |
| Moderate                                                   | 77 (52.0%)     | 64 (44.1%)     | 141 (48.1%)   |                |
| Large                                                      | 42 (28.4%)     | 28 (19.3%)     | 70 (23.9%)    |                |
| Progress                                                   |                |                |               |                |
| Regression                                                 | 1 (0.7%)       | 1 (0.7%)       | 2 (0.7%)      | .020           |
| Stable                                                     | 124 (83.8%)    | 136 (93.8%)    | 260 (88.7%)   |                |
| Progression                                                | 23 (15.5%)     | 8 (5.5%)       | 31 (10.6%)    |                |
| Valid percentages are shown.                               |                |                |               |                |
| LPP, lichen planopilaris; FFA, frontal fibrosing alopecia. |                |                |               |                |