


Supplementary material has been published as submitted. It has not been copyedited, or typeset by Acta Dermato-Venereologica

Lines of treatment used in this cohort of 133 patients with slow-flow vascular malformations

Fig. S1. Number of patients receiving different lines of long-term therapy



APPENDIX 2

Suppl Table 2. Prophylactic treatment prescribed after an infectious episode for four patients with severe or recurrent cellulitis episodes

Prophylaxis	Patient 1	Patient 2	Patient 3	Patient 4
Sex	M	F	M	M
Age at first infection	8 months	5 months	27 months	196 months
Time to antibiotic prophylaxis	After the 3 rd cellulitis episode	After the 2 nd cellulitis episode	After the 3 rd cellulitis episode	After the 1 st cellulitis episode (severe sepsis)
Type of antibiotic	IM benzathine benzylpenicillin	oral phenoxymethylpenicillin	oral phenoxymethylpenicillin	oral phenoxymethylpenicillin
Second choice of antibiotic	Oral cefaclor (for bad tolerance of the first treatment)	Oral cefaclor (after the 4 th episode) then oral amoxicillin-clavulanic acid (after the 5 th episode)	/	oral amoxicillin-clavulanic acid (switch after 6 months)
Antibiotic side-effects reported	none	none	none	none
Total number of cellulitis episodes	5	5	4	2
Other additional treatment	Sirolimus (after the 5 th episode)	Pending alpelisib treatment (BYL719)	/	Sirolimus

M: Male; F: Female; IM: intramuscular