

Table SI. Reasons for therapy discontinuation, according to the specific treatment prescribed or the class of therapy

		N	%
Topical therapies (n=86)	Complete response	2	2.3%
	Partial response	32	37.2%
	No Response	4	4.7%
	AEs	3	3.5%
	Other reasons	1	1.2%
	na	44	51.2%
Topical antibiotics (n=19)	Partial response	10	52.6%
	No Response	1	5.3%
	na	8	42.1%
Topical disinfectants (n=17)	Partial response	2	11.8%
	No Response	1	5.9%
	na	14	82.4%
Topical cortisone (n=22)	Complete response	1	4.5%
	Partial response	12	54.5%
	No Response	1	4.5%
	Other reasons	1	4.5%
	na	7	31.8%
Topical dapsone (n=10)	Partial response	7	70.0%
	na	3	30.0%
Topical isotretinoin (n=4)	AEs	3	75.0%
	na	1	25.0%
Local injection of cortisone (n=1)	na	1	100.0%
Other (n=13)	Complete response	1	7.7%
	Partial response	1	7.7%
	No Response	1	7.7%
	na	10	76.9%
Systemic antibiotics (n=44)	Complete response	4	9.1%
	Partial response	28	63.6%
	No Response	5	11.4%
	AEs	2	4.5%
	Other reasons	1	2.3%
	na	4	9.1%
Rifampicin + Clindamycin (n=15)	Partial response	10	66.7%
	AEs	2	13.3%
	Other reasons	1	6.7%
	na	2	13.3%
Tetracyclines (n=28)	Complete response	4	14.3%
	Partial response	17	60.7%
	No Response	5	17.9%
	na	2	7.1%
Other (n=1)	Partial response	1	100.0%

		N	%
Systemic steroids (n=5)	Partial response	2	40.0%
	AEs	1	20.0%
	na	2	40.0%
Prednisolone (n=1)	na	1	100.0%
Other (n=4)	Partial response	2	50.0%
	AEs	1	25.0%
	na	1	25.0%
Other systemic therapies (n=26)	Partial response	11	42.3%
	No Response	4	15.4%
	AEs	2	7.7%
	Other reasons	1	3.8%
	na	8	30.8%
Acitretin (n=2)	No Response	1	50.0%
	AEs	1	50.0%
Isotretinoin (n=8)	Partial response	7	87.5%
	AEs	1	12.5%
Dapsone (n=2)	Partial response	1	50.0%
	Other reasons	1	50.0%
Mycophenolate Mofetil (n=1)	No Response	1	100.0%
Zinc salts (n=1)	Partial response	1	100.0%
Vitamins/supplements (n=1)	na	1	100.0%
Other (n=11)	Partial response	2	18.2%
	No Response	2	18.2%
	na	7	63.6%

AEs: adverse events, na: not available

Multiple treatment courses are possible for each patient.