

Table SII. Demographics and disease characteristics of patients who became partial responders after achieving PASI 90 at week 16

	PASI90R who maintained PASI 90 response after week 16 (n = 320)	PASI90R who became partial responders at any time-point after week 16 (n = 17)	p-value
Age, years, mean (SD)	43.41 (12.99)	51.12 (13.51)	0.02
Age ≥ 65 years, n (%)	22 (6.88)	3 (17.65)	0.12
Male, n (%)	234 (73.13)	12 (70.59)	0.78
Weight, kg, mean (SD)	78.81 (16.34)	90.72 (17.04)	0.004
Body mass index, kg/m ² , mean (SD)	26.60 (5.06)	30.83 (5.05)	<0.001
Body mass index ≥ 25 kg/m ² , n (%)	188 (58.75)	13 (76.47)	0.21
Waist, cm, mean (SD)	94.21 (14.66)	106.59 (14.03)	<0.001
Nail psoriasis, n (%)	120 (37.50)	8 (47.06)	0.43
Scalp psoriasis, n (%)	254 (79.38)	10 (58.82)	0.05
Age at diagnosis, years, mean (SD)	26.07 (13.59)	29.68 (16.76)	0.49
Time since diagnosis, years, mean (SD)	17.92 (11.95)	22.03 (12.26)	0.14
Psoriatic arthritis, n (%)	50 (15.63)	4 (23.53)	0.49
PASI at baseline, mean (SD)	21.29 (9.55)	22.68 (11.15)	0.56
Metabolic syndrome, n (%)	40 (12.50)	5 (29.41)	0.06

A patient is considered a partial responder if, at any time-point after week 16, they achieve an improvement (reduction) in PASI score compared with baseline between 50% and <75%, and therefore lose the PASI 90 response reached at week 16.

PASI90R: PASI 90 responders at week 16; PASI: Psoriasis Area and Severity Index; SD: standard deviation.

Those values that are statistically significant are highlighted in bold.