

Supplementary material has been published as submitted. It has not been copyedited, typeset or checked for scientific content by Acta Dermato-Venereologica

Table SI. PRO separated by pruritus groups according to the classification of itch (IFSI-groups) before and after the treatment start in a specialized pruritus care unit

	Before treatment start (T0)		After treatment start (T1)	
		p-value		p-value
NRS-average 24 h, mean±SD (median)				
IFSI I	5.35± 2.37 (5.0)		3.44±2.67 (3.0)	
IFSI II	4.83± 2.54 (5.0)		3.76±2.94 (3.0)	
IFSI III	6.23± 2.44 (7.0)	0.003**	3.75±2.72 (4.0)	0.73
DLQI, mean±SD (median)				
IFSI I	10.14±7.40 (8.0)		6.14±6.02 (4.0)	
IFSI II	7.37±5.57 (6.0)		5.04±4.92 (3.0)	
IFSI III	9.84±5.91 (9.0)	0.006**	6.07±5.26 (5.0)	0.34
ItchyQol, mean±SD (median)				
IFSI I	68.20±16.94 (68.2)		56,10±20.46 (57.2)	
IFSI II	59.40±16.06 (56.1)		52.8±17.38 (55.0)	
IFSI III	73.48±14.96 (74.8)	<0.001***	62.92±20.02 (67.1)	0.008**
PBI-P, mean±SD (median)				
IFSI I	1.02±1.20 (0.6)		2.13±1.36 (2.3)	
IFSI II	1.33±1.36 (1.0)		2.08±1.35 (2.2)	
IFSI III	1.14±1.30 (0.7)	0.49	2.06±1.25 (1.8)	0.97

*p<0.05; **p<0.01; ***p<0.001 (Wilcoxon sign-rank test)

NRS: numeric rating scale; VRS: verbal rating scale; DLQI: Dermatology Life Quality Index; ItchyQol: pruritus specific quality of life measure, PBI-P: Patient Benefit Index for Pruritus