

Table SI. Cases of ichthyosis linearis circumflexa (ILC) without congenital erythroderma and trichorrhexis invaginata (TI)^a

Case (Ref)	Age, years/sex	ILC	TI	Atopy
1. Nabai and Mehregan (4)	2/F	Yes	No	Not reported
2. Manabe, et al. (19)	18/M	Yes	No	Increased total serum IgE (3,600 IU/ml)
3. Manabe, et al. (19)	Not reported/F	Yes	Not reported	Not reported
4. Mathys, et al. (5)	24/M	Yes	No	No
5. Tosti, et al. (7)	14/F	Yes	No	Atopic dermatitis. Increased total (600 IU/ml) and specific (milk, eggs) serum IgE
6. Tosti, et al. (7)	15/F	Yes	No	No
7. Tomidei, et al. (6)	46/F	Yes	No	No
8. Malakar, et al. (8)	9/M	Yes	No	No
9. Gambichler, et al. (3)	13/M	Yes	No	Atopic dermatitis. Increased total (347 IU/ml) and specific (ragweed, birch, rye flour, house dust mite) serum IgE
10. Gambichler, et al. (3)	6/F	Yes	Not reported	Not reported
11. Suga, et al. (2)	20/M	Yes	No	No
12. Kumar, et al. (9)	7/F	Yes	No	Not reported

^aAt least 16 additional cases have been previously reviewed by Mevorah et al. (20).