

Table SII. Intraepidermal nerve fibre density (IENFD (fibres/mm)) in different stages of prurigo nodularis stratified according to the whole collective (n = 30) and patients with biopsy obtained from same localization (n = 14)

Localization	Patients <i>n</i>	IENFD (fibres/mm), mean ± SD (median)			
		Lesional skin (LS)	Inter-lesional skin (ILS)	Non-lesional skin (NLS)	Healed skin (HS)
Whole collective					
Arm	14	7.43 ± 4.09 (6.03)	6.24 ± 3.86 (5.53)	10.26 ± 5.93 (10.06)	14.16 ± 7.17 (17.1)
Leg	8	3.07 ± 3.53 (1)	2.92 ± 1.26 (3.38)	5.60 ± 3.53 (5.09)	7.29 ± 4.6 (6.65)
Trunk	8	11.81 ± 7.72 (9.19)	13.30 ± 8 (11.4)	17.44 ± 9.29 (15.7)	13.35 ± 6.77 (16.8)
Total	30	7.43 ± 5.95 (6.03)	7.83 ± 6.69 (5.53)	11.39 ± 7.79 (10.9)	12.25 ± 6.84 (14.5)
Identical biopsy area in each individual patient					
Arm	6	8.47 ± 5.52 (5.5)	5.31 ± 2.84 (5.39)	9.44 ± 4.44 (10.06)	15.71 ± 5.52 (17.75)
Leg	4	2.73 ± 3.92 (0.88)	2.49 ± 1.35 (2.43)	5.86 ± 4.02 (5.71)	7.57 ± 5.27 (6.94)
Trunk	4	10.45 ± 4.54 (8.31)	14.14 ± 10.78 (9.9)	15.72 ± 6.84 (13.55)	15.63 ± 6.85 (16)
Total	14	7.40 ± 5.49 (5.5)	7.03 ± 7.32 (4.34)	10.21 ± 6.13 (10.06)	13.36 ± 6.59 (14.55)