

Table SII. Description of the cases included and results obtained in the 3 prior studies and the present study of shiny white streaks (SWS)

	Balagula et al. (25)	Di Stefani et al. (22)	Liebman et al. (26)	Present study
Lesions evaluated, n	11,225	800	538	800
Lesions with histopathological diagnosis, n	494	502	538	800
SWS in evaluated lesions, %	1.8	16.1	38.7	13.4
Melanomas, n	229	400	154	125
SWS melanoma, %	28.4	31.0	23.4	32.8
SWS invasive melanoma, %	41.0	n.s.	n.s.	50.7
SWS <i>in situ</i> melanoma, %	17.0	n.s.	n.s.	10.7
OR invasive/ <i>in situ</i> (95% CI, $p < 0.05$ )	3.4 (1.9–6.3)	n.s.	n.s.	10.3 (3.8–28.0)
BCC, n	39	n.s.	191	133
SWS in BCC, %	47.6	n.s.	42.4	30.8
OR malignant/benign (95% CI, $p < 0.005$ )	2.5 (1.5–4.3)	n.s.	n.s.	10.5 (6.4–17.5)
Naevi (biopsied naevi), n	34	102	n.s.	305
SWS naevi, %	0.07	1.3	n.s.	1.6

n.s. not specified in the article; OR: odds ratio; CI: confidence interval; BCC: basal cell carcinoma.