

Table SI. Pathological and surgical features of non-pigmented and pigmented basal cell carcinomas (PBCC)

Characteristics	Non-PBCC (n= 59)	PBCC (n= 199)	p-value
Histologic subtypes, n (%)			0.006
Nodular or nodulocystic type, n (%)	20 (33.9)	96 (48.2)	0.052
Superficial type	2 (3.3)	10 (5.0)	0.600
Morpheic type	4 (6.8)	12 (6.0)	0.834
Infiltrative type	18 (30.5)	24 (12.1)	0.001
Micronodular type	2 (3.4)	25 (12.6)	0.043
Mixed type	13 (22.1)	32 (16.1)	0.290
Mohs stage, mean ± SD	2.34 ± 1.15	1.80 ± 0.98	0.001
Mohs stage, n (%)			<0.001
1 <sup>st</sup>	19 (32.3)	92 (46.2)	
2 <sup>nd</sup>	14 (23.7)	74 (37.2)	
3 <sup>rd</sup>	13 (22.0)	19 (9.6)	
≥4 <sup>th</sup>	13 (22.0)	14 (7.0)	
Surgical margin, mm, mean ± SD	6.97 ± 2.15	6.28 ± 2.04	0.026
Surgical margin, n (%)			0.021
≤5 mm	26 (44.1)	114 (57.3)	
6–10 mm	16 (27.1)	60 (30.2)	
11–15 mm	9 (15.2)	16 (8.0)	
≥16 mm	8 (13.6)	9 (4.5)	
Marginal area, mm <sup>2</sup> , mean ± SD	194.4 ± 179.9	128.6 ± 108.7	0.001
Marginal area, n (%)			0.005
≤100 mm <sup>2</sup>	24 (40.7)	97 (48.7)	
101–200 mm <sup>2</sup>	16 (27.1)	77 (38.7)	
201–400 mm <sup>2</sup>	11 (18.6)	16 (8.1)	
≥401 mm <sup>2</sup>	8 (13.6)	9 (4.5)	
Surgical depth, n (%)			0.036
Subcutaneous layer	24 (40.7)	115 (57.8)	0.021
Muscle fascia/muscular layer	13 (22.0)	40 (20.1)	0.747
Perichondrium/cartilage	22 (37.3)	44 (22.1)	0.019