

Table SII. Clinical features of idiopathic bullous pemphigoid (BP) group

Case	Age, years	Severity score			Steroid dosage (mg/kg/day)	
		BPDAI	ABSIS	Acral severity (0–8)	Peak dose	Lowest maintenance dose
1	63	3	3.00	0	0.19	0.03
2	85	21	7.50	1	0.43	NA
3	76	30	7.00	1	0.14	0.02
4	82	23	5.00	1	0.00	NA
5	50	69	17.00	5	0.68	0.04
6	74	1	0.75	0	0.06	0.00
7	85	16	7.00	0	0.21	0.12
8	78	5	3.15	1	0.57	0.05
9	22	43	15.50	1	0.80	0.00
10	74	12	6.30	2	0.93	0.28
11	43	51	15.00	1	0.64	0.00
12	79	37	8.50	3	0.98	0.10
13	82	4	1.50	0	0.26	0.04
14	92	27	4.75	2	0.21	0.21
15	62	47	7.50	0	0.47	0.08
16	69	8	2.00	0	0.31	0.04
17	82	6	3.50	2	0.35	0.09
18	95	31	8.75	1	0.25	0.08
19	88	14	2.00	1	0.13	0.04
20	83	2	0.50	0	0.11	0.05
21	74	18	3.50	0	0.23	0.01
22	55	20	3.00	0	0.32	0.02
23	76	3	0.50	0	0.15	0.00
24	66	17	2.25	1	0.42	NA
25	54	5	1.00	1	0.35	0.01
26	70	12	5.80	0	0.14	NA
27	65	2	1.00	0	0.00	0.00
28	50	1	0.30	0	1.01	0.05
29	72	3	0.20	0	0.00	0.00
30	82	34	9.25	5	0.65	NA
31	88	14	2.75	2	0.42	0.10
32	85	5	0.30	1	0.16	0.08
33	98	58	20.50	5	0.66	0.17
34	86	4	1.50	2	0.22	0.11
35	79	20	4.50	3	0.15	0.08
36	75	3	0.30	1	0.16	NA

NA: patients with insufficient follow-up to decide the maintenance dose (values were not included in the analysis); BPDAI: Bullous Pemphigoid Disease Area Index; ABSIS: Autoimmune Bullous Skin Disorder Intensity Scores.