

Table SII. Clinical characteristics of study population

Patient groups	ASUM	Venous	Vasculitis	PG	Control N	Control H
Age at diagnosis, years, mean (range)	72.2 (50–87)	69.7 (47–88)	59.1 (41–83)	52.8 (22–74)	63.1 (50–79)	70.7 (48–92)
Sex, <i>n</i>						
Female	33 (49.3)	12 (80)	5 (33.3)	10 (66.6)	9 (60)	10 (66.6)
Male	34 (50.7)	3 (20)	10 (66.6)	5 (33.3)	6 (40)	5 (33.3)
Arterial hypertension, <i>n</i>						
Yes	67 (100)	7 (46.7)	6 (40)	5 (33.3)	0 (0)	15 (100)
No	0 (0)	8 (53.3)	9 (60)	14 (66.6)	15 (100)	0 (0)
Vitamin K Antagonist, <i>n</i>						
Yes	24 (35.8)	0 (0)	2 (13.3)	1 (6.7)	0 (0)	0 (0)
No	43 (64.7)	15 (100)	13 (86.7)	14 (93.3)	15 (100)	15 (100)
Diabetes mellitus II, <i>n</i>						
Yes	33 (49.3)	0 (0)	7 (46.7)	2 (13.3)	0 (0)	1 (6.7)
No	34 (50.7)	15 (100)	8 (53.3)	13 (86.7)	15 (100)	14 (93.3)
Renal insufficiency, <i>n</i>						
Yes	12 (17.9)	0 (0)	2 (13.3)	0 (0)	0 (0)	0 (0)
No	55 (82.1)	15 (100)	13 (86.7)	15 (100)	15 (100)	15 (100)
PAOD, <i>n</i>						
Yes	12 (17.9)	2 (13.3)	3 (20)	2 (13.3)	0 (0)	0 (0)
No	55 (82.1)	13 (86.7)	12 (80)	13 (86.7)	15 (100)	15 (100)
CVD, <i>n</i>						
Yes	16 (23.9)	15 (100)	3 (20)	2 (13.3)	1 (6.7)	0 (0)
No	51 (76.1)	0 (0)	12 (80)	13 (86.7)	14 (93.3)	15 (100)
Size of ulceration, cm, mean (range)	7.13 (2–20)	8.55 (1.5–28)	3.94 (0.5–20)	5.35 (1–16)	–	–
Duration of disease, months, mean (range)	7.75 (3–15.5)	16.17 (4–44)	4.25 (0.5–28)	3.73 (1–11.5)	–	–

PAOD: peripheral artery occlusive disease; CVD: chronic venous disease; ASUM: arteriolosclerotic ulcer of Martorell; PG: pyoderma gangraenosum; Control N: non-hypertensive controls; Control H: hypertensive controls.