

Urticaria, Angioedema and Dyspnoea in Adjuvant Therapy of Melanoma with Interferon α -2b

Doris Helbig, Jan Christoph Simon and Regina Treudler

University of Leipzig, Department for Dermatology, Venereology and Allergology, Philipp-Rosenthal-Str. 23, DE-04103 Leipzig, Germany. E-mail: doris.helbig@medizin.uni-leipzig.de

Accepted November 19, 2009.

Adjuvant therapy with interferon (IFN) α -2b has been shown to increase disease-free survival in patients with stage II (3 million IU, 3 times/week, subcutaneous: low-dose) or stage III melanoma (20 million IU/m² body surface area, intravenous 5 times/week: high-dose) without improving overall survival (1). The most frequently reported cutaneous adverse drug reactions (ADR) reported in a group of 33 patients on adjuvant low-dose IFN α -2b therapy because of malignant melanoma were alopecia (48%), hair discoloration (18%), eczematous reactions (39%) and pruritus (30%) (2). Urticaria was reported in only one of these patients (3%) (2). In most reported cases, cutaneous ADR to IFN α -2b do not necessitate cessation of therapy. There are no reports in the literature of urticaria during high-dose IFN α -2b.

CASE REPORT

We report here the case of a 48-year-old man with no history of allergy, who was started on adjuvant high-dose therapy with IFN α -2b because of stage IIb melanoma. On day one, he was treated with IFN α -2b (40 million IU) intravenously, after premedication with 1 g oral paracetamol 1 h previously. Approximately 1 h after the first infusion, he developed slight oedema of the face, which resolved without any further therapy. As an intolerance reaction to paracetamol was suspected, the premedication was changed to ibuprofen on day two. However, within 20 min of the second infusion (after only approximately 10% of the amount had been applied) the patient developed urticarial exanthema and angioedema of both hands, which disappeared within one hour after stopping the infusion and oral antihistamines. Polyvalent intolerance to analgesics was suspected.

Three weeks later, skin-prick tests were negative for a series of analgesics (including paracetamol, diclofenac and ibuprofen), latex and for IFN α -2b. Total IgE, specific IgE for latex and serum tryptase were within normal ranges. Oral provocation testing with ibuprofen was well tolerated, as was paracetamol on another occasion.

On week 4, 1 million IU IFN α -2b was applied subcutaneously after premedication with ibuprofen, which was well tolerated. The following day, 10 min after application of 3 million IU IFN α -2b subcutaneously, the patient again developed generalized urticaria with dyspnoea. Premedication had not been changed. The immediate type symptoms required application of

methylprednisolone and dimetindene intravenously. Therapy with IFN α -2b has not been restarted since.

DISCUSSION

We attributed the patient's immediate type symptoms to application of IFN α . The differential diagnosis included intolerance to analgesics or latex allergy, which could be excluded by *in vitro* and *in vivo* testing, including oral provocation test to analgesics. There were no signs of underlying mastocytosis.

The pathogenesis of the reaction in our patient is not clear. Interestingly, the patient showed a spontaneously resolving facial oedema after 40 million IU IFN α -2b intravenous, but urticarial exanthema and angioedema after the second infusion and after administration of only 3 million IU IFN α -2b subcutaneously. One might speculate that this was due to an immunological sensitization, which was, however, not shown by skin prick test, although non-immunological hypersensitivity might also be suspected.

IFNs are anti-viral and immunoregulatory agents that may increase the activity of macrophages, B- and T cells, as well as of mast cells (3–5). The response of mast cells to IFN might be differentially regulated intra- and inter-individually. This is emphasized by the fact that high concentrations of IFN α -2b were shown to give rise to an IgE-mediated histamine release from basophil granulocytes from patients without formerly known urticaria, but not in a group with pre-existing urticaria (6).

Immediate-type reactions to IFN α have been reported occasionally, i.e. in a patient with hepatitis C who developed angioedema after 22 weeks of IFN α -2b 6 million IU/week (7, 8). In this case, the pathogenesis of IFN-related angioedema was suggested to be similar to that of angioedema induced by angiotensin-converting enzyme (ACE) inhibitors, because there was an increase in the plasma bradykinin level and the onset was delayed after the start of IFN therapy. *In vitro* investigations demonstrated a release of IFN γ and macrophage migration inhibition factor by peripheral blood lymphocytes of patients with drug-induced urticaria, indicating immune sensitization (9). Furthermore, IL1-induced IL8 production was dramatically suppressed by the administration of IFN α -2b (10).

Specific desensitization, as reported with IFN- β in a patient with multiple sclerosis (11), was also considered, but finally not realized in our patient because of

unknown pathogenesis of reaction and lack of compliance.

In summary, we report here a rare case of immediate-type reaction to IFN α -2b in a patient with melanoma, which led to discontinuation of adjuvant therapy with IFN α -2b.

REFERENCES

1. Garbe C, Hauschild A, Volkenandt M, Schadendorf D, Stolz W, Kortmann RD, et al. Deutsche Leitlinie: Malignes Melanom. In: Garbe C (Hrsg.). Interdisziplinäre Leitlinien zur Diagnostik und Behandlung von Hauttumoren. Stuttgart: Georg Thieme Verlag, 2005: p. 23–55.
2. Guillot B, Blazquez L, Bessis D, Dereure O, Guilhou JJ. A prospective study of cutaneous adverse events induced by low-dose alpha-interferon treatment for malignant melanoma. *Dermatology* 2004; 208: 49–54.
3. Mannering GJ, Deloria LB. The pharmacology and toxicology of the interferons: an overview. *Ann Rev Pharmacol Toxicol* 1986; 26: 455–515.
4. Siegel DS, Le J, Vilcek J. Modulation of lymphocyte proliferation and immunoglobulin synthesis by interferon-gamma and “type I” interferons. *Cell Immunol* 1986; 101: 380–390.
5. Knop J, Taborski B, DeMaeyer-Guignard J. Selective inhibition of the generation of T suppressor cells of contact sensitivity in vitro by interferon. *J Immunol* 1987; 138: 3684–3687.
6. Torrelo A, Harto A, Ledo A. Interferon therapy for chronic urticaria. *J Am Acad Dermatol* 1995; 32: 684–685.
7. Marcellin P, Pouteau M, Martinot-Peignoux M, Degos F, Duchatelle V, Boyer N, et al. Lack of benefit of escalating dosage of interferon alfa in patients with chronic hepatitis C. *Gastroenterology* 1995; 109: 156–165.
8. Pardini S, Bosincu L, Bonfigli S, Dore F, Longinotti M. Anaphylactic-like syndrome in systemic mastocytosis treated with alpha-2-interferon. *Acta Haematol* 1991; 85: 220.
9. Livni E, Lapidot M, Halevy S. In vitro release of interferon-gamma and macrophage migration inhibition factor in drug-induced urticaria and angioedema. *Acta Derm Venereol* 1999; 79: 18–22.
10. Reznikov LL, Puren AJ, Fantuzzi G, Muhl H, Shapiro L, Yoon DY, et al. Spontaneous and inducible cytokine responses in healthy humans receiving a single dose of IFN-alpha2b: increased production of interleukin-1 receptor antagonist and suppression of IL-1-induced IL-8. *J Interferon Cytokine Res* 1998; 18: 897–903.
11. Kalpaklioglu AF, Baccioglu KA, Erdemoglu AK. Desensitization in interferon-beta1a allergy: A case report. *Int Arch Allergy Immunol* 2009; 149: 178–180.