

Supplementary material has been published as submitted. It has not been copyedited, typeset or checked for scientific content by Acta Dermato-Venereologica

**Table SI. Comparison of the attitude to photographs of people with skin disease of high school students with or without skin disease, of witnesses of bullying and non-witnesses of bullying and of males and females**

Number of photo	Person on photo	Mean attitude before presentation of patients' stories in students		Mean attitude after presentation of patients' stories in students		Mean attitude before presentation of patients' stories in students		Mean attitude after presentation of patients' stories in students		Mean attitude before presentation of patients' stories in students		Mean attitude after presentation of patients' stories in students	
		With skin disease (n=157)	Without skin disease (n=223)	With skin disease (n=157)	Without skin disease (n=223)	In witnesses of bullying (n=121)	In non-witnesses of bullying (n=257)	In witnesses of bullying (n=121)	In non-witnesses of bullying (n=257)	In males (n=169)	In females (n=200)	In males (n=169)	In females (n=200)
1	Male (control)	2.37±0.98	2.39±0.98	2.22±0.99	2.21±1.03	2.34±1.01	2.40±0.97	2.18±1.07	2.23±0.97	2.30±0.99	2.49±0.95	2.18±1.02	2.24±0.99
2	Female (severe acne)	2.75±1.00	3.05±0.96**	2.39±0.96	2.57±1.04	2.76±1.09	2.98±0.94	2.17±1.05	2.61±0.94***	3.92±0.91	2.63±0.93***	2.88±0.94	2.17±0.92***
3	Female (control, overweight)	2.15±1.02	2.44±1.04**	2.08±1.03	2.24±1.05	2.07±1.09	2.41±1.08**	2.01±1.05	2.22±1.01	2.73±1.06	1.98±0.88***	2.54±1.03	1.86±0.90***
4	Female (vitiligo, African origin)	2.06±1.01	2.32±1.10*	1.86±0.97	2.06±1.02	1.98±1.07	2.30±1.06**	1.69±0.93	2.07±1.00***	2.73±0.99	1.80±0.93***	2.43±0.99	1.59±0.82***
5	Female (control)	2.11±0.97	2.01±0.92	2.06±1.01	2.05±0.93	2.05±1.00	2.05±0.52	1.97±0.97	2.10±0.95	2.12±0.92	2.02±0.98	2.11±0.95	2.03±1.00
6	Female (epidermolysis bullosa, open clothes)	2.26±1.01	2.50±1.06*	2.17±0.99	2.23±1.04	2.25±1.04	2.45±1.06	1.91±1.00	2.31±0.99***	2.81±0.95	2.04±0.95***	2.61±0.95	1.85±1.01***
7	Female (control, unusual nose proportions, open clothes)	2.24±1.01	2.22±1.03	2.17±1.04	2.17±1.02	2.17±1.06	2.24±1.01	2.03±1.10	2.22±0.98	2.38±0.98	2.12±1.02*	2.36±0.99	2.03±1.00**

8	Female (vitiligo, open clothes)	2.29±1.09	2.36±1.04	2.11±1.02	2.13±0.98	2.09±1.06	2.42±1.03**	1.93±1.03	2.20±0.97**	2.67±1.01	2.06±1.02***	2.43±0.90	1.87±0.97***
9	Female (control)	1.99±1.03	1.96±0.95	1.96±0.97	1.97±0.98	1.84±1.03	2.01±0.94	1.86±1.01	1.98±0.95	2.04±0.96	1.92±0.99	2.03±0.98	1.89±0.96
10	Female (alopecia universalis)	1.85±0.99	2.14±1.15**	1.84±1.00	2.00±1.16	1.77±1.01	2.11±1.13**	1.64±1.06	2.01±1.09***	2.41±1.14	1.69±0.94***	2.27±1.11	1.63±0.98***
11	Woman (control)	1.94±0.94	1.98±0.97	1.90±0.95	1.89±0.94	1.92±1.00	1.98±0.93	1.83±0.99	1.91±0.91	2.04±0.95	1.63±0.98	1.98±0.99	1.82±0.89
12	Male (control, mild acne)	2.24±1.07	2.29±1.01	2.13±1.02	2.11±0.97	2.23±1.11	2.31±0.98	2.07±1.05	2.16±0.97	2.22±1.04	2.35±1.04	2.09±0.95	2.17±1.02

\*-  $p < 0.05$ ; \*\*-  $p < 0.01$ ; \*\*\*-  $p < 0.001$