

Table SI. Patients with cutaneous T-cell lymphoma (CTCL) treated with bexarotene

Patient	Age ^a , years	Sex	UV therapy ^b	Radiation therapy ^b	Systemic therapy ^b	Staging	Medical history	Other malignancies	CTCL type
1	59	F	No	No	No	IA	No	No	Follicular
2	53	M	Yes	No	No	IA	No	No	MF
3	51	M	Yes	Yes	Yes	IA	No	No	Follicular
4	85	M	No	No	No	IA	Yes	Yes	Follicular
5	39	F	Yes	No	No	IA	No	No	Follicular
6	87	F	No	No	No	IA	Yes	No	MF
7	57	M	Yes	No	No	IA	No	No	MF
8	50	M	Yes	No	No	IA	No	No	MF
9	23	F	Yes	Yes	Yes	IA	No	No	Follicular
10	65	F	Yes	No	No	IA	Yes	No	MF
11	66	F	No	No	No	IA	Yes	Yes	MF
12	69	M	Yes	No	Yes	IA	No	No	MF
13	80	M	No	No	No	IB	Yes	No	Follicular
14	78	F	No	No	Yes	IB	Yes	No	Follicular
15	70	M	Yes	Yes	No	IB	No	No	CD30+
16	85	M	Yes	No	No	IB	Yes	Yes	MF
17	71	F	Yes	No	No	IB	Yes	Yes	MF
18	51	M	Yes	No	No	IB	Yes	No	MF
19	73	F	Yes	No	Yes	IB	Yes	No	MF
20	56	F	Yes	No	No	IB	No	No	MF
21	60	M	No	No	No	IB	Yes	No	MF
22	64	M	Yes	No	No	IB	Yes	No	MF
23	70	M	No	No	No	IB	Yes	No	Follicular
24	61	F	Yes	No	Yes	IIA	Yes	No	MF
25	58	F	No	Yes	No	IIA	Yes	Yes	MF
26	65	M	Yes	No	No	IIA	Yes	Yes	MF
27	54	M	Yes	No	No	IIB	Yes	No	MF
28	49	M	Yes	Yes	Yes	IIB	No	Yes	MF
29	28	M	Yes	Yes	Yes	IIB	No	Yes	MF
30	60	F	No	Yes	Yes	SPTL	Yes	No	SPTL
31	66	F	No	No	Yes	SPTL	Yes	No	SPTL
32	61	M	No	No	Yes	SS	Yes	Yes	SS
33	76	F	No	No	No	SS	Yes	No	SS
34	66	F	Yes	No	Yes	SS	Yes	Yes	SS
35	52	M	Yes	No	Yes	SS	No	No	SS
36	72	M	No	Yes	Yes	^a	Yes	No	Peripheral T-cell lymphoma ^a
37	61	M	Yes	Yes	?Yes	IB	No	No	MF

^aAt the time of CTCL diagnosis.^bBefore bexarotene therapy.

UV: ultraviolet; MF: mycosis fungoides; SPTL: subcutaneous panniculitis-like T-cell lymphoma.