

Table SII. Side-effects associated with bexarotene treatment

| Side-effect                                 | Patients<br>n (%) | Grade of<br>side-effect <sup>a</sup> |
|---|-------------------|--------------------------------------|
| Hypertriglyceridaemia                       | 19 (51)           | I–IV                                 |
| No side-effects                             | 9 (24)            | –                                    |
| Hypercholesterolaemia                       | 7 (19)            | I–II                                 |
| Leucopaenia/neutropaenia                    | 5 (14)            | II                                   |
| Elevation of liver transaminases (ALT, AST) | 4 (11)            | II–IV                                |
| Hypothyroidism                              | 4 (11)            | I                                    |
| Anaemia                                     | 3 (8)             | I–II                                 |
| Skin symptoms, rash, flush                  | 3 (8)             | I                                    |
| Hypoglycaemia                               | 2 (5)             | II                                   |
| Creatinine kinase (CK) elevation            | 1 (3)             | II                                   |
| Thrombocytopaenia                           | 1 (3)             | II                                   |
| Gastrointestinal symptoms, e.g. nausea      | 1 (3)             | I                                    |
| Other                                       | 1 (3)             | –                                    |

<sup>a</sup>National Cancer Institute common terminology criteria for adverse events.  
ALT: alanine aminotransferase; AST: aspartate aminotransferases.