

Table SI. *Gonorrhoea* patient characteristics: sex, age, sample site, test of cure (TOC), initial method of diagnosis, and result of antimicrobial susceptibility testing

Sex	Age, years	Sample site	Days to cure (1 st or 2 nd TOC)		Initial method of diagnosis		MIC, mg/l	
			Days to 1 st TOC	2 nd TOC)	PCR	Culture	Ciprofloxacin	Ceftriaxone
M	47	Rectum			Neg			
		Pharynx			Neg			
		Urethra	6	6	Pos	Pos	32	0.047
M	22	Rectum	6	6	Pos	Neg		
		Pharynx			Neg			
		Urethra	6	11	Pos	Pos	3	0.023
M	40	Urethra	7	7	Pos	Pos	2	0.002
M	22	Rectum	7	7	Pos	Pos	0.004	0.003
		Pharynx	7	7	Pos	Neg		
		Urethra	7	7	Pos	Pos	0.004	0.003
M	29	Urethra	7	7	Pos	Pos	0.003	0.004
M	43	Urethra	7	7	Pos	Pos	32	0.004
M	43	Urethra	7	7	Pos	Pos	1	0.004
M	39	Urethra	7	7	Pos	Pos	32	0.047
M	53	Rectum			Neg			
		Pharynx			Neg			
		Urethra	7	7	Pos	Pos	0.004	0.003
M	28	Urethra	7	7	Pos	Pos	8	0.047
M	51	Urethra	7	7	Pos	Pos	3	0.032
M	34	Urethra	7	7	Pos	Pos	3	0.008
F	28	Cervix	7	7	Pos	Pos	0.38	0.003
		Urethra	7	7	Pos	Pos	Not performed	Not performed
M	54	Urethra	7	7	Pos	Pos	Not performed	Not performed
M	22	Rectum			Neg			
		Pharynx			Neg			
		Urethra	7	7	Pos	Pos	0.002	0.002
M	29	Urethra	7	7	Pos	Pos	0.002	0.003
M	58	Urethra	7	7	Pos	Pos	Not performed	Not performed
M	49	Urethra	8	8	Pos	Neg		
M	35	Urethra	8	8	Pos	Pos	32	0.016
M	34	Urethra	8	8	Pos	Pos	0.002	0.002
M	37	Urethra	8	8	Pos	Pos	6	0.006
F	21	Cervix	8	8	Pos	Pos	3	0.004
		Urethra	8	8	Pos	Neg		
M	46	Rectum			Neg	Pos		
		Pharynx			Neg			
		Urethra	11	11	Pos	Pos	0.002	0.002
M	25	Urethra	11	11	Pos	Pos	1.5	0.002
M	29	Rectum			Neg			
		Pharynx			Neg			
		Urethra	4	11	Pos	Pos	0.006	0.25
M	44	Urethra	12	12	Pos	Pos	0.38	0.003
M	27	Rectum			Neg			
		Pharynx	12	12	Pos	Neg		
M	42	Rectum	13	13	Pos	Neg		
		Pharynx			Neg			
		Urethra	13	13	Pos	Pos	32	0.004
M	39	Rectum	14	14	Pos	Neg		
M	27	Rectum			Neg			
		Pharynx			Neg			
		Urethra	7	19	Pos	Pos	32	0.047

EUCAST (www.eucast.org) breakpoints were applied.

Neg: negative; Pos: positive; M: male; F: female; MIC: minimal inhibitory concentration.