

Table SI. Reported cases of basal cell carcinoma (BCC) of the nipple-areola complex

| Ref. | Age/sex | Axillary LNs (clinical examination unless otherwise stated) | Further information on histopathology | Mammography | Treatment | Follow-up |
|------|---------|---|--|--|--|---|
| 4 | 49/F | No | Superficial and infiltrating BCC | Normal | MMS | SLNB negative |
| 21 | 61/M | No | | | MMS | |
| 10 | 47/F | No | Superficial BCC | Normal | MMS | |
| 5 | 65/F | | Infiltrating BCC | Normal ("no involvement at deeper level") | MMS | No recurrence after 1 year |
| 23 | 55/M | | | | MMS | |
| 11 | 46/F | Reactive hyperplasia on SLNB axilla | Superficial | Normal | Partial mastectomy as initial resection margin not tumour free | No recurrence after 1 year |
| 19 | 57/M | No | | | Mastectomy | No recurrence after 2 years |
| 16 | 71/M | Yes | | | Simple mastectomy | |
| 20 | ?/F | No | | | Simple mastectomy | No recurrence after 1 year |
| 20 | 43/M | No | | | Simple mastectomy | No recurrence after 2 years |
| 15 | 54/M | Yes at 2 nd presentation | Metastatic BCC demonstrated in axillary LN at 2 nd presentation | | Simple excision with DXT at first presentation | Mastectomy and axillary node dissection 4 years later at 2 nd presentation |
| 6 | 75/F | No | Infiltrating BCC | Involucional changes and dermal thickening | Wedge biopsy with radiotherapy | |
| 18 | 67/F | No | | | Etretinate | |
| 7 | 67/M | No | Invasion of lactiferous ducts | Normal | Simple excision | |
| 12 | 57/M | No | Nodular | | Simple excision | No recurrence after 5 years |
| 12 | 39/M | No | Superficial | | Simple excision | No recurrence after 2 years |
| 12 | 47/F | No | Nodular | | Simple excision | No recurrence after 2 years |
| 8 | 82/F | No | Invasion of lactiferous ducts | Microcalcifications | Simple excision | No recurrence after 2 years |
| 22 | 70/M | | | Calcifications in areola | Simple excision | No recurrence after 14 months |
| 9 | 63/M | No | Invasion of lactiferous ducts | | Simple excision | No recurrence after 16 months |
| 9 | 80/M | No | Nodular with features of fibroepithelioma of Pinkus | | Simple excision | No recurrence after 8 months |
| 4 | 86/M | No | | | Simple excision | |
| 15 | 60/M | | | | Simple excision | No recurrence after 13 years |
| 13 | 51/M | No | Superficial | | Simple excision | |
| 4 | 72/M | | | | Not specified | |
| 4 | 47/M | | | | Not specified | |
| 4 | 72/M | No | | | Simple excision | |
| 4 | 66/F | No | | | Simple excision | |
| 4 | 49/F | | | | Simple excision | |
| 17 | 57/M | Yes | | | Not specified | |
| 4 | 60/M | No (confirmed histologically) | | | Simple excision | No recurrence within 18 months |

MMS: Mohs micrographic surgery; DXT: radiotherapy; LN: lymph nodes; SLNB: sentinel lymph node biopsy.