

Table SIII. Clinical characteristics, overall and by physician-judged disease severity at sampling and by physician-type, excluding patients experiencing a flare (Analysis 1)

| Characteristic | Overall | Physician-judged severity (Analysis 1) | | |
|--------------------------|-------------------|----------------------------------------|------------------|----------------|
| | | Mild | Moderate | Severe |
| Dermatologists | (n = 1,163, 100%) | (n = 969, 83.3%) | (n = 188, 16.2%) | (n = 6, 0.5%) |
| Current BSA | (n = 1021) | (n = 851) | (n = 167) | (n = 3) |
| BSA%, mean \pm SD | 4.2 \pm 6.0 | 2.9 \pm 4.3 | 10.4 \pm 8.6 | 20.7 \pm 9.0 |
| BSA >10, n (%) | 80 (7.8) | 23 (2.7) | 54 (32.3) | 3 (100.0) |
| Current PGA score, n (%) | (n = 1163) | (n = 969) | (n = 188) | (n = 6) |
| 0–1 | 655 (56.3) | 639 (65.9) | 15 (8.0) | 1 (16.7) |
| 2–5 | 508 (43.7) | 330 (34.1) | 173 (92.0) | 5 (83.3) |
| Current PASI score | (n = 713) | (n = 623) | (n = 88) | (n = 2) |
| PASI, mean \pm SD | 3.5 \pm 6.1 | 2.5 \pm 4.7 | 9.7 \pm 9.6 | 16.0 \pm 5.7 |
| PASI >10, n (%) | 39 (5.5) | 10 (1.6) | 27 (30.7) | 2 (100.0) |
| Paediatricians | n = 493, 100%) | (n = 415, 84.2%) | (n = 74, 15.0%) | (n = 4, 0.8%) |
| Current BSA | (n = 363) | (n = 300) | (n = 60) | (n = 3) |
| BSA%, mean \pm SD | 6.7 \pm 9.8 | 3.7 \pm 4.1 | 20.4 \pm 15.7 | 29.3 \pm 5.1 |
| BSA >10, n (%) | 51 (14.0) | 13 (4.3) | 35 (58.3) | 3 (100.0) |
| Current PGA score, n (%) | (n = 493) | (n = 415) | (n = 74) | (n = 4) |
| 0–1 | 247 (50.1) | 247 (59.5) | 0 (0) | 0 (0) |
| 2–5 | 246 (49.9) | 168 (40.5) | 74 (100) | 4 (100) |
| Current PASI score | (n = 274) | (n = 223) | (n = 49) | (n = 2) |
| PASI, mean \pm SD | 6.8 \pm 11.5 | 4.6 \pm 9.2 | 16.3 \pm 15.2 | 20.0 \pm 2.8 |
| PASI >10, n (%) | 50 (18.2) | 22 (9.9) | 26 (53.1) | 2 (100.0) |
| GP/PCPs | (n = 557, 100%) | (n = 456, 81.9%) | (n = 99, 17.8%) | (n = 2, 0.4%) |
| Current BSA | (n = 449) | (n = 360) | (n = 87) | (n = 2) |
| BSA%, mean \pm SD | 10.0 \pm 12.6 | 8.6 \pm 12.5 | 15.4 \pm 11.0 | 35.0 \pm 7.1 |
| BSA >10, n (%) | 117 (26.1) | 72 (20.0) | 43 (49.4) | 2 (100.0) |
| Current PGA score, n (%) | (n = 557) | (n = 456) | (n = 99) | (n = 2) |
| 0–1 | 248 (44.5) | 240 (52.6) | 8 (8.0) | 0 (0) |
| 2–5 | 309 (55.5) | 216 (47.4) | 91 (92.0) | 2 (100.0) |
| Current PASI score | (n = 156) | (n = 126) | (n = 30) | (n = 0) |
| PASI, mean \pm SD | 4.9 \pm 7.2 | 3.8 \pm 4.7 | 9.5 \pm 12.5 | – |
| PASI >10, n (%) | 15 (9.6) | 8 (6.3) | 7 (23.3) | – |

Where the full number of patients is not available, number of patients per group is indicated in the table (n).
 BSA: body surface area; GP: general practitioner; PASI: Psoriasis Area Severity Index; PCP: primary care physician; PGA: Physician's Global Assessment; SD: standard deviation.