

ORIGINAL ARTICLE

## Prevalence among adolescents in Bergen, Western Norway, of temporomandibular disorders according to the DC/TMD criteria and examination protocol

Anne Marit Graue<sup>a</sup>, Asbjørn Jokstad<sup>b</sup>, Jörg Assmus<sup>c</sup> and Marit Slättelid Skeie<sup>a</sup>

<sup>a</sup>Department of Clinical Dentistry – Pedodontics, Faculty of Medicine and Dentistry, University of Bergen, Bergen, Norway; <sup>b</sup>Department of Clinical Dentistry, Faculty of Health Sciences, UiT The Arctic University of Norway, Tromsø, Norway; <sup>c</sup>Centre for Clinical Research, Haukeland University Hospital, Bergen, Norway

### ABSTRACT

**Objective:** The aims of this study were to assess the prevalence of temporomandibular disorders (TMD) among adolescents and to contrast the prevalence of TMD according to the DC/TMD clinical examination protocol versus the prevalence of pain related to TMD according to two screening questions.

**Material and methods:** Two hundred and ten adolescents living in the county of Bergen, Norway, were offered an additional examination for TMD in connection with their regular dental check-up appointment. Five dental clinics were selected with differing socio-economic patient populations, as reflected by stratification of average levels of DMFT, and an equal number of girls and boys were invited to participate. The participants answered two screening questions for pain related to TMD followed by a clinical examination according to the DC/TMD protocol by five calibrated examiners.

**Results:** Acceptable calibration results were obtained. Approximately 80% of eligible participants consented to partake. According to the criteria of DC/TMD, the prevalence of TMD among the study participants was 11.9%, with a peak at 16 years of age. According to the self-reported screening questions for pain related to TMD, 7.2% responded positively. Only seven participants with a TMD diagnosis established according to the DC/TMD clinical examination protocol reported also pain related to TMD based on answering the two screening questions.

**Conclusion:** The prevalence of TMD is higher for girls than for boys and the prevalence of TMD established according to the DC/TMD criteria was higher than the prevalence of pain related to TMD estimated by use of two screening questions for self-reported pain.

### ARTICLE HISTORY

Received 1 March 2016  
Revised 22 April 2016  
Accepted 12 May 2016  
Published online 1 June 2016

### KEYWORDS

Adolescent; DC/TMD; prevalence; temporomandibular disorders; TMD

### Introduction

Temporomandibular disorders (TMDs) are a general term for various clinical signs and symptoms involving the masticatory muscles, the temporomandibular joint (TMJ) and associated structures.[1] TMD may influence daily life negatively by limitation of regular functions of the masticatory system, or because of pain originating in the stomatognathic muscles or the temporomandibular joints or the temple areas. The pattern of pain generally varies over time, but some individuals may suffer from longstanding problems caused by the pain related to TMD irrespective of age.[2] Adolescents with self-reported pain related to TMD report a higher consumption of analgesics and absence from school due to general pain in comparison with peers matched for age and gender.[3] For some individuals, the pain related to TMD constitutes a substantial problem and can influence all aspects of their daily life.[4] It is important to identify the earliest manifestation of TMD in young individuals, and a correct diagnosis based on signs and symptoms of TMD should be considered as an important first step in monitoring these individuals, both in relation to the condition's aetiology and for individualized preventive and therapeutic interventions.

Reported estimates of the prevalence of TMD among adolescents vary widely, ranging between 4% and 68%.[5–7] One of the likely reasons for the large variance is the diversity of diagnostic criteria and clinical examination protocols. Accordingly, the need for calibrated examiners that apply valid and reliable diagnostic methods for TMD has long been recognized and debated in the literature.[8,9] For young individuals, it is of special interest to identify the earliest manifestation of TMD. One clinical examination protocol for TMD that has gained international acceptance is The Research Diagnostic Criteria for Temporomandibular Disorders (RDC/TMD),[10] which has been widely used globally to estimate prevalences of TMD. During later years, some limitations with this protocol were identified, and a revised clinical examination protocol has recently been presented, titled The Diagnostic Criteria for Temporomandibular Disorders (DC/TMD).[11] Some of the items that were questioned in the RDC/TMD were the procedures for diagnosing myofascial pain as well as disc reduction with displacement, and the feasibility and practical application of selected palpation sites.[12] The DC/TMD clinical examination protocol appears to be valid for identifying the most common pain-related TMD diagnoses

with a reported diagnostic sensitivity of  $\geq 0.86$  and specificity  $\geq 0.98$ .<sup>[11]</sup>

Several studies in Scandinavia have been conducted with the aim to estimate the prevalence of TMD among adolescents, and possible impact on daily activities.<sup>[2–4,13]</sup> The diagnostic methods and study designs have varied. In one cross-sectional study, approximately 29 thousand adolescents living in Östergötland County, Sweden, answered two screening questions about any experiences of pain related to TMD.<sup>[13]</sup> The investigators described high reliability scores, which corroborated previous findings.<sup>[14]</sup> Furthermore, excellent diagnostic validation was reported following subsequent clinical examinations undertaken according to the RDC/TMD clinical examination protocol of 60 of the study participants versus 60 control subjects.<sup>[15]</sup> In another cross-sectional study, cohorts of 400 children and adolescents in Jönköping, Sweden, were examined clinically in 1983, 1993 and 2003 and classified according to The Clinical Dysfunction Index (Helkimo 1974).<sup>[16]</sup> To the authors' knowledge, there are no studies that report the prevalence of TMD among adolescents in Norway, or anywhere else in Scandinavia according to clinical examinations and use of diagnostic criteria of the new DC/TMD protocol.<sup>[11]</sup>

The main objective of this study was to assess the prevalence among adolescents aged from 12 up to 19 years located in Bergen, Western Norway, of TMD according to the DC/TMD clinical examination protocol. A secondary study objective was to contrast the prevalence of TMD according to the DC/TMD clinical examination protocol against the prevalence of pain related to TMD identified by use of the two screening questions for pain related to TMD adopted by Nilsson et al. <sup>[14]</sup> and by others.<sup>[15]</sup>

## Materials and methods

### Sample

The Hordaland County Council, Western Norway, offers free dental treatment to all children and adolescents up to 19 years through a public dental health service (PDHS). In the municipality of Bergen, the service includes 13 dental clinics in which adolescents attend, and these maintain records of various dental parameters in their patient populations, including average decayed, missing and filled teeth (DMFT) indices. DMFT is generally considered as a proxy for socio-economic status.<sup>[17–19]</sup> To reflect social gradients of the study sample, the 13 clinics were first ranked by average DMFT of their patient populations and alternate clinics from the top to the bottom of the ranking were invited to participate. The recruitment and clinical examination of adolescents, involved individuals within the age of 12 up to 19 years. Four of the clinics were located in urban areas, whereas one clinic was situated in a rural area.

The total population size of 12 up to 19 years old in Bergen is 21,695 individuals according to the Statistics Norway (Statistisk Sentralbyrå, SSB). The sample size calculation was based on a recent literature report.<sup>[20]</sup> We assumed a prevalence of 5% and used a precision of 3.5% for the 95% confidence interval, which gave a sample size of 149. Due to an anticipated drop-out (missing/cancelled appointments,<sup>[21]</sup>),

the goal was to include 210 individuals in the study. Hence, the five clinics aimed to recruit approximately 40 study participants each, with the objective to obtain a convenience sample of 210, assumed to reflect the social gradients of adolescents managed by the PDHS in Bergen, Western Norway.

### Study participants

All potential study participants received a written invitation to partake in the study, which detailed an additional TMD examination in connection with their regular dental check-up appointment conducted in their customary PDHS clinic. The invitation was sent in advance of the regular dental check-up appointment to those who were next in line for recall to the clinics according to the digital journal system. An equal number of boys and girls in each age group received the information letter about the additional TMD examination.

### The TMD examination

TMD was diagnosed by applying two different diagnostic tools. The study participants answered first two screening questions for pain related to TMD,<sup>[13]</sup> which was followed by a clinical examination according to the DC/TMD protocol.<sup>[11]</sup> The two screening questions for pain related to TMD were “(1) Do you have pain in your temples, face, jaw joint and jaws once a week or more?” and (2) “Do you have pain when you open your mouth wide and chew once a week or more?” If the patient answered yes to one or both the questions, they were diagnosed as having pain related to TMD. The following DC/TMD clinical examination sessions lasted for approximately 20 min.

### Calibration of clinical examiners

Prior to the initiation of the clinical examinations, the principal examiner (specialist candidate in paediatric dentistry) underwent calibration sessions together with an expert trained and educated in the use of the DC/TMD clinical examination protocol at The University of Aarhus, Denmark. Subsequent calibrations of four additional clinical examiners consisted of theory (5 h) and clinical palpation exercises. The clinical calibrations were undertaken in several sessions, and included adult volunteers (mean age 22.6 years) as well as younger volunteers (mean age 16.5 years) examined in their respective local PDHS clinics.

### Clinical examination

All the clinical examinations were conducted from November 2014 to May 2015. In four of the PDHS clinics, the local dentists performed the examinations, while the principal author examined the study participants in the fifth clinic.

The clinical examinations in compliance with the DC/TMD clinical examination protocol involved 10 different examination items: location of pain, incisal relationship, jaw opening pattern, opening movements in millimeters, lateral and protrusive movements in millimeters, TMJ noise during opening/

closing/lateral/protrusive movements, joint locking and muscle and pain from the TMJ upon calibrated palpation. Muscle palpation sites included the masseter, temporalis and supplemental muscles such as the lateral pterygoid area and the posterior and submandibular muscles. All registrations were done separately for the left and the right side of the face.[11]

The examiners used a pressure-measuring instrument regularly to facilitate proper palpation force, and to assure high inter- and intra-examiner reliabilities. Two pressure-measuring instrument instruments (Dentrade, Köln, Germany) measuring 0.5 kg and 1.0 kg were used to produce the proper force in a steady manner before every examination of muscle groups and TMJ. Additionally, the rulers for measuring millimeters were properly prepared, by cutting the end so that the edge was even with the "0" mark and by reducing the width to create space for the lip. In this study, palpation was performed for 2 s, which restricted the diagnoses for referred pain according to the DC/TMD clinical examination protocol.[11]

### The DC/TMD Symptom Questionnaire

The DC/TMD Symptom Questionnaire consists of a minimum of 5 and maximum of 14 questions, focusing on pain in the jaw, temples and ears or in front of the ears. Also items about headache, joint noise and locking of the jaw were included. The time frame that was applied according to the DC/TMD Symptom Questionnaire was consistently "the last 30 days", in line with the time frame used in the clinical examination.[11] The clinical examiner assisted the study participants if they had problems of understanding the item content.

### Diagnosis of TMD

The TMD diagnosis was established based on the DC/TMD diagnostic decision tree, using both the completed clinical examination form and the Symptom Questionnaire. The diagnostic decision tree in DC/TMD, which contains 11 different diagnosis options within three different categories, was applied.[11] This implied that one patient could have multiple diagnoses, and categories such as pain-related TMD in combination with headache, and intra-articular joint disorders in combination with degenerative joint disorders were used. All participants diagnosed with TMD were offered treatment or a referral to a specialist when needed.

### Ethical considerations

This cross-sectional study was approved by The Regional Committee for Medical Research Ethics (REK-number 2012/542). A written consent was obtained from the study participant when being 16 years and older, or from both the study participant and a parent when below 16 years.

### Statistics

Inter-examiner agreements among the five clinical examiners of the clinical measurements (mm) were assessed by applying

Bland–Altman plots. The alternatives for registration of pain upon palpation as well as for joint noise were "yes" or "no", and the inter-examiner agreements were calculated in percentages.

Possible differences between study participant subgroups were tested by using Chi-square statistics. The level of statistical significance was set at 5%. All statistical tests were computed by using a commercial statistical software package (SPSS version 22, IBM, Chicago, IL).

## Results

Out of the 210 eligible dental patients, 23 failed to show up for their appointment and a further 20 declined to partake, resulting in a study sample consisting of 167 adolescents (79.5%) aged from 12 up to 19 years (Figure 1). Fifty-one percent of the study participants were girls ( $n = 86$ ).

### Calibration and reliability of clinical examiners

The limits of agreement were within 3 mm in maximum mouth opening without assistance, and 2.5 mm for maximum mouth opening with assistance. The measurement of pain during free mouth opening showed wider limits of agreement, i.e., 4–5 mm (Figure 2). The inter-examiner agreement in registering pain upon palpation (Yes/No) in the masseter and temporalis muscle was 87.9% of all registrations. For registration of TMJ noise (Yes/No), the agreement was 86.5%.

### Self-reported pain related to TMD

Twelve study participants answered yes to at least one of the two screening questions for pain related to TMD, indicating a prevalence of 7.2%. Five study participants (5/167, 3.0%) answered "yes" to question one, while five (5/167, 3.0%) answered "yes" to question two, and two (2/167, 1.2%) answered "yes" to both questions. The prevalence for pain related to TMD was higher for girls (11.6% versus 2.5%), but did not reach a statistically significant difference (Figure 3).






Figure 1. Flow chart of the participants.



**Figure 2.** Bland–Altman plots demonstrating the deviations in clinical measurements between one principal examiner calibrated in conducting clinical examinations according to DC/TMD protocol at The University of Aarhus, Denmark (horizontal zero-line), versus the other five clinical examiners involved in the study. The clinical examinations included volunteers diagnosed with and without TMD ( $n = 18$ ). Clinical measurements focused on item 4 in the DC/TMD examination protocol, i.e., the opening movements measured in millimeters; (4a) Measurement of pain during free mouth opening, (4b) maximum mouth opening without assistance and (4c) maximum mouth opening with assistance. Horizontal red lines indicate mean difference (solid line) and limits of agreement (dash lines) between the principal examiner versus the other five clinical examiners.



**Figure 3.** Prevalence in relation to age and gender of TMD identified by use of the DC/TMD clinical examination protocol (red,  $n = 20/167$ ), and prevalence of TMD-pain identified by use of two screening questions for TMD pain (green,  $n = 12/167$ ).


**Clinical examination according to the DC/TMD protocol**

Twenty study participants were identified with a TMD diagnosis, suggesting a prevalence of 11.9%. The prevalence was higher for the girls compared with the boys (19.8%,  $n = 17/86$  vs. 3.7%,  $n = 3/81$ ,  $p < 0.002$ ). The age group showing the highest prevalence of TMD (28.0%,  $n = 7/25$ ) was at 16 years (Figure 3).

The most common TMD diagnosis was disc displacement with reduction, with a prevalence of 5.4% (9/167), followed by myalgia, 3.0% (5/167) (Figure 4). Three participants (1.8%)

had more than one TMD diagnosis. The combination of myalgia and arthralgia occurred in two participants (1.2%) and a combination of arthralgia and headache attributed to TMD in one participant (0.6%). Disc displacement with intermittent locking occurred in one participant (0.6%).

Out of the 20 participants diagnosed with TMD, 12 (60.0%) reported pain from the jaw, temple or ear alternatively in front of the ear during the last 30 d. Of those, 10 participants (eight girls) reported worsening of the intensity of pain during activities like chewing, opening the mouth, chewing and



**Figure 4.** Frequency distribution of TMD diagnoses according to the DC/TMD clinical examination protocol among 12 up to 19 years old adolescents in Bergen, Western Norway ( $n = 20/167$ ).

talking. Nine out of the 20 participants (45.0%) with a DC/TMD diagnosis reported temple headache in the DC/TMD symptom questionnaire. Among the participants with no DC/TMD diagnosis, 29 out of 147 participants (19.7%) reported headache within the last 30 d in the DC/TMD questionnaire. None of the study participants complained of any adverse effects after the clinical examination according to the DC/TMD protocol.

#### **Self-reported pain related to TMD versus clinical examination**

Five study participants whom responded positively to the two screening questions for pain related to TMD did not qualify for a diagnosis according to DC/TMD clinical examination protocol. Four of these had responded in the DC/TMD symptom questionnaire that they suffered from headache.

#### **Discussion**

Practically all children make use of the public dental health services in Norway, regardless of socioeconomic background. The present study sample should, therefore, represent different socio-economic patient backgrounds, particularly by sampling the clinics with differences in the clinics' DMFT-indices. The recruitments for study participation followed also the recall lists for regular dental appointments. The response rate of approximately 80% was deemed satisfactory for being a cross-sectional clinical study.[22] In sum, the TMD data collected in the current study are considered as likely representative for adolescents living in the Bergen municipality.

Calibration over multiple sessions has been shown to improve the reliability of clinical examinations for TMD,[23] and this was endeavoured by the adoption of various calibration exercises. Acceptable calibration results were obtained for vertical range of motion of the mandible in the maximum- and maximum-assisted mouth opening, while wider limits of agreement were seen when asking for unassisted opening without pain. The observation corroborate previous findings,[23] and is likely a consequence of difficulties for the patient to know exactly at what level of mouth opening the pain onset will occur.

The higher prevalence for girls than for boys (19.8% versus 3.7%) was expected and in accordance with other studies, as female gender is identified as a risk factor for TMD.[13,24,25] That girls constituted the vast majority of those who reported

a worsening of pain during activities (chewing, talking and opening the mouth) are also in accordance with previous findings.[3] It has been suggested that self-reported TMD symptoms in girls are likely related to pubertal development.[26] Not all the study participants diagnosed with TMD according to DC/TMD experienced pain due to their condition. Especially for those with disc displacement with reduction, pain was not a common problem. It is important to be aware of the fact since pain often is the main cause when seeking help with TMD problems.[3]

The peak of TMD among the 16-year-old study participants was interesting in the context that the majority of Norwegian adolescents at that age change from compulsory secondary school to high school. Greater expectations from both the students themselves and their environment on their achievements with the schoolwork can be stressful and perhaps might be a factor in onset of TMD. A Swedish study from 2008 on mental health among adolescents from 16 to 18 years old found that a large proportion associated self-perceived stress with high pressure and demands from school, and also that perceived stress correlated strongly with health complaints like tiredness, headache, musculoskeletal pain and sleeping difficulties. Especially the girls reported significantly higher levels of stress.[27] A comparable study has recently been completed in Norway in 16 and 17-years old and corroborates the Swedish study in that high levels of perceived stress and musculoskeletal pain seem to prevail.[28] The highest levels of stress and pain values were reported in the group with head pain. Bodily pain elsewhere than in the orofacial region in adolescents is associated with pain related to TMD,[24] and the probability of facial pain increases with the number of other pain conditions.[25]

The prevalence of pain related to TMD in adolescents in Bergen in 2015 according to the two screening questions for pain related to TMD was 7.2%, and thus higher than in a group of adolescents in Sweden in 2000.[13] The cause of the higher value in the present study is unclear. Given that both cohorts are representative and the examiners calibrated, the observation may indicate an increase in the prevalence of pain related to TMD among adolescents today.

That the two screening questions for pain related to TMD differed from the DC/TMD examination with regard to identifying individuals likely to have TMD are probably because these may prompt the reporting of various forms of headaches that are likely unrelated to TMD. Moreover, since there is no time limit for the two screening questions, there is no way of distinguishing between chronic and acute pains. In

contrast, the DC/TMD clinical examination protocol is limited to the examination of structures within the stomatognathic domain, and only to experienced symptoms registered within the last 30 d.[11] Moreover, the spectrum of TMD diagnoses includes also asymptomatic conditions such as disc displacements, which the two pain screening questions will likely fail to identify.

Only seven participants were diagnosed with TMD according to the DC/TMD clinical examination protocol, and with pain related to TMD according to the two screening questions. Knowing that the most frequent diagnosis according to the DC/TMD clinical examination protocol was disc displacement with reduction and that this condition often is pain free, this may be a reason why several participants did not get a TMD diagnosis according to two screening questions for pain related to TMD.

Four out of the five participants that reported pain according to the two screening questions for pain related to TMD were not diagnosed with TMD according to the DC/TMD criteria and examination protocol. However, they had marked for headache in the temple areas in the DC/TMD symptom questionnaire. The positive response to the two screening questions for pain related to TMD might be actually due to tension-type headaches.[14] Common for these study participants was that they reported that the headache had been there for a long period, varying for 1–6 years. Tension-type headaches are relatively common among adolescents [29] and in the current cross-sectional study, self-reported headache was more than twice as frequent in the TMD group established according to the DC/TMD clinical examination protocol in comparison with those without a TMD diagnosis. Our findings thus corroborate the observations that TMD and tension-type headache often are coexisting conditions [30] and that headache seems to precede pain related to TMD in adolescents.[31]

### Limitations of the study

A larger sample size would have strengthened the reliability of the current study results. However, conducting clinical examinations according to the DC/TMD protocol is time consuming, and it was not possible to achieve a larger sample size within the existing logistical and financial limits.

### Conclusion

In this study, it was found that a substantial part of adolescents had a TMD diagnosis according to the DC/TMD protocol. Girls were found to have a significantly higher prevalence of TMD diagnoses according to the DC/TMD protocol than boys, and girls also reported more pain related to TMD than boys

### Acknowledgements

The authors would like to thank the Public Dental Health Services in Hordaland, Norway, for contributing with time and personnel resources, and especially the dentists who contributed as clinical examiners. Our

gratitude also goes to all the participants accepting to be included in the study.

### Disclosure statement

The authors report that they have no conflicts of interests. The authors alone are responsible for the content and writing of the paper.

### Funding information

This research was carried out with funding from The Oral Health Centre of Expertise in Western Norway, Hordaland.

### Notes on contributors

**Anne Marit Graue** contributed to the study design, calibration and conducted the clinical examination. She was also the main contributor of the manuscript.

**Asbjørn Jokstad** contributed with his expert advice and critical review of the manuscript.

**Jörg Assmus** carried out statistical analyses and the figures in the manuscript.

**Marit Slåttestad Skeie** contributed to the original study design of the project, the calibration of examiners and in the writing of the manuscript.

### References

- [1] Okeson J. Management of temporomandibular disorders and occlusion, 7 ed. St. Louis (MI): Elsevier; 2013.
- [2] Nilsson IM, List T, Drangsholt M. Incidence and temporal patterns of temporomandibular disorder pain among Swedish adolescents. *J Orofac Pain.* 2007;21:127–132.
- [3] Nilsson IM, Drangsholt M, List T. Impact of temporomandibular disorder pain in adolescents: differences by age and gender. *J Orofac Pain.* 2009;23:115–122.
- [4] Nilsson IM, List T, Willman A. Adolescents with temporomandibular disorder pain – the living with TMD pain phenomenon. *J Orofac Pain.* 2011;25:107–116.
- [5] Nydell A, Helkimo M, Koch G. Craniomandibular disorders in children – a critical review of the literature. *Swed Dent J.* 1994;18:191–205.
- [6] Thilander B, Rubio G, Pena L, et al. Prevalence of temporomandibular dysfunction and its association with malocclusion in children and adolescents: an epidemiologic study related to specified stages of dental development. *Angle Orthod.* 2002;72:146–154.
- [7] Sena MF, Mesquita KS, Santos FR, et al. Prevalence of temporomandibular dysfunction in children and adolescents. *Rev Paul Pediatr.* 2013;31:538–545.
- [8] Mohl ND. Reliability and validity of diagnostic modalities for temporomandibular disorders. *Adv Dent Res.* 1993;7:113–119.
- [9] Dahlstrom L, Keeling SD, Friction JR, et al. Evaluation of a training program intended to calibrate examiners of temporomandibular disorders. *Acta Odontol Scand.* 1994;52:250–254.
- [10] Dworkin SF, LeResche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. *J Craniomandib Disord.* 1992;6:301–355.
- [11] Schiffman E, Ohrbach R, Truelove E, et al. Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) for Clinical and Research Applications: recommendations of the International RDC/TMD Consortium Network\* and Orofacial Pain Special Interest Groupdagger. *J Oral Facial Pain Headache.* 2014;28:6–27.
- [12] List T, Greene CS. Moving forward with the RDC/TMD. *J Oral Rehabil.* 2010;37:731–733.
- [13] Nilsson IM, List T, Drangsholt M. Prevalence of temporomandibular pain and subsequent dental treatment in Swedish adolescents. *J Orofac Pain.* 2005;19:144–150.
- [14] Nilsson IM, List T, Drangsholt M. The reliability and validity of self-reported temporomandibular disorder pain in adolescents. *J Orofac Pain.* 2006;20:138–144.

- [15] Wahlund K, List T, Dworkin SF. Temporomandibular disorders in children and adolescents: reliability of a questionnaire, clinical examination, and diagnosis. *J Orofac Pain* 1998;12:42–51.
- [16] Kohler AA, Helkimo AN, Magnusson T, et al. Prevalence of symptoms and signs indicative of temporomandibular disorders in children and adolescents. A cross-sectional epidemiological investigation covering two decades. *Eur Arch Paediatric Dentistry: Off J Eur Acad Paediatric Dentistry* 2009;10:16–25.
- [17] Christensen LB, Twetman S, Sundby A. Oral health in children and adolescents with different socio-cultural and socio-economic backgrounds. *Acta Odontol Scand*. 2010;68:34–42.
- [18] Petersen PE. Sociobehavioural risk factors in dental caries – international perspectives. *Commun Dent Oral Epidemiol*. 2005;33:274–279.
- [19] Socialstyrelsen. Sociala skillnader i tandhälsa bland barn och unge. 2013; Artikelnr 2013-5-34; [cited 2016 May 27]. Available from: <http://www.socialstyrelsen.se/publikationer2013/2013-5-34>.
- [20] List T, Ekberg E, Ernberg M, et al. Ny diagnostik för de vanligaste tempromadibulära dysfunktionerna för användning i allmän-tandvården- DC/TMD. *Nor Tannlegeforen Tid*. 2015;125:142–150.
- [21] Skaret E, Raadal M, Berg E, et al. Dental anxiety and dental avoidance among 12 to 18 year olds in Norway. *Eur J Oral Sci*. 1999;107:422–428.
- [22] Salant P. How to conduct your survey. New York: John Wiley & Sons; 1994.
- [23] List T, John MT, Dworkin SF, et al. Recalibration improves inter-examiner reliability of TMD examination. *Acta Odontol Scand*. 2006;64:146–152.
- [24] Fernandes G, van Selms MK, Goncalves DA, et al. Factors associated with temporomandibular disorders pain in adolescents. *J Oral Rehabil*. 2015;42:113–119.
- [25] LeResche L, Mancl LA, Drangsholt MT, et al. Predictors of onset of facial pain and temporomandibular disorders in early adolescence. *Pain* 2007;129:269–278.
- [26] Hirsch C, Hoffmann J, Turp JC. Are temporomandibular disorder symptoms and diagnoses associated with pubertal development in adolescents? An epidemiological study. *J Orofac Orthop*. 2012;73:6–8, 10–18.
- [27] Wiklund M, Malmgren-Olsson EB, Ohman A, et al. Subjective health complaints in older adolescents are related to perceived stress, anxiety and gender – a cross-sectional school study in Northern Sweden. *BMC Public Health* 2012;12:993.
- [28] Osteras B, Sigmundsson H, Haga M. Perceived stress and musculoskeletal pain are prevalent and significantly associated in adolescents: an epidemiological cross-sectional study. *BMC Public Health* 2015;15:1081.
- [29] Xavier MK, Pitangui AC, Silva GR, et al. Prevalence of headache in adolescents and association with use of computer and videogames. *Ciencia Saude Coletiva* 2015;20:3477–3486.
- [30] Speciali JG, Dach F. Temporomandibular dysfunction and headache disorder. *Headache* 2015;55:72–83.
- [31] Nilsson IM, List T, Drangsholt M. Headache and co-morbid pains associated with TMD pain in adolescents. *J Dent Res*. 2013;92:802–807.