

ORIGINAL ARTICLE

Self-reported temporomandibular disorder symptoms and severity of malocclusion in prospective orthognathic-surgical patients

Anna-Liisa Svedström-Oristo^{a,b}, Heidi Ekholm^c, Mimmi Tolvanen^d and Timo Peltomäki^{e,f}

^aDepartment of Oral Development and Orthodontics, Institute of Dentistry, University of Turku, Turku, Finland; ^bDepartment of Oral and Maxillofacial Diseases, Turku University Hospital, Turku, Finland; ^cInstitute of Dentistry, University of Turku, Turku, Finland; ^dDepartment of Community Dentistry, Institute of Dentistry, University of Turku, Turku, Finland; ^eSchool of Medicine, University of Tampere, Tampere, Finland; ^fOral and Maxillofacial Unit of Tampere University Hospital, Tampere, Finland

ABSTRACT

Objective: The objective of this study is to analyze the association between self-reported symptoms of temporomandibular joint disorder (TMD) and the severity of malocclusion in prospective orthognathic-surgical patients.

Material and methods: The subjects consisted of 50 consecutive patients (13 males and 37 females) referred to two university clinics for assessment of orthodontic-surgical treatment need. Data considering self-reported TMD symptoms were gathered using a semi-structured diary. At the first appointment, all patients rated the importance of treatment (on a scale of 1–10) and assessed self-perceived dental appearance using a VAS scale. The scale was anchored with photographs 1 and 10 from the Aesthetic Component (AC) of the Index of Orthodontic Treatment Need (IOTN). Study models were assessed by an experienced orthodontic specialist using the Peer Assessment Rating (PAR) index and the Index of Complexity, Outcome and Need (ICON). Association between the PAR and ICON scores and the number of reported symptoms was analyzed statistically.

Results: Seventy-one percent of patients reported experiencing TMD symptoms. The most prevalent symptoms were pain in the head and/or neck region and fatigue in the TMJ region. The number of symptoms was highest in the morning. Ninety percent of males and 86% of females rated the importance of treatment as high; males experiencing TMD symptoms tended to rate surgery as more important compared with males with no symptoms ($p = 0.056$).

Conclusions: In this sample, the results cannot unambiguously confirm an association between self-reported symptoms of TMD and objectively defined severity of malocclusion.

ARTICLE HISTORY

Received 29 September 2015
Revised 30 May 2016
Accepted 3 June 2016
Published online 23 June 2016

KEYWORDS

Index of Complexity, Outcome and Need; peer assessment rating; self-perception

Introduction

Temporomandibular disorder (TMD) incorporates numerous signs and symptoms involving masticatory musculature, temporomandibular joints (TMJs), and associated structures. TMD has been found to be a major cause of non-dental pain in the orofacial region.[1] The most frequent symptoms are pain in masticatory muscles and in TMJs, TMJ sounds, and limitations in mandibular movements.[2,3] Several predisposing, triggering, and sustaining factors for TMD have been identified: female gender,[4–7] mandibular instability,[4] depression,[8] and parafunctional habits.[4,6] In a recent study by Sierwald et al. [9], TMD patients reported two to three times higher percentages of both daytime and nighttime bruxism than controls. Although these two types of bruxism reflect various backgrounds,[10–12] both types are regarded as contributing to the development of TMD.[9]

TMD has previously been linked with strong evidence to malocclusion, e.g., open bite, deep bite, and cross bite.[2,4,6,13] However, there are also studies that have not considered malocclusion a risk factor [5,14] or that have suggested that it should only be considered a contributing factor in the multifactorial aetiology of TMD.[15] In Finland, three

out of four patients seeking orthodontic-surgical treatment suffer from TMD symptoms.[16,17] After combined orthodontic-surgical treatment, a significant reduction has been found especially in TMD-related headaches. Even other symptoms, such as TMJ crepitation and muscle soreness, have been found to decrease.[17–20] The results are, however, somewhat ambiguous: in 4–41% of patients, the subjective symptoms have been found to increase post-operatively,[2,16,18,21] or there has been no significant change between pre- and postoperative symptoms.[22,23] It has been suggested that patients suffering from preoperative TMD and undergoing orthognathic surgery have an increased risk for postoperative worsening of the symptoms.[21]

The aim of this study was to analyze the association between self-reported symptoms of TMD and the severity of malocclusion in prospective orthognathic-surgical patients. The hypothesis suggested a correlation between self-reported TMD symptoms and the severity of malocclusion.

Material and methods

The subjects comprised 50 consecutive patients (13 males and 37 females, mean age 33.6 years, age range 18–61 years)

referred to two university clinics for assessment of orthognathic-surgical treatment need.

Data considering TMD symptoms were gathered using a semi-structured diary that included seven statements with two response alternatives, yes and no: (1) I have pain in my head and/or neck region, (2) I have pain in the area of my jaw joint, (3) I hear sounds like clicking or crepitation from my jaw joint, (4) My jaws feel tired or stiff, (5) I have difficulties in opening my mouth, (6) I suffer from teeth clenching or grinding, (7) I have difficulties in chewing, and (8) Some other problem, please explain. Statements 1–7 were selected from the Research Diagnostic Criteria for Temporomandibular Disorder (RDC-TMD) History Questionnaire [24] and the Diagnostic Criteria for Temporomandibular Disorder (DC/TMD) Symptom Questionnaire.[25] They were slightly modified to focus on the patient's current experience. The respondents were instructed to fill in the diaries during one working day and one day off work as soon as possible following the appointment. The diaries were to be filled in four times a day: (a) after waking up or at 8 am, (b) after lunch or at noon, (c) after work or at 4 pm, and (d) when going to sleep or at 8 pm.[26] Prior to the study, the diary was tested among eight students for possible revision. However, no amendments were needed.

In the clinical examination, patients were asked to rate the importance of treatment on a scale from 1 (not important) to 10 (extremely important). Further, they were asked to assess self-perceived dental appearance on a visual analogue scale (VAS) anchored with two colour photographs: number 1 (no aesthetically based treatment need) and number 10 (definite treatment need) from the Aesthetic Component (AC) of the Index of Orthodontic Treatment Need (IOTN).[27]

Impressions for study models were taken and the study models were assessed by an experienced, calibrated orthodontic specialist using the Peer Assessment Rating (PAR) index [28] and the Index of Complexity, Outcome and Need (ICON).[29] The PAR scores five occlusal components: (1) relationship between dental midlines in upper and lower jaw, (2) overjet, (3) overbite, (4) alignment in upper and lower anterior segments, and (5) sagittal, vertical, and transversal relationships on the left and right sides. Individual scores are weighted with coefficients from 1 to 6, depending on the component. The weighted scores are added up to give a total PAR score. Similarly, the ICON includes five assessments: (1) overbite, (2) upper arch crowding and spacing, (3) quality of buccal segment interdigitation (both sides), (4) transversal relationship between upper and lower dental arch, and (5) assessment of dental appearance using the AC. In evaluation of study models, dental appearance is assessed using the black and white AC scale. In the ICON, the weightings range from 3 to 7 and the weighted sub-scores comprise the final ICON score. Seventeen of the study models (34%) were rescored using both PAR and ICON. The intra-examiner reliability was good (the intraclass correlation coefficients were 0.86 and 0.89, respectively).

The study protocol was approved by the Ethics Review Committees of the Hospital District of South-West Finland and the Joint Municipal Authority of the Pirkanmaa Hospital

District. All patients signed an informed consent form before the study.

Statistical analyses

To describe prevalence, TMD symptoms were analyzed according to the replies, i.e., symptoms/no symptoms. Moreover, a new sum score variable describing the number of symptoms was calculated by adding up the answers for all the different symptoms. Because all patients did not indicate which of the diaries were filled in during a working day or a day off, the responses in the two diaries were combined and the mean values were used in further analyses. The importance of treatment was analyzed using both continuous and dichotomized values (scale 1–10, values 1–6 “not important” and values 7–10 “important”). AC and VAS values of 1–4 were interpreted as “no treatment need”, values of 5–7 as “borderline” and values of 8–10 as “definite treatment need.” In addition, the ratings were dichotomized into two categories, “no aesthetic treatment need” (values 1–4) and “aesthetic treatment need” (values 5–10).

The differences in the number of symptoms, importance of treatment, self-perceived dental aesthetics and PAR, ICON, and AC scores between females and males were evaluated using the Mann–Whitney *U* test. Differences in the prevalence of symptoms, dichotomized importance of treatment, and dental aesthetics were evaluated with the chi square test. Changes in the frequency of symptoms during the day were analyzed using general linear modelling for repeated measures using Huynh–Feldt correction ($\epsilon > 0.75$). Associations among the PAR, ICON, and AC scores and among age, AC scores, and the number of TMD symptoms were evaluated with Spearman correlation coefficients. *p* Values of < 0.05 were interpreted as statistically significant. The analyses were conducted using the SPSS Statistical Package (IBM SPSS Statistics, V22.0, Armonk, NY).

Results

A total of 71% of patients (76% of females, 55% of males, $p = 0.163$) reported TMD symptoms. The total number of symptoms was highest in the morning and decreased during the day ($p = 0.001$). The most prevalent symptom was pain in the head and/or neck region, followed by fatigue or stiffness in the TMJ region, difficulties in chewing, and joint sounds. The prevalence details of the different symptoms are given in Table 1. The mean PAR score was 26.46 and the mean ICON score was 62.42. There were no statistically significant associations between PAR or ICON scores and the number of self-reported TMD symptoms (Table 2).

The majority of all respondents (86% of females and 90% of males) rated the importance of treatment as 7 or higher. Males reporting TMD symptoms tended to rate treatment as more important than males with no symptoms ($p = 0.056$). Subjective assessment of dental appearance was 5 or more in 66% of females and 64% of males ($p = 0.905$), while 59% of females and 69% of males were assessed as having AC scores of 5 or more by the orthodontic professional ($p = 0.532$). The

Table 1. The prevalence of different symptoms in 34 females and 11 males who replied to symptom statements. A total of 26 females and six males reported symptoms.

		Morning	Noon	Afternoon	Evening
Pain in the head and/or neck region	Day 1	20	12	12	10
	Day 2	13	9	8	10
Fatigue and/or stiffness in TMJ region	Day 1	19	9	11	10
	Day 2	12	8	5	11
Difficulty chewing	Day 1	13	8	10	8
	Day 2	6	5	6	6
TMJ sounds	Day 1	13	6	5	2
	Day 2	5	4	5	2
Pain in the area of TMJ	Day 1	12	8	9	8
	Day 2	6	4	4	6
Teeth clenching or grinding	Day 1	9	2	3	3
	Day 2	3	0	1	2
Difficulty opening mouth	Day 1	7	5	3	3
	Day 2	4	2	2	3
Some other reason	Day 1	3	2	1	2
	Day 2	1	4	1	1
Number of those reporting symptoms	Day 1	27	14	14	12
	Day 2	18	13	13	16

The highest prevalences given in bold. TMJ: temporomandibular joint.

Table 2. Descriptive statistics for the distribution of TMD-symptoms, self-rated importance of surgery, and self-perceived dental aesthetics among all patients (37 females and 13 males), and separately among those patients who reported symptoms (26 females and six males).

	All patients			Patients with symptoms		
	Mean	Median	Range	Mean	Median	Range
Mean number of symptoms in the morning ^a						
Females	1.8	1.5	0–7	2.3	2	0–7
Males	1.0	0.25	0–3.5	1.7	1.5	0–3.5
Mean number of symptoms at noon ^a						
Females	1.1	0	0–7	1.5	0.75	0–7
Males	0.7	0	0–3	1.3	1.25	0–3
Mean number of symptoms in the afternoon ^a						
Females	1.2	0	0–7	1.5	1	0–7
Males	0.6	0	0–3	1.2	0.75	0–3
Mean number of symptoms in the evening ^a						
Females	1.1	0	0–4.5	1.3	1	0–4.5
Males	0.8	0.25	0–3	1.5	1.25	0–3
Mean number of all symptoms ^a						
Females	1.3	0.5	0–5.8	1.7	1	0.3–5.8
Males	0.8	0.25	0–3	1.4	1.1	0.1–3.0
Importance of treatment (scale 1–10)						
Females	8.4	9	5–10	8.8	9	6–10
Males	8.1	8	6–10	9.3	9.5	8–10
Subjective dental aesthetics (scale 1–10)						
Females	5.3	5.5	1–9	5.3	6	1–9
Males	5.0	5	3–8	5.2	5	3–8
PAR score						
Females	26.4	29	6–42	27.5	29.5	14–42
Males	26.5	28	8–43	23.2	21.5	8–39
ICON score						
Females	62.7	63	13–94	62.3	64.5	25–94
Males	61.6	60	35–92	52.2	53	35–71
AC of the IOTN						
Females	5.9	6	1–10	6.0	6.5	2–10
Males	5.7	5	3–9	4.5	4.5	3–7

No statistically significant differences were observed between males and females.

^aTo questions about symptoms, only 34 females and 11 males replied.

professionally assessed AC scores were lower (indicating a more pleasing dental appearance and lower treatment need) among males with TMD symptoms than among males without symptoms (mean 4.5, range 3–7 versus mean 6.6, range 4–9) ($p = 0.078$). Further, ICON scores were lower among males with TMD symptoms than among males without

symptoms (mean 52.2, range 35–71 versus mean 70.4, range 53–92) ($p = 0.100$).

Respondents' age was not correlated with the number of TMD symptoms or with the professionally defined AC (scale 1–10). Correlation between the PAR and ICON scores was 0.763 ($p < 0.001$), and 0.782 ($p < 0.001$) between the PAR and AC scores. By gender, statistically significant differences were found neither in the PAR, ICON, or AC scores nor in the number of symptoms, importance of treatment or self-perceived dental aesthetics (Table 2).

Discussion

In line with previous studies, the most frequently reported symptoms of TMD included pain, joint sounds, and functional limitations.[2,3,7] The finding that these symptoms and the severity of objectively assessed malocclusions had no obvious association is in accordance with the results by Sipilä et al.[30] who also found no association between facial pain and occlusal traits assessed using the PAR. However, Mohlin et al.[31] analyzed the association between TMD and occlusal features and reported that young adults with TMD had significantly higher PAR scores than controls. This difference is presumably due to the fact that in the two former studies, comparisons were made using subjective symptoms of TMD, while Mohlin et al. [31] compared clinically determined, objective signs and occlusal characteristics.

A total of 71% of subjects reported TMD symptoms, which is in line with earlier reports.[16,17] A higher number of symptoms were reported in the morning than at other times of day. Also Nishiyama et al. [32] found that the number of symptoms decreased during the day. They concluded that habitual behavioural factors, such as sleep bruxism, exert a strong effect on TMD-related symptoms. In the current sample, however, only a few patients reported teeth clenching/grinding, emphasizing the need to further analyze temporal differences in the prevalence of various symptoms.

The majority of the patients rated orthognathic-surgical treatment as highly important. Given that the condition-specific quality of life has been found to be lower in preoperative orthodontic-surgical patients with TMD than in controls,[33] it is not surprising that males reporting TMD symptoms rated the treatment as more important than did their symptom-free counterparts. In females, the difference was negligible. The difference between male and female patients may be explained by the finding that in males, occlusal traits have been shown to be directly associated with health-related quality of life, while in females, the association is indirect.[34]

The present study had some limitations. First, the sample size was small, especially in terms of the number of males. However, the numbers were comparable with another recent study investigating the effects of orthodontic and orthodontic-surgical treatments on severe malocclusion.[35] It is possible that the low participation rate of males reflects previous findings that females experience TMD symptoms more frequently than males.[4,7,36] Therefore, the threshold for seeking treatment may be lower among women.[34] Despite the small number, the difference in self-rated

importance of orthognathic-surgical treatment almost reached statistical importance when comparing symptomatic and asymptomatic males.

The second limitation was related to the diary. It included selected questions from the RDC/TMD History Questionnaire [24] and the DC/TMD Symptom Questionnaire.[25] The RDC/TMD diagnostic system has been a widely used method, and its application has been recommended.[37] Among the history-taking questions, attention has been paid especially to self-reported pain, limitations in mouth opening, and joint sounds.[7,37] In the present study, we used slightly modified symptom-related statements focusing exactly on a patient's most recent experiences. Although the reliability and the validity of the original questionnaires have been tested, it is possible that application of the current subset has affected these values. Therefore, the results should be interpreted with some reservation.

Although PAR and ICON have been proposed as suitable in evaluating occlusion in orthognathic-surgical patients,[38] it should be kept in mind that their focus is in static occlusion. Assessment of functional factors is not incorporated, even though they are a primary reason for seeking orthognathic-surgical treatment.[39] PAR emphasizes especially overjet and deviation of dental midlines [28] while ICON's highest weightings focus on dental appearance and crowding.[29] It is possible that some of the patients referred for assessment of orthognathic-surgical treatment need already have aligned dental arches. Therefore, they may not get high scores according to the indices (Table 2). To obtain a more comprehensive view on the dimensions of severe malocclusion, the index scores should be complemented by patients' own assessment of pain and other everyday consequences.

To summarize, the small sample size and the fact that not all respondents reported symptoms prevent definitive conclusions. The results cannot unambiguously confirm an association between self-reported symptoms of TMD and objectively defined severity of malocclusion.

Disclosure statement

The authors report that they have no conflicts of interest.

Funding information

This study was supported by grants from the Finnish Dental Society Apollonia, Orthodontic Division of the Finnish Dental Society, Finnish Women Dentists' Association, University Hospital of Turku, and Competitive Research Funding from the Pirkanmaa Hospital District.

Notes on contributors

A.-L. Svedström-Oristo is Associate Professor at Institute of Dentistry, University of Turku.

H. Ekholm is Dental Student at Institute of Dentistry, University of Turku.

M. Tolvanen is Statistician at Institute of Dentistry, University of Turku.

T. Peltomäki is Professor at School of Medicine, University of Tampere.

References

- [1] Irving J, Wood GD, Hackett AF. Does temporomandibular disorder pain dysfunction syndrome affect dietary intake? *Dent Update*. 1999;26:405–407.
- [2] Dujoncqoy JP, Ferri J, Raoul G, et al. Temporomandibular joint dysfunction and orthognathic surgery: a retrospective study. *Head Face Med*. 2010;6:27.
- [3] Molina OF, dos Santos J Jr, Nelson SJ, et al. Profile of TMD and bruxer compared to TMD and nonbruxer patients regarding chief complaint, previous consultations, modes of therapy, and chronicity. *Cranio*. 2000;18:205–219.
- [4] Marklund S, Wänman A. Risk factors associated with incidence and persistence of signs and symptoms of temporomandibular disorders. *Acta Odontol Scand*. 2010;68:289–299.
- [5] Bonjardim LR, Lopes-Filho RJ, Amado G, et al. Association between symptoms of temporomandibular disorders and gender, morphologic occlusion, and psychological factors in a group of university students. *J Dent Res*. 2009;20:190–194.
- [6] Magnusson T, Egermark I, Carlsson GE. A prospective investigation over two decades on signs and symptoms of temporomandibular disorders and associated variables. A final summary. *Acta Odontol Scand*. 2005;63:99–109.
- [7] Bagis B, Ayaz EA, Turgut S, et al. Gender difference in prevalence of signs and symptoms of temporomandibular joint disorders: a retrospective study on 243 consecutive patients. *Int J Med Sci*. 2012;9:539–544.
- [8] Yap AU, Tan KB, Chua EK, et al. Depression and somatization in patients with temporomandibular disorders. *J Prosthet Dent*. 2002;88:479–484.
- [9] Sierwald I, John MT, Schierz O, et al. Association of temporomandibular disorder pain with awake and sleep bruxism in adults. *J Orofac Orthop*. 2015;76:1–13.
- [10] Feu D, Catharino F, Quintão CC, et al. A systematic review of etiological and risk factors associated with bruxism. *J Orthod*. 2013;40:163–171.
- [11] Lobbezoo F, Ahlberg J, Glaros AG, et al. Bruxism defined and graded: an international consensus. *J Oral Rehabil*. 2013;40:2–4.
- [12] Manfredini D, Lobbezoo F. Role of psychosocial factors in the etiology of bruxism. *J Orofac Pain*. 2009;23:153–166.
- [13] Abrahamsson C, Henrikson T, Nilner M, et al. TMD before and after correction of dentofacial deformities by orthodontic and orthognathic treatment. *Int J Oral Maxillofac Surg*. 2013;42:752–758.
- [14] Manfredini D, Guarda-Nardini L, Marchese-Ragona R, et al. Theories on possible temporal relationship between sleep bruxism and obstructive sleep apnea events. An expert opinion. *Sleep Breath*. 2015;19:1459–1465.
- [15] Gesch D, Bernhardt O, Kocher T, et al. Association of malocclusion and functional occlusion with signs of temporomandibular disorders in adults: results of the population-based study of health in Pomerania. *Angle Orthod*. 2004;74:512–520.
- [16] Pahkala R, Kellokoski JK. Surgical-orthodontic treatment and patients' functional and psychosocial well-being. *Am J Orthod Dentofacial Orthop*. 2007;132:158–164.
- [17] Panula K, Somppi M, Finne K, et al. Effects of orthognathic surgery on temporomandibular joint dysfunction. A controlled prospective 4-year follow-up study. *Int J Oral Maxillofac Surg*. 2000;29:183–187.
- [18] Al-Riyami S, Cunningham SJ, Moles DR. Orthognathic treatment and temporomandibular disorders: a systematic review. Part 2. Signs and symptoms and meta-analyses. *Am J Orthod Dentofacial Orthop*. 2009;136:626.e1–626.e16.
- [19] Al-Riyami S, Moles DR, Cunningham SJ. Orthognathic treatment and temporomandibular disorders; a systematic review. Part 1. A new quality-assessment technique and analysis of study characteristics and classifications. *Am J Orthod Dentofacial Orthop*. 2009;136:624.e1–624.e15.
- [20] Silvola AS, Rusanen J, Tolvanen M, et al. Occlusal characteristics and quality of life before and after treatment of severe malocclusion. *Eur J Orthod*. 2012;34:704–709.
- [21] Wolford L, Reiche-Fischel O, Mehra P. Changes in temporomandibular joint dysfunction after orthognathic surgery. *J Oral Maxillofac Surg*. 2003;61:655–660.
- [22] Farella M, Michelotti A, Bocchino T, et al. Effects of orthognathic surgery for class III malocclusion on signs and symptoms of

- temporomandibular disorders and on pressure pain thresholds of the jaw muscles. *Int J Oral Maxillofac Surg.* 2007;583–587.
- [23] Onizawa K, Schmelzeisen R, Vogt S. Alteration of temporomandibular joint symptoms after orthognathic surgery: comparison with healthy volunteers. *J Oral Maxillofac Surg.* 1995;53:117–121.
- [24] Dworkin SF, Le Resche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. *J Craniomandib Disord.* 1992;6:301–355.
- [25] Schiffman E, Ohrbach R, Truelove E, et al. Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) for Clinical and Research Applications: Recommendations of the International RDC/TMD Consortium Network and Orofacial Pain Special Interest Group. Individual Instruments and Forms. *J Oral Facial Pain Headache.* 2014;28:6–27.
- [26] Alanko OM, Svedström-Oristo AL, Peltomäki T, et al. Psychosocial well-being of prospective orthognathic-surgical patients. *Acta Odontol Scand.* 2014;72:887–897.
- [27] Brook PH, Shaw WC. The development of an index of orthodontic treatment priority. *Eur J Orthod.* 1989;11:309–320.
- [28] Richmond S, O'Brien K, Buchanan I, et al. An introduction to occlusal indices. Manchester: Mandent Press; 1992.
- [29] Daniels C, Richmond S. The development of the Index of Complexity, Outcome and Need (ICON). *J Orthod.* 2000;27:149–162.
- [30] Sipilä K, Ensio K, Hanhela H. Occlusal characteristics in subjects with facial pain compared to a pain-free control group. *Cranio.* 2006;24:245–251.
- [31] Mohlin BO, Derweduwen K, Pilley R, et al. Malocclusion and temporomandibular disorder: a comparison of adolescents with moderate to severe dysfunction with those without signs and symptoms of temporomandibular disorder and their further development to 30 years of age. *Angle Orthod.* 2004;74:319–327.
- [32] Nishiyama A, Kino K, Sugisaki M, et al. Influence of psychosocial factors and habitual behavior in temporomandibular disorder-related symptoms in a working population in Japan. *Open Dent J.* 2012;6:240–247.
- [33] Al-Ahmad H, Al-Bitar Z. The effect of temporomandibular disorders on condition-specific quality of life in patients with dentofacial deformities. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2014;117:293–301.
- [34] Rusanen J, Silvola AS, Tolvanen M, et al. Pathways between temporomandibular disorders, occlusal characteristics, facial pain, and oral health-related quality of life among patients with severe malocclusion. *Eur J Orthod.* 2012;34:512–517.
- [35] Silvola AS, Varimo M, Tolvanen M, et al. Dental esthetics and quality of life in adults with severe malocclusion before and after treatment. *Angle Orthod.* 2014;84:594–599.
- [36] Yekkalam N, Wänman A. Prevalence of signs and symptoms indicative of temporomandibular disorders and headaches in 35-, 50-, 65- and 75-year-olds living in Västerbotten, Sweden. *Acta Odontol Scand.* 2014;72:458–465.
- [37] John MT, Dworkin SF, Mancl LA. Reliability of clinical temporomandibular disorder diagnoses. *Pain.* 2005;118:61–69.
- [38] Templeton K, Powell R, Moore M, et al. Are the Peer Assessment Rating Index and the Index of Treatment Complexity, Outcome, and Need suitable measures for orthognathic outcomes? *Eur J Orthod.* 2006;28:462–466.
- [39] Alanko OM, Svedström-Oristo AL, Tuomisto MT. Patients' perceptions of orthognathic treatment, well-being, and psychological or psychiatric status: a systematic review. *Acta Odontol Scand.* 2010;68:249–260.