

ORIGINAL ARTICLE

The cost of survival: an exploration of colorectal cancer survivors' experiences of pain

Amanda Drury^a , Sheila Payne^b  and Anne-Marie Brady^a 

^aSchool of Nursing & Midwifery, Faculty of Health Sciences, Trinity College Dublin, Dublin, Ireland; ^bInternational Observatory on End of Life Care, Division of Health Research, Lancaster University, Lancaster, UK

ABSTRACT

Background: The Institute of Medicine report 'From Cancer Patient to Cancer Survivor' has drawn widespread attention to the experiences of cancer survivors. Research examining the symptom experiences of survivors are proliferative within the literature but limited by samples which include multiple tumor groups and varying inclusion criteria. This cross-sectional quantitative study seeks to examine pain and quality of life (QoL) in the context of colorectal cancer (CRC) survivorship, as defined by the Institute of Medicine.

Material and methods: A purposive sample of CRC survivors (n = 252) attending hospitals and cancer support centers in the Republic of Ireland were recruited between September 2014 and January 2016. Self-rated health (SRH), QoL and pain were assessed in the sample using the EuroQOL questionnaire, the Functional Assessment of Therapy-Colorectal (FACT-C) questionnaire, and symptom experience items.

Results: One hundred participants (40%) indicated they had pain on the day of the survey or in the past week. Of those with pain, many also experienced a lack of energy (95%), bowel dysfunction (74%), sleep disturbance (76%) or interference with their ability to enjoy life (75%). Pain was associated with younger age, female gender, current chemotherapy treatment, and previous radiotherapy treatment. Although participants reported positive QoL scores, statistical analysis revealed pain was linked to significantly poorer SRH and overall QoL, and poorer physical, emotional, functional, social/family and CRC-specific wellbeing, compared to those who did not indicate pain.

Discussion: Pain was experienced by almost two fifths of CRC survivors up to five years after treatment and was associated with poorer SRH and QoL. In light of these findings, healthcare professionals must endeavor to manage cancer survivors' needs in a manner which is cognizant of the burden of pain and in the context of other symptoms and morbidities experienced by long-term cancer survivors.

ARTICLE HISTORY

Received 15 August 2016

Accepted 14 November 2016

Pain is a complex, multifactorial phenomenon which impacts physical, psychological and social domains of living [1]. Modern evolution of cancer treatments means many cancer patients may look forward to long-term cancer survivorship, as cancer is managed as a chronic illness. However, growing rates of cancer survivorship bring new concerns, as the chronic effects of cancer treatment gain prominence. A growing discourse surrounding the unmet needs of cancer survivors has emerged since the publication of 'From Cancer Patient to Cancer Survivor: Lost in Transition' [2] a decade ago.

Previous studies concluded that colorectal cancer (CRC) survivors regain quality of life (QoL) of similar or greater levels than those reported by normative populations [3–5]. The positive reframing of QoL in the presence of chronic symptoms is a widely discussed and accepted concept within the chronic illness literature [3,6]. However, it is estimated one third of cancer survivors experience pain following curative cancer treatment [5,7–11], compared to almost two thirds of those receiving anti-cancer treatment, or living with advanced, metastatic or terminal cancer [7]. Pain is ranked as

one of the most common symptoms experienced by cancer survivors [4,9,10], and has been associated with poorer physical [12] functional [11–14], psychological [14,15] and overall [16] QoL, and may contribute to social isolation in survivorship [17].

Several factors have demonstrated an influence on the presence and intensity of pain, including age [9,13,18], female gender [10,11,13,14,16,18], ethnicity [13,14], symptom clusters [13], previous chemotherapy treatment [13,18], previous radiotherapy treatment [18], time since treatment [13,16], stage of disease [19] and income [11]. Furthermore, pain has been shown to occur in clusters with fatigue, insomnia [19], depression [8,11] and anxiety [11]. The influence of age on pain is the subject of debate. Although almost one third of cancer survivors report pain [5,7–11], approximately one tenth of older CRC survivors attribute pain to cancer-related factors [9], compared to almost two fifths of general CRC survivor samples [11]. Studies have revealed older cancer survivors are reluctant to attribute symptoms of pain, fatigue or cognitive dysfunction to cancer [17], whereas age-related factors have greater predictive

ability than cancer-related factors for symptoms of pain, low energy levels and weakness [13]. Previous research suggests that ageing does not influence increased risk for pain, fatigue or insomnia in elderly cancer survivors [19]; however, no consensus has been reached regarding the intensity or frequency of pain flares [15,18].

Although under-treatment of pain during cancer treatment has reduced over the past decade [20], the prevalence of pain and its impact on CRC survivors' QoL in the survivorship period remains poorly evaluated. Many studies of cancer survivors' pain focus on samples which include multiple tumor groups and varying inclusion criteria [8,9,13–15], resulting in small samples of each tumor group or a lack of analysis of data pertaining to particular tumor groups. Meanwhile, studies which evaluate the pain of CRC survivors' have limited generalizability and scope due to small samples sizes [11,16], lack of clear definition of cancer survivorship [15], or sampling of older [12,19], early stage (<2 years) [18,19] or long-term cancer survivors (>5 years) [12]. Therefore, the present study sought to explore the prevalence and experience of pain in a sample of CRC survivors up to five years following diagnosis.

Material and methods

Participant selection

The current study is part of a larger mixed methods study to examine CRC survivors' QoL and healthcare experiences. This report details the results of the cross-sectional survey of CRC survivors attending three hospitals and 21 cancer support centers for routine follow-up care in the Republic of Ireland. All adult CRC survivors who were between six months and five years post-diagnosis, over the age of 18 years, resident in the Republic of Ireland, able to speak, read and comprehend English and able to provide informed consent were eligible to participate in the study. In keeping with the Institute of Medicine [2] definition of cancer survivorship, CRC survivors were considered eligible for participation in the study regardless of their disease status. To minimize the effect of acute toxicities on the study findings, CRC survivors were excluded if they were receiving primary cancer treatment, or less than six months post-diagnosis. Ethical approval for this study was granted by the Research Ethics Committees of the participating hospitals and university.

Data collection

Cancer-related QoL in the seven days preceding the study was measured using the 36-item Functional Assessment of Cancer Therapy-Colorectal Cancer (FACT-C) questionnaire [21]. The questionnaire evaluates four subscales of the core FACT-General (FACT-G) questionnaire; physical wellbeing, social wellbeing, emotional wellbeing, and functional wellbeing subscales, and the CRC concerns subscale. FACT-C items are assessed using a Likert-scale format, with item scores ranging from 0, 'not at all' to 4, 'very much', higher scores on each subscale reflect better QoL.

Cancer survivors' self-rated health (SRH) was assessed using the five-item EuroQOL EQ-5D-5L questionnaire and visual analog scale which measures SRH on the day of the survey on a scale of 0–100 [22]. Each health dimension of the EuroQOL questionnaire (Mobility, Self-Care, Usual Activities, Pain/Discomfort, and Anxiety/Depression) is measured on a five-level Likert scale, ranging from 0, 'no problem,' to 4, 'extreme problems'. Both the EuroQOL and the FACT-C questionnaires are widely used and have demonstrated validity and reliability with CRC populations.

The presence and intensity of pain was assessed: (1) on the day of the survey using the EuroQOL item 'Pain/Discomfort'; and (2) in the week prior to the survey using the FACT-C item 'in the past seven days, I have pain'. Each item was rated on a five-point Likert scale, with higher scores reflecting greater pain. Clinical and demographic characteristics assessed in the questionnaire included diagnosis, previous and current treatments, time since diagnosis, age, gender, living arrangements, employment status, ethnicity and healthcare insurance status.

All eligible CRC survivors identified by gatekeepers during the study period between October 2014 and January 2016 were offered written study information and invited to participate. Consenting participants were invited to complete an online or paper-based survey. Reminder letters were sent to those who had not returned a completed questionnaire after four weeks.

Data analysis

For the purpose of analysis, participants who had missing responses to the EuroQOL and FACT-C pain items were excluded. Incomplete data from the FACT-C and EuroQOL questionnaires were recorded as missing and excluded from relevant analysis. Responses to the EuroQOL and FACT-C pain items were collapsed into binary items, a pain score of 0 was coded as 'no pain', whereas scores ranging between 1 and 4 were coded as 'any pain' in the corresponding binary items 'pain today' (EuroQOL) or 'pain in the past week' (FACT-C).

Data was analyzed using SPSS v21. Cronbach's alpha was used to assess the internal consistency of the FACT-C scale and subscales. Descriptive statistics were used to characterize the demographic and cancer-related attributes of the sample. The prevalence of pain within the sample are reported using proportions. QoL and pain intensity scores are reported using descriptive measures of center and spread. Interval estimates of the proportion of the population experiencing pain and population mean scores for QoL and SRH are reported using 95% confidence intervals.

The relationship between pain and demographic and medical characteristics were examined using χ^2 analysis. Where significant values were recorded, odds ratios (ORs) were calculated using a standardized formula (see Table 3). The results of t-tests to examine differences in SRH and QoL between groups with and without pain on the day of the survey are reported, and Spearman's Rank Order Correlations were calculated to estimate the relationship between pain and QoL. All analysis were undertaken at the level $\alpha \leq 0.05$ level of significance unless otherwise stated.

Table 1. Demographic and clinical characteristics of respondents.

Characteristic	Response	n	%	N
Age groups	≤54 years	32	13	239
	55–64 years	50	21	
	65–74 years	107	45	
	≥75 years	50	21	
Gender	Male	132	55	240
	Female	108	45	
Health insurance	Medical card	144	60	241
	Private health insurance	121	50	
	None	17	7	
Employment status	Retired	114	48	239
	Other	57	24	
	Full/part-time employment	51	21	
	Unemployed and unable to work for health reasons	17	7	
Diagnosis	Colon	159	64	248
	Rectum	60	24	
	Other	29	12	
Time since diagnosis	≤1 year	39	17	234
	1–2 years	57	24	
	2–3 years	48	21	
	3–4 years	38	16	
	4–5 years	52	22	
Treatment history	Surgery	228	91	250
	Chemotherapy	151	60	
	Radiotherapy	62	25	
Current treatment	None	225	91	247
	Chemotherapy	18	7	
	Other	11	5	
Cancer status	Remission	199	84	236
	Still present	19	8	
	Uncertain	18	8	

Results

Sample characteristics

Of the 404 CRC survivors screened and eligible for the study, 304 provided informed consent and returned a completed questionnaire (raw response rate: 75.2%). Fifty-two participants failed to complete one or both of the EuroQOL and FACT-C pain items and were therefore excluded. A final sample of 252 CRC survivors were included (response rate: 62.3%). Cronbach alpha results for the FACT-C subscales were acceptable at $\alpha \geq 0.74$, and for the FACT-G and FACT-C scales at $\alpha = 0.94$.

The demographic characteristics and the clinical characteristics of the sample are presented in Table 1. The mean age of participants was 66.4 years [standard deviation (SD) = 11.4]. The average time since diagnosis for the sample was 3.0 years (SD = 1.4). Fifty-five percent were male, 80% lived with family or friends and more than half had private health insurance (50%) or a medical card (60%), which allows the holder to receive healthcare free of charge or at a discounted rate in the public healthcare system. Almost half of participants were retired (48%).

Pain

Frequencies and mean pain ratings reported by the participants are reported in Table 2. On the day of the survey, 90 participants (36%, CI 30–42%) reported pain, while 77 participants had experienced pain in the seven days prior to the study (31%, CI 25–36%). The low mean pain scores reported by the sample on the day of the survey ($\bar{x}=0.5$, SD = 0.7) and in the week prior to the survey ($\bar{x}=0.5$, SD = 0.8) are

reflective of the large proportions reporting no pain or low-moderate levels of pain.

Overall, 40% ($n = 100$) reported pain at one or both time points. Figure 1 presents the frequency of the most commonly reported symptoms experienced by CRC survivors who had any pain. Compared to participants who did not experience pain, those with pain were more likely to report a lack of energy ($p \leq 0.0005$, OR = 17.8), body image disturbance ($p = 0.002$, OR = 2.8), or an inability to work ($p \leq 0.0005$, OR = 5.6). High proportions of those with pain indicated challenges with enjoyment of hobbies (79%, $n = 78$), ability to work (79%, $n = 77$) and enjoyment of life (75%, $n = 73$). In total 76% of participants with pain were discontent with their QoL, compared to 41% of those with no pain. A greater proportion of CRC survivors with pain were more likely to indicate problems with all FACT-C and symptom items ($p \leq 0.05$), with six exceptions; feeling close to one's partner ($p = 0.236$), satisfaction with one's sex life ($p = 0.178$), difficulty urinating ($p = 0.068$), urinary frequency ($p = 0.118$), fear of cancer recurrence ($p = 0.269$) and fear of cancer spread ($p = 0.078$).

Table 3 presents the results of χ^2 analysis to determine differences between CRC survivors based on demographic and medical characteristics. Younger age ($p = 0.003$, OR = 2.3), receipt of current anti-cancer treatment ($p = 0.002$, OR = 3.5), previous chemotherapy treatment ($p \leq 0.0005$, OR = 3.6) and previous radiotherapy treatment ($p = 0.006$, OR = 2.2) were each significantly associated with cancer survivors experience of pain on the day of the survey. Likewise, younger age ($p = 0.027$, OR = 1.8), receipt of current anti-cancer treatment ($p = 0.001$, OR = 4.0) and previous chemotherapy treatment ($p = 0.001$, OR = 2.8) were significantly

Table 2. Results of descriptive frequencies and central measures of pain.

Characteristic	Total sample			Year 1			Year 2			Year 3			Year 4			Year 5		
	Response	Mean (SD)	n	%	Mean (SD)	n	%	Mean (SD)	n	%	Mean (SD)	n	%	Mean (SD)	n	%	Mean (SD)	n
Pain – today (range 0–4)	None	0.5 (0.7)	162	64	0.5 (0.7)	25	64	0.5 (0.7)	37	65	0.4 (0.6)	33	69	0.6 (0.9)	23	61	0.4 (0.7)	34
	Slight		65	26		10	26		14	25		12	25		9	24		14
	Moderate		21	8		4	10		6	11		3	6		3	8		3
	Severe		4	2		0	0		0	0		0	0		3	8		1
Pain – past week (range 0–4)	Extreme		0	0		0	0		0	0		0	0		0	0		0
	Not at all	0.5 (0.8)	175	69	0.4 (0.7)	26	67	0.5 (0.8)	39	68	0.5 (0.9)	32	67	0.5 (1.0)	26	68	0.3 (0.7)	41
	A little bit		51	20		9	23		12	21		12	25		7	18		6
	Somewhat		16	6		4	10		3	5		1	2		3	8		4
	Quite a bit		8	3		0	0		3	5		2	4		1	3		1
	Very much		2	1		0	0		0	0		1	2		1	3		0
Total			252	100		39	100		57	100		48	100		38	100		52

associated with cancer survivors experience of pain during the week prior to the survey.

Quality of life

Table 4 presents results of descriptive and inferential analysis of QoL and SRH for the sample based on pain experience. The sample reported positive perceptions of their health ($\bar{x}=81.4$, $SD=16.2$) and CRC-related QoL (FACT-C: $\bar{x}=111.9$, $SD=18.9$). CRC survivors who reported pain on the day of the survey and in the week preceding the survey reported significantly poorer SRH and QoL scores on all scales ($p \leq 0.05$). The results of correlation analysis between pain and QoL indicators are presented in Table 5. Increasing pain today or in the past week was negatively associated with SRH and QoL on all subscales ($p \leq 0.05$). The relationships between pain today or in the past week and QoL were moderate for all subscales with two exceptions. Physical well-being had a strong correlation with pain on the day of the survey ($r_s=-0.7$) and in the past week ($r_s=-0.7$). Social well-being had a weak correlation with pain on the day of the survey ($r_s=-0.3$) and in the past week ($r_s=-0.3$).

Discussion

This study sought to explore the prevalence and experience of pain in a sample of CRC survivors up to five years following diagnosis and the effect of pain on CRC survivors' QoL. Two fifths of the sample reported pain on the day of the survey or in the past week. When the groups were compared based on the demographic and medical variables under study, CRC survivors who reported pain today and pain in the last week were more likely to be younger, receiving current anti-cancer treatment and have previous chemotherapy treatment. Previous radiotherapy treatment was associated with experience of pain on the day of the survey only. CRC survivors who reported pain on the day of the survey or the week preceding participation in the survey indicated significantly poorer SRH, lower overall QoL scores, worse physical, social, emotional and functional wellbeing and a greater degree of CRC-specific QoL concerns compared to CRC survivors who did not report pain.

The prevalence of pain in this sample of CRC survivors mirror the findings of previous research of cancer survivors which include multiple tumor groups [5,7–11], albeit slightly higher than the prevalence reported in CRC survivor samples of other studies [5,9]. However, the finding that gender and time since diagnosis are not associated with CRC survivors' experience of pain refutes the findings of several studies [10,11,13,14,16,18]. These differences may be attributed to differences in study designs, as studies of CRC survivors tend to have had smaller samples, and may be at risk of type I errors [10,11,16], whereas others examined short-term [18], long-term [13] and diverse groups of cancer survivors [13,14].

The impact of age on cancer survivors' experiences of pain is debatable. The nature of CRC means the profile of survivors is skewed to middle-older age males and females. This study has found younger cancer survivors were twice as likely as




Figure 1. Clustered bar chart of the 10 most common symptoms experienced by colorectal cancer survivors who report pain and comparison with the proportions of those without pain who report these symptoms.

Table 3. Results of chi-square analysis and odds ratios comparing pain experiences of colorectal cancer survivors grouped by demographic and clinical characteristics.

Characteristic	Reference group	Comparison group(s)	Pain – today				Pain – past week			
			χ ²	φ	p	OR ^a	χ ²	φ	p	OR ^a
Age	≤65	>65 years	8.7	–0.2	0.003**	2.3	4.9	–0.1	0.027*	1.8
Gender	Male	Female	2.8	0.1	0.095		0.8	0.1	0.386	
Diagnosis	Colon	Rectum, other	0.6	0.1	0.729		1.2	0.1	0.562	
Healthcare insurance status	Private insurance	None	0.3	0.0	0.555		0.2	0.0	0.642	
Primary treatments:										
Radiotherapy	Yes	No	7.5	0.2	0.006*	2.2	1.0	0.1	0.316	
Chemotherapy	Yes	No	19.2	0.3	0.000**	3.6	11.6	0.2	0.001**	2.8
Surgery	Yes	No	2.2	–0.9	0.140		0.4	0.0	0.524	
Current treatments	Yes	No	9.5	–0.2	0.002**	3.5	12.0	–0.2	0.001**	4.0
Time since diagnosis	≤2.5 years	> 2.5 years	0.0	0.0	0.920		0.4	0.0	0.508	

*p ≤ 0.05; **p ≤ 0.005.

^aOdds ratio formula: OR = {(n) Reference Group Reporting Pain [a] × (n) Comparison Group No Pain [d]} / {(n) Reference Group No Pain [b] × (n) Comparison Group Reported Pain [c]}

Table 4. Results of descriptive and inferential analysis to compare quality of life scores between colorectal cancer survivors who do and do not have pain.

Scale (score range)	Total sample			Pain response	Pain – today			Pain – past week		
	N	Mean (SD)	95% Confidence interval		N	Mean (SD)	t	N	Mean (SD)	t
Self-rated health (0–100)	247	81.4 (16.2)	79.4–83.4	No pain	159	86.6 (12.2)	6.7**	171	86.0 (13.1)	6.7**
				Pain	88	72.0 (18.3)		76	70.9 (17.7)	
Physical well-being (0–28)	250	24.6 (4.3)	24.1–25.1	No pain	162	26.5 (1.9)	9.2**	174	26.4 (2.1)	10.0**
				Pain	88	21.2 (5.2)		76	20.5 (5.0)	
Social well-being (0–28)	248	23.5 (5.3)	22.8–24.1	No pain	159	24.3 (5.1)	3.5**	171	24.3 (5.2)	3.5**
				Pain	89	22.0 (5.2)		77	21.8 (5.1)	
Emotional well-being (0–24)	250	20.1 (4.0)	19.6–20.5	No pain	160	21.0 (2.8)	4.8**	173	20.9 (3.2)	4.5**
				Pain	90	18.3 (5.0)		77	18.2 (4.9)	
Functional well-being (0–28)	248	21.1 (6.3)	20.4–21.9	No pain	159	23.1 (5.4)	6.9**	173	23.0 (5.7)	8.0**
				Pain	89	17.6 (6.3)		75	16.8 (5.6)	
Colorectal cancer subscale (0–28)	248	22.2 (4.9)	21.5–22.8	No pain	160	23.7 (3.9)	7.2**	172	23.5 (4.1)	7.0**
				Pain	88	19.3 (5.1)		76	19.0 (5.0)	
FACT – general (0–108)	243	89.3 (15.7)	87.3–91.3	No pain	156	95.1 (11.0)	7.8**	169	94.8 (12.0)	8.7**
				Pain	87	79.0 (17.4)		74	76.8 (15.9)	
FACT – colorectal (0–136)	240	111.9 (18.9)	109.5–114.3	No pain	155	119.3 (12.7)	8.3**	167	118.9 (14.1)	9.2**
				Pain	85	98.5 (21.1)		73	96.0 (19.1)	

*p ≤ 0.05; **p ≤ 0.005.

Table 5. Results of correlation analysis to establish the relationship between pain and quality of life in colorectal cancer survivors.

Scale	Pain – today		Pain – past week	
	n	r_s	n	r_s
Self-rated health	247	–0.5**	247	–0.5**
Physical well-being	250	–0.7**	250	–0.7**
Social well-being	248	–0.3**	248	–0.3**
Emotional well-being	250	–0.3**	250	–0.3**
Functional well-being	248	–0.5**	248	–0.5**
Colorectal cancer subscale	248	–0.5**	248	–0.4**
FACT – general	243	–0.5**	243	–0.6**
FACT – colorectal	240	–0.5**	240	–0.6**

* $p \leq 0.05$; ** $p \leq 0.005$.

Values shown are Spearman's Rho.

older cancer survivors to report pain on the day of the survey and in the week preceding the survey. Previous studies report older cancer survivors have expressed difficulty in ascribing pain and other symptoms to cancer- or age-related factors [9,11,17], whereas poorer QoL and greater symptom burden in younger cancer survivors have been attributed to fewer coping strategies and greater perceived threat of recurrence [4,6]. To understand this finding, it is not unreasonable to hypothesize that younger age, and fewer comorbidities may permit more aggressive treatment strategies, contributing to a higher rate of late toxicities [6]. This is tentatively supported by the findings of the current study which associate previous radiotherapy and current chemotherapy with a greater likelihood of experiencing pain. Furthermore, younger CRC survivors may also experience a greater disturbance in their work life arising from chronic effects. This study showed that those who experience pain were more likely to indicate they were unable to work or gain fulfillment from their work compared to those who did not experience pain.

Although the mean QoL scores reported within this study are comparable to population norms [3–5], pain was significantly associated with poorer SRH and QoL on all subscales. Furthermore, differences in FACT-C scores between those who did and did not report pain are clinically significant, with disparities greater than the minimally important differences suggested by Yost and colleagues [23]. In addition, almost 95% of those who reported pain also experienced a lack of energy in the week preceding the survey, this is comparable to previous studies [13]. Pain and fatigue are potential deterrents to activity and have implications for the psychological wellbeing of the cancer survivor. Pain and fatigue may signify potential recurrence or disease progression [24,25], whereas late effects of cancer may also contribute to social withdrawal, functional disability, resentment and anger in the long term [17].

Strengths/limitations

The results must be interpreted with cognizance of study limitations. Although this sample was drawn from multiple research sites serving various socioeconomic groups, the cross-sectional nature presents a limitation to the generalizability of this study. Furthermore, the questionnaire items used to assess pain in this sample limit this study, as they

are not considered profound measures of pain. However, they are drawn from instruments which have established validity and reliability for CRC survivor populations. Moreover, the measures provide suitable means to assess point prevalence of pain, consistent with prior research and within the objectives of this study.

This study has addressed many of the previously discussed limitations of cancer survivorship research including clarity of disease status, definition of survivorship and purposive sampling strategies [11]. The strong response rate of 62.3% and rigor of inclusion criteria are strengths of this study. The sample is representative of CRC survivors in Ireland, as participants were recruited from routine follow-up care in hospitals and support centers which reflect the variety of settings in which CRC follow-up care is delivered in Ireland. The inclusion criteria ensure that the sample is consistent with the definition of survivorship set forth by the Institute of Medicine [2], and the symptoms reported are not contaminated by acute consequences of treatment. The homogenous sample of CRC survivors ensure the results are not biased by issues pursuant to other cancer survivor groups. The findings of this study add to the limited knowledge available regarding CRC survivors' experience of pain and its impact on QoL.

These findings present a basis to direct larger scale epidemiological research of QoL and symptom prevalence among CRC survivors. The results of this work guide the development of a qualitative research approach within an ongoing body of mixed methods research. The qualitative component of this study provides an opportunity to fully evaluate CRC survivors' symptom experiences, providing depth, context and coherence to the picture painted by these results.

Conclusion

The findings of this study add to the growing evidence base surrounding symptom burden and management in CRC survivors. The high prevalence of symptoms which co-occur with pain and the association between pain and poorer QoL is a cause for concern, and one which healthcare professionals must remain mindful of. It is not enough to assess and manage CRC survivors' pain in isolation. Indeed, considering pain within the bigger picture of survivors' comorbidities and cancer-related complaints is imperative to address their needs. Such an approach will ensure healthcare professionals may develop appropriate pain management strategies which will assist CRC survivors to sustain reasonable QoL levels in the long term.

Acknowledgments

The authors would like to thank the physicians, nurses and staff at the Colorectal Cancer Surgery and Medical Oncology Departments of the participating hospitals, and the voluntary cancer support services who facilitated data collection.

Disclosure statement

The authors report no conflict of interest. The authors alone are responsible for the content and writing of this paper.

Funding

This research is supported by funding from the Health Research Board of Ireland [grant number HPF.2014.715].

ORCID

Amanda Drury  <http://orcid.org/0000-0002-1720-5673>

Sheila Payne  <http://orcid.org/0000-0001-6982-9181>

Anne-Marie Brady  <http://orcid.org/0000-0002-7112-6810>

References

- [1] Clark D. 'Total pain', disciplinary power and the body in the work of Cicely Saunders, 1958–1967. *Soc Sci Med*. 1999;49:727–736.
- [2] Institute of Medicine, National Research Council. From cancer patient to cancer survivor: lost in transition. Washington (DC): The National Academies Press, 2006.
- [3] Pucciarelli S, Del Bianco P, Toppan P, et al. Health-related quality of life outcomes in disease-free survivors of mid-low rectal cancer after curative surgery. *Ann Surg Oncol*. 2008;15:1846–1854.
- [4] Arndt V, Merx H, Stegmaier C, et al. Restrictions in quality of life in colorectal cancer patients over three years after diagnosis: a population based study. *Eur J Cancer*. 2006;42:1848–1857.
- [5] Rauch P, Miny J, Conroy T, et al. Quality of life among disease-free survivors of rectal cancer. *J Clin Oncol*. 2004;22:354–360.
- [6] Jansen L, Herrmann A, Stegmaier C, et al. Health-related quality of life during the 10 years after diagnosis of colorectal cancer: a population-based study. *J Clin Oncol*. 2011;29:3263–3269.
- [7] van den Beuken-van Everdingen M, de Rijke J, Kessels A, et al. Prevalence of pain in patients with cancer: a systematic review of the past 40 years. *Ann Oncol*. 2007;18:1437–1449.
- [8] Moye J, June A, Martin LA, et al. Pain is prevalent and persisting in cancer survivors: differential factors across age groups. *J Geriatr Oncol*. 2014;5:190–196.
- [9] Deimling G, Sterns S, Bowman K. The health of older-adult, long-term cancer survivors. *Cancer Nurs*. 2005;28:415–424.
- [10] Phipps E, Braitman LE, Stites S, et al. Quality of life and symptom attribution in long-term colon cancer survivors. *J Eval Clin Pract*. 2008;14:254–258.
- [11] Lowery AE, Starr T, Dhingra LK, et al. Frequency, characteristics, and correlates of pain in a pilot study of colorectal cancer survivors 1–10 years post-treatment. *Pain Med* (Malden, Mass). 2013;14:1673–1680.
- [12] Koltyn K, Colbert L, Trentham-Dietz A. Pain and function in older, long-term colorectal cancer survivors. *J Pain*. 2009;10:57.
- [13] Deimling GT, Bowman KF, Wagner LJ. The effects of cancer-related pain and fatigue on functioning of older adult, long-term cancer survivors. *Cancer Nurs*. 2007;30:421–433.
- [14] Green CR, Hart-Johnson T, Loeffler DR. Cancer-related chronic pain examining quality of life in diverse cancer survivors. *Cancer*. 2011;117:1994–2003.
- [15] Green CR, Hart-Johnson T. Cancer pain: an age-based analysis. *Pain Med*. 2010;11:1525–1536.
- [16] Esnaola NF, Cantor SB, Johnson ML, et al. Pain and quality of life after treatment in patients with locally recurrent rectal cancer. *J Clin Oncol*. 2002;20:4361–4367.
- [17] Foley KL, Farmer DF, Petronis VM, et al. A qualitative exploration of the cancer experience among long-term survivors: comparisons by cancer type, ethnicity, gender, and age. *Psycho-Oncology*. 2006;15:248–258.
- [18] Kenzik K, Pisu M, Johns SA, et al. Unresolved pain interference among colorectal cancer survivors: implications for patient care and outcomes. *Pain Med*. 2015;16:1410–1425.
- [19] Kozachik SL, Bandeen-Roche K. Predictors of patterns of pain, fatigue, and insomnia during the first year after a cancer diagnosis in the elderly. *Cancer Nurs*. 2008;31:334–344.
- [20] Greco MT, Roberto A, Corli O, et al. Quality of cancer pain management: an update of a systematic review of undertreatment of patients with cancer. *J Clin Oncol*. 2014;32:4149–4154.
- [21] Ward WL, Hahn EA, Mo F, et al. Reliability and validity of the Functional Assessment of Cancer Therapy-Colorectal (FACT-C) quality of life instrument. *Qual Life Res*. 1999;8:181–195.
- [22] Brooks R. EuroQol: the current state of play. *Health Policy* (Amsterdam, Netherlands). 1996;37:53–72.
- [23] Yost KJ, Cella D, Chawla A, et al. Minimally important differences were estimated for the Functional Assessment of Cancer Therapy-Colorectal (FACT-C) instrument using a combination of distribution- and anchor-based approaches. *J Clin Epidemiol*. 2005;58:1241–1251.
- [24] LeMay K, Wilson KG, Buenger U, et al. Fear of pain in patients with advanced cancer or in patients with chronic noncancer pain. *Clin J Pain*. 2011;27:116–124.
- [25] Beech N, Arber A, Faithfull S. Restoring a sense of wellness following colorectal cancer: a grounded theory. *J Adv Nurs*. 2012;68:1134–1144.