

Pathology diagnosis is achieved through biopsy. As in our case, biopsies do not always yield the expected results. We should not forget that this test may also sometimes give false positive results, especially in low-grade AS that cannot be readily distinguished from radiotherapy-induced changes [3].

Standard therapy entails surgery with complete tumour resection. However, as in our case, chemotherapy becomes the first treatment of choice when surgery is not possible.

Results are far from alone promising for the most widely used drugs. Anthracyclines alone or with iphosphamide have led to disease control after several months (between 7 and 24 months) [8], always in combination with previous surgery. Over the past few years, paclitaxel has been used for used in advanced stage and/or metastatic AS [9,10].

In our case, the weekly taxane kept the disease under control for five months. Our case reveals the sound profile of paclitaxel as a single agent in the initial treatment of unresectable, radiotherapy-induced AS in an anthracycline-naïve patient.

## References

- [1] De Bree E, Van Coevorden F, Peterse J, Russell N, Rutgers E. Bilateral angiosarcoma of the breast after conservative treatment of bilateral invasive carcinoma: Genetic predisposition? *EJSO* 2002;28:392–5.
- [2] Luini A, Gatti G, Galimberti V, Zurrida S, Intra M, Gentilini O, et al. Conservative treatment of breast cancer: Its evolution. *Breast Cancer Res Treat* 2005;94:195–8.
- [3] Moe M, Bertelli G. Breast angiosarcoma following lumpectomy and radiotherapy for breast cancer: A case with short latent period and false negative result on biopsies. *Ann Oncol* 2007;18:801–9.
- [4] Deutsch M, Sayfan E. Angiosarcoma of the breast occurring soon after lumpectomy and breast irradiation for infiltrating ductal carcinoma: A case report. *Am J Clin Oncol* 2003;49: 532–8.
- [5] Strobbe LJ, Peterse HL, van Tinteren H, Wijnmaalen A, Rutgers EJ. Angiosarcoma of the breast after conservation therapy for invasive cancer, the incidence and outcome. An unforeseen sequela. *Breast Cancer Res Treat* 1998;47: 101–9.
- [6] Kirova YM, Vilcoq JR, Asselain B, Sastre-Garau X, Fourquet A. Radiation-induced sarcomas after radiotherapy for breast carcinoma. *Cancer* 2005;104:856–63.
- [7] Tomasini C, Grassi M, Pippione M. Cutaneous angiosarcoma arising in an irradiated breast. *Dermatology* 2004;209: 208–14.
- [8] Sher T, Hennessy BT, Valero V, Trent J, Broglio K, Woodward WA, et al. Angiosarcoma of the breast. *J Clin Oncol* 2006;24:9554 (Abstract).
- [9] Schlemmer M, Reichardt P, Verweij J, Hartmann JT, Judson I, Thyss A, et al. Paclitaxel in patients with advanced angiosarcomas. *J Clin Oncol* 2007;25:10033 (Abstract).
- [10] Penel N, Bui Nguyen B, Bay JO, Cupissol D, Ray-Coquard I, Piperno-Neumann S, et al. Weekly paclitaxel in metastatic angiosarcoma. A FNCLCC French Sarcoma Group (GSF-GETO) phase II trial. *J Clin Oncol* 2007;25:10002 (Abstract).

## A multimodality approach to reversible paraneoplastic encephalitis associated with ovarian teratomas

TIFFANY TANG<sup>1,2,\*</sup>, KAY YAW TAY<sup>2,\*</sup>, JOSIAH CHAI<sup>2</sup>, ANUPRIYA AGARWAL<sup>3</sup>, JEFFREY LOW<sup>3</sup>, EE LIAN LEE<sup>4</sup>, KHOON LEONG CHUAH<sup>5</sup>, KESAVAN S/O SITTAMPALAN<sup>6</sup>, CHIN FONG WONG<sup>6</sup>, JOSEP DALMAU<sup>7</sup>, JOANNE NGEOW<sup>1</sup>, LAY TIN SOH<sup>1</sup> & MIN-HAN TAN<sup>1,8</sup>

<sup>1</sup>Department of Medical Oncology, National Cancer Centre, Singapore, <sup>2</sup>Department of Neurology, National Neuroscience Institute, Singapore, <sup>3</sup>Department of Obstetrics and Gynaecology, National University Hospital, Singapore, <sup>4</sup>Department of Psychiatry, Singapore General Hospital, Singapore, <sup>5</sup>Department of Pathology, Tan Tock Seng Hospital, Singapore, <sup>6</sup>Department of Pathology, Singapore General Hospital, Singapore, <sup>7</sup>Division of Neuro-oncology, Department of Neurology, University of Pennsylvania, Philadelphia and <sup>8</sup>NCCS-VARI Laboratory of Translational Cancer Research, National Cancer Centre, Singapore.

Paraneoplastic encephalitis associated with ovarian teratomas and antibodies to the N-methyl-D-aspartate receptor (NMDA) is a recently described

phenomenon and resection of the tumour appears significant in achieving sustained neurological recovery [1]. We describe two patients who presented

Correspondence: Min-Han Tan, Department of Medical Oncology, National Cancer Centre, Singapore. Tel: +65 6436 8000. Fax: 65 6227 2759. E-mail: tan.min.han@nccs.com.sg (\* The authors contributed equally to this work)

(Received 16 November 2008; accepted 16 February 2009)

ISSN 0284-186X print/ISSN 1651-226X online © 2009 Informa UK Ltd. (Informa Healthcare, Taylor & Francis AS)  
DOI: 10.1080/02841860902829221

with paraneoplastic encephalitis associated with immature teratomas and discuss the role of chemotherapy in the treatment of this condition.

The first patient was a 15-year-old girl who presented with lower abdominal pain. A CT scan of her pelvis revealed bilateral ovarian masses suggestive of teratomas. These were resected and histology of the left ovarian tumour revealed a grade (2/3) immature ovarian teratoma and the right ovarian cyst had features suggestive of a benign cystic teratoma. With further omental foci of immature teratoma she had a final pathologic stage of IIIA. Twenty-five days post operatively, she began to exhibit several neuro-psychiatric symptoms which include marked short-term memory loss, paranoid psychosis, agitation, catatonia and myoclonic jerks. No organic cause was found to explain her symptoms; extensive metabolic and imaging studies were normal, serum anti-neuronal antibodies (anti-Hu, anti-Yo and anti-Ri) were negative and an EEG done was unremarkable. After failing treatment with anti-psychotic agents, she was started on standard BEP (bleomycin, etoposide, cisplatin) chemotherapy; 14 days after her symptoms began. Two weeks after initiation of chemotherapy her paranoid psychosis resolved. Six weeks later and after completion of three cycles of BEP, her memory returned to normal and she recovered completely.

The second patient was a 24-year-old woman who presented with vomiting and diarrhoea preceding fever, fits and altered mental status. On examination, she had gaze-evoked nystagmus and dystonia. She was also seen with intermittent eye deviation and version to the right associated with right upper limb rigidity and clenching of her fist. She would shout, speak incomprehensibly and she would not respond appropriately to external verbal or visual cues. CSF examination




Figure 1. MRI showing T2 hyperintensities in bilateral cerebellar hemispheres in Patient 2




Figure 2. CT pelvis showing the left sided teratoma in Patient 2

revealed leukocyte pleocytosis, with a nucleated cell count of 14/ul (95% lymphocytes, 5% monocytes, 0% neutrophils); CSF glucose was 3.9 mmol/L with a blood glucose level of 5.4 mmol/L; CSF protein was within normal range. Microbiological assessment of CSF for bacteria, mycobacterium, fungi and viruses were negative and MRI scans of the brain revealed several small T2 hyperintense foci in bilateral cerebellar hemispheres (Figure 1). The patient was treated empirically for bacterial and herpes simplex encephalitis and in view of her failure to respond to the initial antibiotic regimen, she was also treated for tuberculous meningoencephalitis. In search of a paraneoplastic cause to explain her symptoms, a CT scan of her thorax, abdomen and pelvis was performed. This revealed a 10.8 by 9.7 by 8.5 cm cystic lesion suggestive of a left-sided ovarian teratoma (Figure 2). Examination of the patient's serum for paraneoplastic markers (anti-Hu, anti-Yo and anti-Ri) was negative. Immunological characterisation of her serum and CSF demonstrated the presence of anti-NMDA receptor autoantibodies. She underwent a resection of the left-sided teratoma (Figure 3) and intra-operatively, a right ovarian cyst was seen and removed. Histology revealed a grade (3/3) immature teratoma and a mature teratoma respectively with a final



Figure 3. Macroscopic appearance of the left ovarian teratoma from Patient 2

Table I. Paraneoplastic Encephalitis associated with Immature Ovarian Teratomas (IOT)–Outcomes with and without chemotherapy.

| Reference      | Age/<br>Gender | Tumour Diagnosis and stage as<br>published                                         | Anti-Neuronal antibodies                                                                                 | Treatment                                                                                                  | Outcomes                                                                                                                       |
|----------------|----------------|------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------|
| Dalmau [1]     | Case 1 (30/F)  | 10 cm right IOT                                                                    | NMDA R (+)                                                                                               | Corticosteroids<br>Plasma exchange<br>IVIG                                                                 | Back to work as an internal<br>medicine resident                                                                               |
|                | Case 4 (17/F)  | 7 cm left IOT                                                                      | NMDA R (+)                                                                                               | Tumour Removal<br>Plasma Exchange<br>IVIG<br>Tumour<br>Removal Corticosteroids and<br>Cyclophosphamide     | Back to school with good grades                                                                                                |
|                | Case 7 (19/F)  | 22 cm right IOT                                                                    | NMDA R (+)                                                                                               | Corticosteroids<br>IVIG<br>Tumour<br>removal <b>Chemotherapy</b>                                           | Alternating insomnia/hypersomnia<br>for 7 months;<br>total recovery of motor function<br>but MMSE 24/30                        |
|                | Case 10 (14/F) | 1.9 cm left IOT                                                                    | NMDA R (+)                                                                                               | Tumour removal<br>Plasma Exchange<br>Corticosteroids IVIG                                                  | Unexpected death after mild<br>improvement                                                                                     |
| Aydiner [2]    | 39/F           | 6.8 cm benign ovarian<br>teratoma with 1 mm<br>focal area of immature<br>component | Anti-Hu and<br>Anti-Yo from<br>serum and CSF<br>negative                                                 | Tumour removal<br><b>Chemotherapy (BEP)</b>                                                                | Mood disorder controlled with<br>Lithium<br>Persistent amnesia                                                                 |
| Sansing [3]    | 34/F           | 7.9 cm left IOT                                                                    | NMDA R (+)                                                                                               | Tumour removal<br>Methylprednisolone<br>Plasma Exchange<br>IVIG<br>Cyclophosphamide<br><b>Chemotherapy</b> | Mild generalized weakness but<br>could walk<br>unaided. Normal<br>cognitive function, memory and<br>psychiatric<br>evaluation. |
| van Altena [4] | 32/F           | 5 cm IOT in both ovaries<br>Grade 1, Stage Ib                                      | Anti-Hu, Anti-Yo, Anti-Ri,<br>Anti-Tr,<br>Anti-amphi, Anti CV2,<br>anti-Ma2 in<br>CSF and serum negative | Bilateral<br>salpingo-oophorectomy only                                                                    | Normal Neurological and<br>Performance<br>status                                                                               |
| Lee [5]        | 15/F           | 20 cm left IOT and 12 cm<br>mature teratoma in the right<br>ovary                  | Not reported                                                                                             | Tumour removal<br>Methylprednisolone                                                                       | Expressive aphasia Disorientation in<br>time<br>and date Continued to drop out of<br>school                                    |
| Nokura [6]     | 19/F           | IOT                                                                                | Negative for Anti-Hu                                                                                     | Tumour removal Corticosteroids                                                                             | Memory problem and poor score on<br>IQ<br>testing at 6 months                                                                  |

pathologic stage of IA. Her neuropsychiatric symptoms failed to abate despite surgery and a course of intravenous immunoglobulins. Fourteen days post-operatively, she was started on adjuvant chemotherapy with a standard chemotherapeutic regime of BEP, receiving a total of three cycles. Her dystonia became less prominent rapidly after initiation of her first cycle, but it was only after 20 days after commencement of chemotherapy that she regained orientation and was able to answer questions in a relevant and appropriate manner. She returned to work four months after completion of chemotherapy.

These two cases highlight the role of chemotherapy as an adjunct to surgery in a multimodality approach to paraneoplastic encephalitis associated with teratoma. In the first patient, the temporal association is definitive where neuropsychiatric symptoms only developed after the teratomas were resected. These symptoms resolved completely following chemotherapy. Given that tumour foci were identified on the omentum and in the peritoneal fluid, it is conceivable there may have been residual disease after surgery resulting in a delayed onset of encephalitis. This would further support the role of chemotherapy as an adjunct to surgery. In the second patient, since both surgery and intravenous immunoglobulin was administered prior to chemotherapy, one cannot exclude the possibility that her improvement may have represented a delayed effect of treatment. However, a strong temporal association between initiation of chemotherapy and symptom improvement was noted.

In a review of patients reported to have paraneoplastic encephalitis associated with immature ovarian teratomas in the English literature (Table I), only three of the ten patients received chemotherapy, with a fourth receiving cyclophosphamide only [1–6]. The grade and stage of the disease as well as the details of treatment

were however not reported for all ten patients so it is unclear whether chemotherapy would have been recommended [7,8]. The current difficulty in establishing ideal treatment modalities lies in the small number of patients that have been identified to date and further reports may be helpful to determine this.

### Acknowledgements

Dr. Min-Han Tan has received support for research by the Singapore Millenium Foundation.

### References

- [1] Dalmau J, Tüzün E, Wu HY, Masjuan J, Rossi JE, Sansing LH, et al. Paraneoplastic anti N-methyl-D aspartate receptor encephalitis associated with ovarian teratomas. *Ann Neurol* 2007;61:25–36.
- [2] Aydiner A, Gürvit H, Baral I. Paraneoplastic limbic encephalitis with immature ovarian teratoma—a case report. *J Neurooncol* 1998;37:63–6.
- [3] Sansing LH, Tüzün E, Ko MW, Baccon J, Lynch DR, Dalmau J. A patient with encephalitis associated with NMDA receptor antibodies. *Nat Clin Pract Neurol* 2007;3:291–6.
- [4] van Altena AM, Wijnberg GJ, Kolwijck E, de Hullu JA, Massuger LF. A patient with bilateral immature ovarian teratoma presenting with paraneoplastic encephalitis. *Gynecol Oncol* 2008;108:445–8.
- [5] Lee AC, Ou Y, Lee WK, Wong YC. Paraneoplastic limbic encephalitis masquerading as chronic behavioural disturbance in an adolescent girl. *Acta Paediatr* 2003;92:506–9.
- [6] Nokura K, Yamamoto H, Okawara Y, Koga H, Osawa H, Sakai K. Reversible limbic encephalitis caused by ovarian teratoma. *Acta Neurol Scand* 1997;95:367–73.
- [7] Williams S, Blessing JA, Liao SY, Ball H, Hanjani P. Adjuvant therapy of ovarian germ cell tumours with cisplatin, etoposide and bleomycin; a trial of the Gynecologic Oncology Group. *J Clin Oncol* 1994;12:701–6.
- [8] Gershenson DM, Morris M, Cangir A, Kavanagh JJ, Stringer CA, Wharton JT, et al. Treatment of malignant germ cell tumours of the ovary with bleomycin, etoposide and cisplatin. *J Clin Oncol* 1990;8:715–20.

## Somatic mutation of GNAQ gene is rare in common solid cancers and leukemias

HYEON SEOK EOM<sup>2</sup>, MIN SUNG KIM<sup>1</sup>, SOO YOUNG HUR<sup>3</sup>, NAM JIN YOO<sup>1</sup> & SUG HYUNG LEE<sup>1</sup>

<sup>1</sup>Department of Pathology, College of Medicine, The Catholic University of Korea, Seoul, Korea, <sup>2</sup>Hematology-Oncology Clinic, National Cancer Center of Korea, Goyang, Korea and <sup>3</sup>Department of Obstetrics/Gynecology, College of Medicine, The Catholic University of Korea, Seoul, Korea

Correspondence: Sug Hyung Lee, Department of Pathology, College of Medicine, The Catholic University of Korea, 505 Banpo-dong, Socho-gu, Seoul 137-701, Korea. Tel: +82 2 2258 7311. Fax: +82 2 2258 7765. E-mail: suhulee@catholic.ac.kr

(Received 16 November 2008; accepted 16 February 2009)

ISSN 0284-186X print/ISSN 1651-226X online © 2009 Informa UK Ltd. (Informa Healthcare, Taylor & Francis AS)  
DOI: 10.1080/02841860902882444