

ORIGINAL ARTICLE

The potential of proton beam radiation therapy in prostate cancer, other urological cancers and gynaecological cancers

BENGT JOHANSSON¹, MONA RIDDERHEIM² & BENGT GLIMELIUS^{3,4}

¹Department of General Oncology, Örebro University Hospital, Örebro, Sweden, ²Department of Oncology, University Hospital, Lund, Sweden, ³Department of Oncology, Radiology and Clinical Immunology, University Hospital, Uppsala, Sweden and ⁴Department of Oncology and Pathology, Karolinska Institutet, Stockholm, Sweden

Abstract

A group of Swedish oncologists and hospital physicists have estimated the number of patients in Sweden suitable for proton beam therapy. The estimations have been based on current statistics of tumour incidence, number of patients potentially eligible for radiation treatment, scientific support from clinical trials and model dose planning studies and knowledge of the dose-response relations of different tumours and normal tissues. In prostate cancer it is estimated that annually about 300 patients and in gynaecological cancer about 50 patients, are candidates for proton beam therapy. Owing to major uncertainties, it has not been possible to give an estimate of the number of potential patients with urinary bladder cancer.

The incidence of prostate cancer has increased during recent years, due to increased use of PSA testing. Recent statistics reveal over 9000 new cases annually in Sweden [1]. If public PSA screening is implemented the number will temporarily increase even more.

Median age at diagnosis is presently somewhat lower, about 70 years, than in the past and there is a stage migration towards more early stages T1–T2, reflecting the increase in PSA testing. This change has gone much further in countries recommending PSA screening, such as USA [2].

Cancer of the bladder is diagnosed in about 2300 patients annually [1]. Seventy per cent of these cases are superficial and quite amenable to some form of intravesical treatment. Patients with muscularly infiltrative bladder cancer are treated with surgery, radiotherapy and cytostatics or combinations of these.

About 2700 cases of gynaecological cancer are diagnosed in Sweden every year [1]. Of these about 1300 are corpus cancer, about 825 ovarian cancer and about 450 cervical cancer. Vaginal cancer and vulva cancer are more unusual, totalling about 160 cases together.

Prostate cancer

Optimal treatment of prostate cancer is controversial especially for early stages (T12N0M0) where conclusive clinical studies still are awaited. Different alternatives are expectancy until tumour-related symptoms arise, radical prostatectomy or radical radiation therapy. One Swedish randomized study, SPCG 4 [3,4], showed that radical prostatectomy reduced the risk of overall cancer-related deaths and the rate of distant metastases. Another cohort study with a follow-up time >20 years, where primary expectancy was applied, showed an increased cancer-related death rate in patients who survived for more than 15 years [5]. Thus, there is fairly strong evidence that reasonably young patients with an early prostate cancer should be offered curative treatment. The strongest evidence is then for surgery [4,6,7]. The choice of therapeutic strategy in a patient with a newly diagnosed early prostate cancer is still difficult [8–10].

In more advanced stages, like T3N0M0 (and in stages T2N0M0 with unfavourable prognostic signs) radiation therapy is given to control local tumour growth and risk of tumour dissemination and death [7,11]. The intermediate and high risk group of

Correspondence: Bengt Johansson, Department of General Oncology, Örebro University Hospital, S-701 85 Örebro, Sweden. E-mail: bengt.johansson@orebroll.se

(Received 23 August 2005; accepted 15 September 2005)

prostate cancer benefits from a high target dose (>78 Gy) than previously used [12], whereas there is little evidence for this in the low risk group [13]. Patients with distant metastases are commonly treated by hormonal therapy. Hormonal therapy is also frequently used as (neo-) adjuvant therapy in localized disease [7,14].

Radiation therapy may be external beam therapy, brachytherapy or a combination of both [7]. A special feature of all radiation therapy of the prostate is the fact that the prostate gland will move between each fraction, so that a security margin of 15–20 mm has to be added to the prostate gland to safely cover the whole prostate volume. The urethra runs through the prostate and the anterior rectal wall is located dorsally in direct contact with the prostate gland. These are the most important organs at risk.

In the report from the Swedish Council on Technology Assessment in Health Care (SBU), the practice of curatively intended radiation therapy on prostate cancers was measured in the year 2001. At least 1625 patients were treated annually with curative intent [15].

Early and late reactions from the urethra, the rectal wall and neurovascular structures are the dominant sites giving rise to adverse effects, well described in many publications [16–32]. Increased total doses will result in more side effects. More sophisticated radiation techniques, particularly compared to those used in the past, may reduce the side effects, but randomised studies which compare the level of side effects using different techniques are almost entirely lacking [7,33].

Clinical experience on proton beam therapy in prostate cancer

Using protons, a great deal of experience on prostate cancer has been published from Boston and Loma Linda. A survey of 1255 patients with mainly T1–T2 tumours treated at Loma Linda between 1991 and 1997 with protons of 74 cGy (cobalt-gray equivalent) showed bNED = 73% at eight years [34]. Freedom from grade three to four toxicity from the rectum and the bladder were 99%. In a previous publication on 911 patients, the role of grade two toxicity from the rectum was 3.5% and from the bladder 5.4% [35]. It was concluded that protons at a dose of 75 Gy yielded disease-free survival rates comparable with other forms of local radiation therapy, and with minimal morbidity, based, however, upon a non-randomised comparison.

In a randomised study from Boston between 1982 and 1992, 202 patients were treated with 50.4 Gy photons using a four-field box technique. Half the

patients were then given a photon boost of 17.2 Gy and the other half a proton boost of 25.2 Gy. After eight years, the local control was better in the proton group, 80% vs 60%, but the difference was not statistically significant ($p=0.09$). The difference was, however, statistically significant in a subgroup analysis of high grade tumours [36]. The incidence of rectal bleeding was higher in the proton group. Some urinary and gastrointestinal tract toxicity persisted [21].

Preliminary experience using carbon ions was recently reported in 175 patients treated between 2000 and 2003 at Chiba, Japan [37].

Comparative dose-planning model studies

A comparative study of dose planning with photons and protons by Lee et al. [38] showed improved TCP and NTCP with protons compared to photons. More recent studies [39] have also confirmed the potential advantages of proton beams which result in a more homogeneous dose distribution. Only a limited number of patients have been planned with both proton and photon techniques and biological models have not been used to evaluate to what extent proton therapy is better than modern photon techniques like IMRT.

An estimation of suitable number of patients for proton beam therapy in Sweden

If protons were generally available it is likely that more patients would be selected for this therapy, given the physical advantages in the proton beam, permitting escalated doses with improved tumour control or reduced toxicity. However, it is far from clear whether the dose distribution advantages would be sufficient to be cost-effective.

T1–T2. Even if randomised comparative studies are lacking, it is reasonable to consider radical prostatectomy, monobrachytherapy, external beam radiotherapy (>72 Gy) or a combination of external beam + brachytherapy as isoeffective, yielding a local tumour control of 80–85% [7]. There is support in the literature for improved tumour control with dose escalation (>75 Gy) [6,40]. Dose escalation, unless the radiation technique is improved with better normal tissue sparing, also increases the risk of adverse effects [30,33] and there is still limited knowledge about the magnitude of these late toxicity effects using modern techniques.

In this report we do not consider that protons will have sufficient advantage in T1–T2, low risk cases. There is now also a strong trend in favour of the unique features of brachytherapy with relative dose

sparing of the urethra and the anterior rectal wall, although this is not universally agreed upon. Brachytherapy also offers a short treatment time.

About 1/3 of the patients who undergo a radical prostatectomy have positive surgical margins. About 900 prostatectomies were done during 2004 in Sweden. For a fraction of these non-radically operated patients, we expect proton beam therapy to be indicated, due to the better physical dose distributions. Brachytherapy is not suitable for these patients. Thus, we estimate about 100 patients per year to be suitable for proton beam therapy.

T3. Approximately 1/3 of the patients are in clinical stage T3NX at detection. After lymphadenectomy 60% of these are T3N0M0. Clinical data support a better outcome when dose escalation (≥ 78 Gy) is applied. Prostatectomy is seldom possible and not recommended for this group. Approximately 40–50% of these patients have a life expectancy of less than 10 years and about 60% are treated by endocrine therapy only, so approximately 40% are considered for external beam radiotherapy.

T3 cases with a prostate volume exceeding about 50 cm³ are not suitable for brachytherapy because of pubic arch interferences. We estimate that about 200 patients in stage T3N0M0 with particular features are suitable for proton beam therapy.

Cost-effectiveness calculations

In a separate study a calculation of cost-effectiveness was performed [41]. The calculations were based on a 65 years male treated by either photons or protons. It was assumed, however, based on uncertain data from clinical studies, that proton beam treatment would result in 20% fewer tumour-related deaths and 40% fewer treatment-related side effects.

The risk of death from prostate cancer is 2.5% per year. The cost per QALY will then be 239 000 SEK. In the discussion above we assumed proton beam treatment to be used mainly for high risk cases where it would be more appropriate to assume a 5% annual risk of cancer death. This assumption will result in a cost of 151 000 SEK per gained QALY. Thus, this calculation reveals that protons are cost effective, especially for patients with a somewhat worse prognosis than average.

Future research areas

Before protocols for clinical studies of proton therapy are initiated it would be appropriate to perform comparative dose planning studies between IMRT photon and proton beam techniques. It could

also be possible to compare such studies with brachytherapy dose plans even if this is more complicated, due to dose inhomogeneities in the brachytherapy situation. These studies can predict the outcome in future clinical studies and thus aid in the dimension of the trials.

Recent radiobiological studies have estimated the α/β ratio for prostate cancer to be low (1.5–3) [42–44]. If this is true, hypo-fractionation with strictly conformal techniques like proton beam and brachytherapy could be very cost effective and also improve the therapeutic ratio. This issue must be evaluated in randomized protocols.

When a national proton facility is in clinical use all patients should be evaluated in prospective protocols with careful long-term follow-up. Some studies on more frequent occurring tumour groups could be evaluated in randomized protocols for more conclusive results. We consider patients after non-radical prostatectomy and high risk prostate cancers to be suitable in these aspects.

Summary assessment

We estimate that 300 patients every year will be suitable for proton beam therapy in Sweden, if treatment resources are available. Proton therapy dose escalation might give improved probability for tumour control without increased side effects. We consider large (T3), high risk tumours to be the most important group (about 200 patients per year). The cost for this treatment is estimated to 150 000 SEK per gained QALY.

Urinary bladder cancer

The role of radiotherapy in primary bladder cancer is unclear [45]. It has been used as pre-operative treatment to augment the likelihood of tumour control and as definitive treatment, either on its own or combined with cytostatics in cases where surgery is impossible or unsuitable. According to the review of literature undertaken recently by SBU [45], and an educational review [46], the literature on the subject is limited, with mostly small and inconclusive trials. Preoperative radiotherapy and surgery was standard treatment until about 1990, but this has since been removed in favour of cystectomy alone [45,47]. An SBU survey for 12 weeks in 2001 showed radiotherapy was given to about 50 bladder cancer patients against the primary tumour region. On an annual basis this means upwards of 200 patients irradiated. Just over 60% of these received “curative” radiation doses [48].

Clinical experience of proton beam therapy for urinary bladder cancer

There are no studies reported in which protons have been used for bladder cancer, but there have been four randomised studies comparing neutrons and photons. These studies [49–52] show that neutron irradiation does not increase tumour control or survival, and that it entails a heightened risk of serious side-effects [45].

Model studies

No studies have been reported. Since the small intestine and rectum are adjacent risk organs, it is possible that better dose distribution can be achieved with protons. It has not been studied whether this means greater tumour control through the feasibility of higher doses or through a reduction of long-term morbidity.

Assessment of the number of cases eligible for proton beam therapy

The number of cases eligible for proton beam therapy cannot be assessed, because the role of radiotherapy in the treatment of bladder cancer is poorly defined and no experience has been accumulated of model studies or of proton treatment of patients. If treatment is given, it would probably only be applied within clinical trials, apart from one or two exceptional patients where good dose distribution cannot be achieved with conventional irradiation.

Research needed

Model studies can give some idea of any potential for protons compared with 3D CRT and IMRT. There is a great need for clinical studies of radiotherapy for bladder cancer.

Summary assessment

It is estimated that between 100 and 150 bladder cancer patients annually in Sweden receive radiotherapy with a curative purpose. It is impossible to judge the fraction of these patients who could benefit from proton therapy.

Gynaecological cancer

Radiotherapy plays an important role in the treatment of cancer of the corpus and cervix uteri, vagina and vulva but is being used more and more rarely for ovarian cancer [53–55]. Radiotherapy still has a prominent role for cervical and vaginal cancer, both

primary and recurrent. In the SBU survey [48] 359 radiation treatments were given, including 158 (44%) as external radiotherapy against the primary tumour. On an annual basis this implies about 1500 treatments, including 660 external with a principally curative purpose. The latest figures give a diagnostic breakdown of eight ovarian cancer cases, 133 vulva and vaginal cancer, 190 cervical cancer and 310 corpus cancer.

Clinical experience of proton beam therapy for gynaecological cancer

Protons have been used for irradiation for gynaecological cancer and the first cancer patient in the world to be treated, for example, was a woman with recurrent cervical cancer [56]. Since then a couple of minor studies have been reported from Japan [57–59]. The patients treated have not been judged possible to treat with intracavitary radiation, owing to very large tumours, anatomic deviations or other difficult medical conditions. The results are judged favourable, better than those which external photon therapy can achieve, and on a par with those yielded by intracavitary radiotherapy at the corresponding stage (if this had been possible to give) [59]. As an alternative to a conventional brachytherapy boost [60], 16 patients have been treated with stereotactic photon radiotherapy. The treatments were well tolerated.

Model studies

In a model study better dose distributions were seen in a patient with stage IB squamous cell carcinoma of the uterine cervix [61]. In the above-mentioned clinical trial [60] a comparative dose-planning study was performed between brachytherapy and stereotactic radiotherapy (dynamic-arc and IMRT) using photons. The external techniques improved dose homogeneity to the target and reduced the maximum dose to the rectum, when compared to brachytherapy.

Assessment of the number of cases eligible for proton beam therapy

This is very difficult to assess. One or two special situations may primarily come to be considered, e.g. primarily advanced cervical cancer where brachytherapy is not possible owing to anatomic conditions or does not provide adequate tumour coverage (e.g. stage III with bilateral parametrium extent (about 10–20 cases per annum), in isolated local/regional recurrences (about 20 cases per annum) or in isolated cases of cancer of the vulva and vagina, where tumour coverage cannot be achieved with

interstitial, intracavitary or conventional external therapy (about five cases per annum). A maximum of 50 cases annually may come to be considered. However, if stereotactic techniques gain more popularity in the future, see [60], many more patients can be potential candidates.

Summary assessment

As boost therapy towards volumes at risk to contain tumour cells, brachytherapy has an established role since very long. In cases where brachytherapy is for some reason difficult to perform, an external boost with protons may be a valid alternative. This may be of importance in up to about 50 patients annually in Sweden. However, there is a potential for treatment of many more patients, if clinical studies show that external boost irradiation has advantages over brachytherapy.

References

- [1] Cancer Incidence in Sweden. Epidemiologiskt Centrum 2003; www.sos.se.
- [2] Albertsen PC. Is screening for prostate cancer with prostate specific antigen an appropriate public health measure? *Acta Oncol* 2005;44:255–64.
- [3] Holmberg L, Bill-Axelson A, Helgesen F, Salo JO, Folmerz P, Haggman M, et al. A randomized trial comparing radical prostatectomy with watchful waiting in early prostate cancer. *N Engl J Med* 2002;347:781–9.
- [4] Bill-Axelson A, Holmberg L, Ruutu M, Haggman M, Andersson SO, Bratell S, et al. Radical prostatectomy versus watchful waiting in early prostate cancer. *N Engl J Med* 2005;352:1977–84.
- [5] Johansson JE, Andren O, Andersson SO, Dickman PW, Holmberg L, HO A. Natural history of early, localized prostate cancer. *JAMA* 2004;291:2713–9.
- [6] Glimelius B, Steineck G. Low evidence of radiation therapy in prostate cancer—a plea for intensified scientific activity. *Acta Oncol* 2004;43:311–5.
- [7] Nilsson S, Widmark A, Norlén BJ. A systematic overview of radiation therapy effects in prostate cancer. *Acta Oncol* 2004;43:316–81.
- [8] Albertsen PC, Nease RF, Jr, Potosky AL. Assessment of patient preferences among men with prostate cancer. *J Urol* 1998;159:158–63.
- [9] Saigal CS, Gornbein J, Reid K, Litwin MS. Stability of time trade-off utilities for health states associated with the treatment of prostate cancer. *Qual Life Res* 2002;11:405–14.
- [10] Krahn M, Ritvo P, Irvine J, Tomlinson G, Bremner KE, Bezjak A, et al. Patient and community preferences for outcomes in prostate cancer: implications for clinical policy. *Med Care* 2003;41:153–64.
- [11] Foroudi F, Tyldesley S, Barbera L, Huang J, Mackillop WJ. Evidence-based estimate of appropriate radiotherapy utilization rate for prostate cancer. *Int J Radiat Oncol Biol Phys* 2003;55:51–63.
- [12] Pollack A, Zagars GK, Starkschall G, Antolak JA, Lee JJ, Huang E, et al. Prostate cancer radiation dose response: results of the M. D. Anderson phase III randomized trial. *Int J Radiat Oncol Biol Phys* 2002;53:1097–105.
- [13] Cheung R, Tucker SL, Lee AK, de Crevoisier R, Dong L, Kamat A, et al. Dose-response characteristics of low- and intermediate-risk prostate cancer treated with external beam radiotherapy. *Int J Radiat Oncol Biol Phys* 2005;61:993–1002.
- [14] Damber J-E. Endocrine therapy for prostate cancer. *Acta Oncol* 2005;44:605–9.
- [15] Möller TR, Einhorn N, Lindholm C, Nyhlén U, Svensson S, for the SBU Survey Group. Radiotherapy techniques in current use in Sweden. *Acta Oncol* 2003;42:376–86.
- [16] Hanlon AL, Watkins Bruner D, Peter R, Hanks GE. Quality of life study in prostate cancer patients treated with three-dimensional conformal radiation therapy: Comparing late bowel and bladder quality of life symptoms to that of the normal population. *Int J Radiat Oncol Biol Phys* 2001;49: 51–9.
- [17] Zelefsky MJ, Wallner KE, Ling CC, Raben A, Hollister T, Wolfe T, et al. Comparison of the 5-year outcome and morbidity of three-dimensional conformal radiotherapy versus transperineal permanent iodine-125 implantation for early-stage prostatic cancer. *J Clin Oncol* 1999;17:517–22.
- [18] Zelefsky MJ, Fuks Z, Hunt M, Yamada Y, Marion C, Ling CC, et al. High-dose intensity modulated radiation therapy for prostate cancer: early toxicity and biochemical outcome in 772 patients. *Int J Radiat Oncol Biol Phys* 2002;53:1111–6.
- [19] Kupelian PA, Reddy CA, Carlson TP, Willoughby TR. Dose/volume relationship of late rectal bleeding after external beam radiotherapy for localized prostate cancer: absolute or relative rectal volume? *Cancer J* 2002;8:62–6.
- [20] Kupelian PA, Reddy CA, Carlson TP, Altsman KA, Willoughby TR. Preliminary observations on biochemical relapse-free survival rates after short-course intensity-modulated radiotherapy (70 Gy at 2.5 Gy/fraction) for localized prostate cancer. *Int J Radiat Oncol Biol Phys* 2002;53:904–12.
- [21] Gardner BG, Zietman AL, Shipley WU, Skowronski UE, McManus P. Late normal tissue sequelae in the second decade after high dose radiation therapy with combined photons and conformal protons for locally advanced prostate cancer. *J Urol* 2002;167:123–6.
- [22] al-Abany M, Helgason AR, Cronqvist AK, Svensson C, Wersall P, Steineck G. Long-term symptoms after external beam radiation therapy for prostate cancer with three or four fields. *Acta Oncol* 2002;41:532–42.
- [23] Widmark A, Fransson P, Tavelin B. Self-assessment questionnaire for evaluating urinary and intestinal late side effects after pelvic radiotherapy in patients with prostate cancer compared with an age-matched control population. *Cancer* 1994;74:2520–32.
- [24] Dearnaley DP, Khoo VS, Norman AR, Meyer L, Nahum A, Tait D, et al. Comparison of radiation side-effects of conformal and conventional radiotherapy in prostate cancer: a randomised trial. *Lancet* 1999;353:267–72.
- [25] Lilleby W, Fossa SD, Waehre HR, Olsen DR. Long-term morbidity and quality of life in patients with localized prostate cancer undergoing definitive radiotherapy or radical prostatectomy. *Int J Radiat Oncol Biol Phys* 1999;43:735–43 (R1/154).
- [26] Chism DB, Horwitz EM, Hanlon AL, Pinover WH, Mitra RK, Hanks GE. Late morbidity profiles in prostate cancer patients treated to 79–84 Gy by a simple four-field coplanar beam arrangement. *Int J Radiat Oncol Biol Phys* 2003;55: 71–7.
- [27] Huang EH, Pollack A, Levy L, Starkschall G, Dong L, Rosen I, et al. Late rectal toxicity: dose-volume effects of

- conformal radiotherapy for prostate cancer. *Int J Radiat Oncol Biol Phys* 2002;54:1314–21.
- [28] Schultheiss TE, Lee WR, Hunt MA, Hanlon AL, Peter RS, Hanks GE. Late GI and GU complications in the treatment of prostate cancer. *Int J Radiat Oncol Biol Phys* 1997;37:3–11.
 - [29] Crook J, Esche B, Futter N. Effect of pelvic radiotherapy for prostate cancer on bowel, bladder, and sexual function: the patient's perspective. *Urology* 1996;47:387–94.
 - [30] Peeters ST, Heemsbergen WD, van Putten WL, Slot A, Tabak H, Mens JW, et al. Acute and late complications after radiotherapy for prostate cancer: results of a multicenter randomized trial comparing 68 Gy to 78 Gy. *Int J Radiat Oncol Biol Phys* 2005;61:1019–34.
 - [31] Heemsbergen WD, Hoogeman MS, Hart GA, Lebesque JV, Koper PC. Gastrointestinal toxicity and its relation to dose distributions in the anorectal region of prostate cancer patients treated with radiotherapy. *Int J Radiat Oncol Biol Phys* 2005;61:1011–8.
 - [32] al-Abany M, Helgason AR, Cronqvist AK, Lind B, Mavroidis P, Wersall P, et al. Toward a definition of a threshold for harmless doses to the anal-sphincter region and the rectum. *Int J Radiat Oncol Biol Phys* 2005;61:1035–44.
 - [33] Morris DE, Emami B, Mauch PM, Konski AA, Tao ML, Ng AK, et al. Evidence-based review of three-dimensional conformal radiotherapy for localized prostate cancer: an ASTRO outcomes initiative. *Int J Radiat Oncol Biol Phys* 2005;62:3–19.
 - [34] Slater JD, Rossi CJ, Jr, Yonemoto LT, Bush DA, Jabola BR, Levy RP, et al. Proton therapy for prostate cancer: the initial Loma Linda University experience. *Int J Radiat Oncol Biol Phys* 2004;59:348–52.
 - [35] Schulte RW, Slater JD, Rossi CJ, Slater JM. Value and perspectives of proton radiation therapy for limited stage prostate cancer. *Strahlenther Onkol* 2000;176:3–8.
 - [36] Shipley WU, Verhey LJ, Munzenrider JE, Suit HD, Urie MM, McManus PL, et al. Advanced prostate cancer: the results of a randomized comparative trial of high dose irradiation boosting with conformal protons compared with conventional dose irradiation using photons alone. *Int J Radiat Oncol Biol Phys* 1995;32:3–12.
 - [37] Ishikawa H, Tsuji H, Kamada T, Yanagi T, Wakatsuki M, Dhimazaki J, et al. A phase II trial using carbon ion radiotherapy (C-ion RT) for prostate cancer. *Proc Am Soc Clin Oncol* 2005;Abstr 4630.
 - [38] Lee M, Wynne C, Webb S, Nahum AE, Dearnaley D. A comparison of proton and megavoltage X-ray treatment planning for prostate cancer. *Radiother Oncol* 1994;33:239–53.
 - [39] Cella L, Lomax A, Miralbell R. Potential role of intensity modulated proton beams in prostate cancer radiotherapy. *Int J Radiat Oncol Biol Phys* 2001;49:217–23.
 - [40] Kupelian P, Kuban D, Thames H, Levy L, Horwitz E, Martinez A, et al. Improved biochemical relapse-free survival with increased external radiation doses in patients with localized prostate cancer: the combined experience of nine institutions in patients treated in 1994 and 1995. *Int J Radiat Oncol Biol Phys* 2005;61:415–9.
 - [41] Lundkvist J, Ekman M, Rehn Ericsson S, Jönsson B, Glimelius B. Proton therapy of cancer: Potential clinical advantages and cost-effectiveness. *Acta Oncol* 2005;44:850–61.
 - [42] Brenner DJ, Hall EJ. Fractionation and protraction for radiotherapy of prostate carcinoma. *Int J Radiat Oncol Biol Phys* 1999;43:1095–101.
 - [43] Brenner DJ, Martinez AA, Edmundson GK, Mitchell C, Thames HD, Armour EP. Direct evidence that prostate tumors show high sensitivity to fractionation (low alpha/beta ratio), similar to late-responding normal tissue. *Int J Radiat Oncol Biol Phys* 2002;52:6–13.
 - [44] Fowler JF. The radiobiology of prostate cancer including new aspects of fractionated radiotherapy. *Acta Oncol* 2005;44:265–76.
 - [45] Widmark A, Flodgren P, Damber J, Hellsten S, Cavallin-Stahl E. A systematic overview of radiation therapy effects in urinary bladder cancer. *Acta Oncol* 2003;42:567–81.
 - [46] Muren LP, Smaaland R, Dahl O. Conformal radiotherapy of urinary bladder cancer. *Radiother Oncol* 2004;73:387–98.
 - [47] Shelley MD, Barber J, Wilt T, Mason MD. Surgery versus radiotherapy for muscle invasive bladder cancer. *Cochrane Database Syst Rev* 2002;CD002079.
 - [48] Möller TR, Brorsson B, Ceberg J, Frodin JE, Lindholm C, Nylen U, et al. A prospective survey of radiotherapy practice 2001 in Sweden. *Acta Oncol* 2003;42:387–410.
 - [49] Battermann JJ. Results of d+T fast neutron irradiation on advanced tumors of bladder and rectum. *Int J Radiat Oncol Biol Phys* 1982;8:2159–64.
 - [50] Duncan W, Williams JR, Kerr GR, Arnott SJ, Quilty PM, Rodger A, et al. An analysis of the radiation related morbidity observed in a randomized trial of neutron therapy for bladder cancer. *Int J Radiat Oncol Biol Phys* 1986;12:2085–92.
 - [51] Errington RD, Ashby D, Gore SM, Abrams KR, Myint S, Bonnett DE, et al. High energy neutron treatment for pelvic cancers: study stopped because of increased mortality. *Bmj* 1991;302:1045–51.
 - [52] Pointon RS, Read G, Greene D. A randomised comparison of photons and 15 MeV neutrons for the treatment of carcinoma of the bladder. *Br J Radiol* 1985;58:219–24.
 - [53] Einhorn N, Trope C, Ridderheim M, Boman K, Sorbe B, Cavallin-Ståhl E. A systematic overview of radiation therapy effects in uterine cancer (corpus uteri). *Acta Oncol* 2003;42:557–61.
 - [54] Einhorn N, Trope C, Ridderheim M, Boman K, Sorbe B, Cavallin-Ståhl E. A systematic overview of radiation therapy effects in ovarian cancer. *Acta Oncol* 2003;42:562–6.
 - [55] Einhorn N, Trope C, Ridderheim M, Boman K, Sorbe B, Cavallin-Ståhl E. A systematic overview of radiation therapy effects in cervical cancer (cervix uteri). *Acta Oncol* 2003;42:546–56.
 - [56] Sténson S. Effects of high-energy protons on healthy organs and malignant tumours. Including a study of radiation protection by local hypoxia. Thesis. Uppsala University; 1969.
 - [57] Arimoto T, Kitagawa T, Tsujii H, Ohhara K. High-energy proton beam radiation therapy for gynecologic malignancies. Potential of proton beam as an alternative to brachytherapy. *Cancer* 1991;68:79–83.
 - [58] Tsujii H, Tsuji H, Inada T, Maruhashi A, Hayakawa Y, Takada Y, et al. Clinical results of fractionated proton therapy. *Int J Radiat Oncol Biol Phys* 1993;25:49–60.
 - [59] Kagei K, Tokuuye K, Okumura T, Ohara K, Shioyama Y, Sugahara S, et al. Long-term results of proton beam therapy for carcinoma of the uterine cervix. *Int J Radiat Oncol Biol Phys* 2003;55:1265–71.
 - [60] Molla M, Escude L, Nouet P, Popowski Y, Hidalgo A, Rouzaud M, et al. Fractionated stereotactic radiotherapy boost for gynecologic tumors: an alternative to brachytherapy? *Int J Radiat Oncol Biol Phys* 2005;62:118–24.
 - [61] Suit HD, Becht J, Leong J, Stracher M, Wood WC, Verhey L, et al. Potential for improvement in radiation therapy. *Int J Radiat Oncol Biol Phys* 1988;14:777–86.