

*Supplementary material for Tuxen M. K. et al. Phase II study of neoadjuvant pegylated liposomal doxorubicin and cyclophosphamide ± trastuzumab followed by docetaxel in locally advanced breast cancer. Acta Oncol 2014; 53:1440–1445.*

Supplementary Table I. Grade 3–4 toxicities (47 patients).

Adverse events grade 3–4	Number (%)
Palmar-plantar erythrodysesthesia (PPE)	16 (34)
Pain (myalgia + arthralgia)	15 (31)
Neurotoxicity	10 (21)
Fatigue	10 (21)
Febrile neutropenia	9 (19)
Mucositis	4 (8)
Nausea, vomiting	2 (4)
Hypersensitivity	1 (2)
Nail disorders	1 (2)

Supplementary Table II. Trials with neoadjuvant chemotherapy in primary breast cancer.

Author	Number of patients	Regimen	PR+ CR (%)	CR (%)	pCR (%)
Buzdar et al. [7]	87	P d1 q3w x 4	80	27	8
	87	FAC d1 q3w x 4	79	24	17
Miller et al. [8]	19	D d1 q2w x 3 → A d1 q2w x 3	89	32	16
	21	AD d1 q3w x 4	81	10	10
Bujanda et al. [9]	49	EC d1 +T d1,8,15,22 q4w x 3–4	84	33	15
Evans et al. [10]	180	AC d1 q3w x 6	61	17	24
	183	AD d1 q3w x 6	66	20	21
Kelly et al. [11]	37	AC d1 q3w x 4 → TP d1 q1w x 12	86	56	19
Rocca et al. [12]	58	E d1,2 + CIS d1 q3w + FU daily x 6	58		41
Tiezzi et al. [13]	60	ED d1 q3w	77	15	5
Gogas et al. [14]	35	P+ PLD d1 q3w x 6	71	17	9
Rossi et al. [15]	35	P d1,8 + PLD d1 q2w x 4	74	8	9
Artioli et al. [16]	50	G d1,8 + D d8 + PLD d8 q3w x 4–6	74	26	16
Dellapasqua et al. [17]	29	PLD d1 q2w x 8 + C daily	62	0	3
Taguchi et al. [18]	47	P d1,8,15 q3w x 4 → FEC d1 q3w x 4	85	38	28
Therasse et al. [19]	224	C d1 – 14 + EF d1,8 q4w x 6	89	31	14
	224	EC (high dose) d1 q3w x 6	91	27	10
Gianni et al. [20,21]	117 (HER2 positive)	T+AP d1 q3w x 3 → T+P d1 q3w x 4 → T+CMF d1 q4w x 3	87	60	38
	118 (HER2 positive)	AP d1 q3w x 3 → P d1 q3w x 4 → CMF d1 q4w x 3	74	51	19
	99 (HER2 negative)	AP d1 q3w x 3 → P d1 q3w x 4 → CMF d1 q4w x 3	71	25	16

A, doxorubicin; C, cyclophosphamide; CIS, cisplatin; CR, complete response; D, docetaxel; E, epirubicin; F, 5-fluorouracil; G, gemcitabine; M, methotrexate; P, paclitaxel; pCR, pathological complete response; PLD, pegylated liposomal doxorubicin; PR, partial response; T, trastuzumab; V, vincristine.



Supplementary Figure 1. Patient with palmar-plantar erythrodysesthesia.