

ORIGINAL ARTICLE

Clinical outcome with bevacizumab in patients with recurrent high-grade glioma treated outside clinical trials

SILVIA HOFER¹, KATARZYNA ELANDT³, RICHARD GREIL⁴, ANDREAS F. HOTTINGER⁵, URS HUBER⁶, DIETER LEMKE⁷, CHRISTINE MAROSI³, ADRIAN OCHSENBEIN⁸, JOSEF PICHLER⁹, ULRICH ROELCKE¹⁰, PATRIK WEDER¹¹, THILO ZANDER¹², WOLFGANG WICK⁷ & MICHAEL WELLER²

¹Department of Oncology, University Hospital Zurich, Switzerland, ²Department of Neurology, University Hospital Zurich, Switzerland, ³Department of Internal Medicine I, University Hospital Vienna, Austria, ⁴Third Medical Department at the Private Medical University Hospital Salzburg, Austria, ⁵Department of Oncology University Hospital Geneva, Switzerland, ⁶Swiss Tumor Institute, Onkozentrum Klinik im Park, Zurich, Switzerland, ⁷Department of Neurooncology, National Center for Tumour Disease and Neurology Clinic, University Hospital Heidelberg, Germany, ⁸Department of Oncology, University Hospital Bern, Switzerland, ⁹Department of Internal Medicine and Neurooncology, Landesnervenklinik Linz, Austria, ¹⁰Department of Neurooncology and Medical Oncology, Kantonsspital Aarau, Switzerland, ¹¹Department of Oncology, Kantonsspital St. Gallen, Switzerland and ¹²Department of Oncology, Kantonsspital Luzern, Switzerland

Abstract

Background. Patients with recurrent high-grade glioma (HGG) have a poor prognosis and there is no defined standard of care. High levels of vascular endothelial growth factor (VEGF) expressed in HGG make the anti-VEGF monoclonal antibody bevacizumab (BEV) of particular interest. **Patients and methods.** In an ongoing registry data were collected from patients who have received BEV for the treatment of recurrent HGG. The primary objective was the identification of any clinical benefit as assessed by change in Karnofsky Performance Score (KPS), decreased steroid use and duration of treatment. **Results.** Two hundred and twenty-five patients with HGG were included (176 glioblastoma; 49 anaplastic glioma; median age 52 years). KPS improved in 10% of patients and remained stable in 68%. Steroids were stopped in 37.6% of patients. Median duration of treatment was 5.5 months; 19.1% of patients were treated for more than 12 months. Median overall survival from beginning of BEV treatment was 8.5 months. At the time of analysis, 169 patients (75.1%) had died and 56 patients (24.9%) were alive. Only 21 patients (9.3%) discontinued treatment due to toxicity. **Conclusions.** Our data reveal valuable palliation with preservation of KPS and an option for steroid withdrawal in patients treated with BEV, supporting the role of this therapy in late-stage disease.

High-grade gliomas (HGG) account for approximately 23.9% of all primary brain and central nervous system tumours and have an annual incidence rate of 5–7 cases per 100 000 population. The most common type of HGG is glioblastoma (WHO grade IV), accounting for approximately 53.8% of all gliomas [1].

Glioblastoma is associated with poor prognosis, with patients displaying a median survival of 15 months only when treated with current standard of care, involving maximal safe surgical resection followed by radiotherapy with concomitant and adjuvant temozolomide (TMZ) and individual therapy at recurrence [2,3].

Regardless of initial treatment, recurrence of HGG is inevitable. A variety of therapies have been investigated for patients with recurrent disease. According to the NCCN guidelines (www.nccn.org), patients can be offered drugs such as TMZ, nitrosoureas, PCV (procarbazine, CCNU and vincristine), cyclophosphamide, platinum-based regimens or bevacizumab (BEV; Avastin®, Genentech/Roche, Switzerland) alone or in combination with chemotherapy. Second surgery, second courses of radiation therapy and experimental treatment options are also offered at some centres. An overview by the North American Brain Tumour Consortium (NABTC)

examined data from 596 patients (159 patients with grade III tumours and 437 patients with grade IV tumours) enrolled in phase II trials for recurrent HGG. Median survival durations, measured from time of registration, were 39 weeks (8.9 months) and 30 weeks (6.9 months) for grade III and grade IV tumours, respectively (Table I) [4].

To date, there is no established standard of care for patients with recurrent HGG. Salvage treatment remains palliative and should therefore meet palliative needs. High levels of vascular endothelial growth factor (VEGF) expression found in HGG [5] have made the VEGF monoclonal antibody BEV of particular interest in therapeutic strategies, both as an adjunct to radiotherapy [6] and in combination with chemotherapy, initially with irinotecan in analogy to previous experiences in colorectal cancer [7–9].

In 2009, the US Food and Drug Administration (FDA) and Swissmedic granted accelerated approval of BEV for the treatment of recurrent glioblastoma, whereas there is currently no such approval in the European Union [10]. The FDA and Swissmedic approval of BEV in recurrent glioblastoma is mainly based on data from two uncontrolled phase II trials [11,12]. The only randomised phase II trial was an open-label, non-comparative, multi-centre trial of BEV alone and in combination with irinotecan in 167 recurrent glioblastoma patients. Median progression-free survival (PFS) was 4.2 months (BEV alone) and 5.6 months (BEV with irinotecan). Median overall survival (OS) was 9.2 months (BEV alone; 95% CI, 8.2–10.7) and 8.7 months (BEV with irinotecan; 95% CI, 7.8–10.9) [11]. The second phase II trial, a single-centre study, investigated single-agent BEV followed by BEV plus irinotecan in 48 heavily pre-treated patients with recurrent glioblastoma. Median PFS was 16 weeks (3.7 months) and median OS was 31 weeks (7.1 months) [12].

Outside of clinical trials, BEV-based regimens are now widely used in several countries where the drug is reimbursed for recurrent malignant glioma.

Our ongoing registry aims at addressing clinical outcome of BEV-based treatment in recurrent HGG after one or multiple prior therapies, using community-based data from patients treated at 30 centres in Switzerland, Austria and Germany.

Methods

In an ongoing and regularly updated registry (starting in August 2006), data on patient characteristics are collected from unselected patients with recurrent HGG treated at 30 centres in Switzerland, Austria and Germany. The decision to treat a patient with recurrent HGG with a BEV-based regimen and the frequency of treatment assessment either radiologically or clinically is the treating physician's choice. Patients are registered at the time of first BEV application, follow-up is until death.

Anonymous data are sampled using an institutional review board-approved written questionnaire. Baseline patient characteristics, medical history regarding the course of HGG, prior treatment regimens and patient outcomes focusing on BEV treatment are recorded.

The main objective of this analysis was the identification of clinical benefit as assessed by change in Karnofsky Performance Score (KPS; baseline KPS and best KPS during BEV-based treatment) and steroid use, as well as duration of treatment. We partitioned KPS into three groups and defined group I as good to excellent (KPS 80–100%) group II as moderate (KPS 60–80%) and group III as poor (KPS < 60%). OS from start of BEV treatment is also of interest; other parameters include reasons for BEV discontinuation and safety. Response rates are not assessed in this cohort due to concerns

Table I. Common treatment regimens for recurrent glioblastomas.

	NABTC Phase II [4]	TMZ 7/7 [16]	CCNU [15]	BEV [11]	BEV/ CPT-11 [11]
Number of patients (n)	437*	64*	92	85	82
Median age (years)	52 (21–84)	51	55.3	54 (23–78)	57 (23–78)
KPS (%)					
90–100	40	72	49	44.7	37.8
70–80	54	–	50	55.3	62.2
In first relapse (%)	50	–	77	81.2	80.5
Time from diagnosis to treatment (months)	–	8.5	12	8.6	9.8
Median duration of response (months)	–	–	2.8–9.6	5.6	4.3
Median PFS (months)	1.6/3.4**	5.5	1.64	4.2	5.6
Median OS (months)	6/7**	8.7	7.1	9.2	8.7

*Patient cohort with grade IV tumors; **Non-temozolomide/temozolomide regimen, respectively.

NABTC, North American Brain Tumor Consortium; TMZ 7/7 temozolomide one week on, one week off; CCNU, lomustine; BEV, bevacizumab; CPT-11, irinotecan; KPS, Karnofsky Performance Score; PFS, progression-free survival; OS, overall survival.

Table II. Patient demographics.

	Glioblastoma	Other HGG	All
Number of patients, n (%)	176 (100)	49 (100)	225 (100)
Gender, n (%)			
Male	107 (60.8)	30 (61.2)	137 (60.9)
Female	69 (39.2)	19 (38.8)	88 (39.1)
Age (years)			
Median	55	46	52
Range	19–79	20–70	19–79
Median time from initial diagnosis to start of BEV, days (months)	305 (10.0)	1328 (43.5)	335 (11.0)

HGG, high-grade glioma (anaplastic astrocytoma, oligodendroglioma and mixed glioma); BEV, bevacizumab.

regarding response criteria for anti-angiogenic treatment and ‘pseudoresponses’, whereby the pseudoresponse reflects a normalisation of abnormally permeable microvessels rather than true tumour shrinkage [13]. In addition, there is a discrepancy between high response rates on conventional radiographic criteria and moderate OS in published phase II trials.

Statistical analysis

Patient characteristics such as gender, age, tumour histology at initial diagnosis and steroid use at the start and during treatment with BEV were assessed. Descriptive statistics using SAS version 8.0 were used to assess: median duration from diagnosis to the date of first BEV treatment administration (days), median duration of BEV therapy (months), KPS before therapy and best score during BEV therapy, and reason for discontinuation of BEV. Time to death after initiation of BEV therapy (days) and

time from diagnosis to death (days) were determined using Kaplan–Meier analysis. All analyses are based on a data cut-off point of 31 August 2010.

Results

A total of 225 patients with recurrent HGG were included in this registry [176 patients with glioblastoma and 49 patients with other HGG (anaplastic gliomas of astrocytic, oligodendroglial or mixed phenotypes)]. These diagnoses refer to the histological diagnoses made at first surgery. Patient demographics are shown in Table II. The median age of patients was 52 years (range 19–79). Overall, 20% of patients received BEV monotherapy and 80% were treated with chemotherapy plus BEV. The main combination partner was irinotecan (82% of all combinations), followed by TMZ (9%), lomustine (4.5%) and pegylated liposomal doxorubicin (4.5%). Four of 225 patients received concurrent surgery or irradiation during BEV treatment.

Table III. Changes in KPS and corticosteroid use in patients prior to and during BEV.

	Glioblastoma (n = 176)	Other HGG (n = 49)	All (n = 225)
Steroids at start of treatment, n (%)			
Yes	123 (69.9)	26 (53.1)	149 (66.2)
No	42 (23.9)	15 (30.6)	57 (25.3)
Missing	11 (6.2)	8 (16.3)	19 (8.5)
Steroid interruption during treatment, n (%)			
Yes	48 (27.3)	8 (16.3)	56 (24.9)
No	73 (41.5)	17 (34.7)	90 (40.0)
Not applicable* or missing	55 (31.2)	24 (49.0)	79 (35.1)
KPS prior to treatment, n (%)			
I (80–100)	69 (39.2)	21 (42.9)	90 (40.0)
II (60–70)	63 (35.8)	13 (26.5)	76 (33.8)
III (<60)	30 (17.1)	9 (18.4)	39 (17.3)
Missing	14 (8.0)	6 (12.2)	20 (8.9)
KPS change during treatment, n (%)			
Improved	20 (11.4)	3 (6.1)	23 (10.2)
Stable	120 (68.1)	34 (69.4)	154 (68.4)
Worse	16 (9.1)	4 (8.2)	20 (8.9)
Missing	20 (11.4)	8 (16.3)	28 (12.5)

HGG, high-grade glioma (anaplastic astrocytoma, oligodendroglioma and mixed glioma); KPS, Karnofsky Performance Score, BEV bevacizumab.

*not applicable, where no steroids at start of BEV treatment.

Table IV. Duration of BEV treatment.

	Glioblastoma (n = 176)	Other HGG (n = 49)	All (n = 225)
Duration of BEV treatment (months)			
Median	5.45	6.2	5.5
Range	0.5–39	0.5–36	0.5–39
Patients on treatment, n (%)			
≥ 6 months	86 (48.9)	25 (51.0)	111 (49.3)
≥ 1 year	31 (17.6)	12 (24.5)	43 (19.1)
≥ 2 year	6 (3.4)	4 (8.2)	10 (4.4)

HGG, high-grade glioma (anaplastic astrocytoma, oligodendroglioma and mixed glioma); BEV, bevacizumab.

Overall, 40% of patients had a KPS of $\geq 80\%$ at the start of BEV treatment; 10% of patients showed improvement in their KPS during treatment (from one group to a better group), whereas 9% showed a decline; no data were available from 13% of patients (Table III). No difference was found between WHO grade III and IV tumours in this respect. During treatment, steroids were stopped in 56 of 149 HGG patients (37.6%) who were receiving steroids at baseline (Table III).

Median duration of BEV-based treatment (with or without additional chemotherapy) was 5.5 months (range 0.5–39; Table IV). Almost half of the patients (49.3%) were treated for more than six months, 19% for more than one year and 10 patients (4.4%) were treated for more than two years.

Overall, median OS from beginning of BEV treatment was 8.5 months, with a median OS of 8.3 months in patients with glioblastoma and 9.1 months in patients with other HGG (Table V). A Kaplan–Meier plot of time-to-death analysis from start of BEV treatment is shown in Figure 1. At the time of this analysis (31 August 2010), 169 patients (75.1%) had died and 56 patients (24.9%) remained alive.




Figure 1. Overall survival from beginning of bevacizumab (BEV) treatment.

There was no clear difference in median duration of BEV-based treatment or median OS from beginning of BEV treatment between the following age groups; <50 years, 51–60 years and >60 years (data not shown).

The main reason for discontinuation of BEV treatment was disease progression. Overall, 76% of patients stopped treatment because of clinical or radiological disease progression or both. A total of 21 patients (9.3%) discontinued treatment due to toxicity (Table VI). Three fatal CNS bleeding complications and two fatalities due to impaired wound healing accounted for toxic deaths in 2.2% of the patient population. Other treatment-related toxicities, such as hypertension and thromboembolic events, were within the range expected based on phase II trials [11].

Discussion

To date, there is no standard of care in recurrent HGG. Recurrences often become symptomatic with increased intracranial pressure and steroids are regularly the treatment of choice to relieve these symptoms [14]. However, steroids are not generally able to maintain their beneficial effect for a long period of time and can also cause disabling side-effects. In contrast to chemotherapeutic agents available for recurrent HGG, the VEGF antibody BEV has the potential to rapidly lower intracranial pressure by modifying vascular permeability. BEV is a well-tolerated drug, causing severe side-effects in only a low percentage of patients.

Our observational registry of an unselected patient cohort with recurrent and pre-treated HGG from 30 different centres aims to reflect community-based experience with BEV. Evidence for the efficacy of BEV-containing treatment regimens for recurrent HGG stems primarily from unicentric or oligocentric phase II trials. Our experience of a larger and less selected population seems of importance and reproduces benefits for patients with recurrent HGG. Within this patient population only a few patients showed a decline in KPS during BEV treatment. Median duration of treatment was 5.5 months and half of all patients were judged to derive benefit from BEV therapy for at least six months and 19% for at least one year. Median OS of over eight months from the start of BEV treatment for all patients confirms the results from phase II trials. In this cohort of recurrent HGG patients, BEV was given as up to and including the last line of therapy. Therefore, it is not likely that OS was confounded by additional therapeutic approaches given after BEV treatment. However, the analysis of such registry data has limitations, including the possibility of a reporting bias of patients in general and underreporting of adverse events.

Table V. Overall survival in HGG patients treated with BEV at recurrence.

	Glioblastoma (n = 176)	Other HGG (n = 49)	All (n = 225)
OS from start of BEV treatment, days			
Median, days (months)	252 (8.3)	277 (9.1)	259 (8.5)
95% CI	216; 305	190; 372	224; 305
OS from diagnosis			
Median, days (months)	668 (21.9)	1972 (64.7)	732 (24.0)
95% CI	569; 762	890; 3485	660; 846

HGG; high-grade glioma (anaplastic astrocytoma, oligodendroglioma and mixed glioma); BEV, bevacizumab; OS, overall survival; CI, confidence interval.

Although an often rapid symptomatic improvement with BEV in recurrent HGG is appealing there are some issues that have not yet been resolved, including mechanisms of action and resistance to BEV in HGG, dosing, optimal drug combination and response assessment [13]. Some of these issues will be addressed in a phase II trial from the EORTC Brain Tumour Group which will explore the sequence of BEV and lomustine in glioblastoma patients at first recurrence. In this four-arm study of 249 patients, OS at 12 months will be the primary endpoint (EORTC 26101; www.eortc.be).

In summary, BEV met with the criteria which are claimed for a palliative treatment for HGG patients, that is the preservation of a stable neurological status, withdrawal of steroids and a good tolerability with an acceptable incidence of side-effects. OS confirms

Table VI. Reasons for discontinuation of BEV treatment.

	All (n = 225)
Progression (total PD), n (%)	171 (76.0)
Clinical PD	33 (14.7)
Radiological PD	38 (16.9)
Clinical and radiological PD	100 (44.4)
Toxicity , n (%)	21 (9.3)
Toxicities that led to death (fatal)	5 (2.2)
CNS bleeding (3 fatal, included above)	6 (2.6)
Bleeding other location	2 (0.8)
Wound-healing complications (2 fatal, included above)	5 (2.2)
Thromboembolism (venous or arterial)	4 (1.7)
Hand-foot syndrome [‡]	1 (0.4)
Proteinuria	2 (0.8)
Diarrhea [¥]	1 (0.4)
Other reasons , n (%)	24 (10.6)
Not assessed	9 (4.0)
Patient wish	11 (4.8)
Sustained tumor response	4 (1.7)

PD, progressive disease.

*Multiple reasons for BEV discontinuation were possible.

[‡]Bevacizumab combined with pegylated liposomal doxorubicin.

[¥]Bevacizumab with irinotecan.

findings from phase II clinical trials [11], suggesting that BEV enriches the limited repertoire of medical treatment options for HGG.

Acknowledgements

We sincerely thank Barbara Hofstetter for her invaluable support during the project and critical review of the manuscript. We also thank Laura McDonagh for her editorial support. This work was supported in part by an unrestricted grant from Roche (USZ-NEURO-15122008). SH receives research support and honoraria for advisory board participation and lectures from Roche and Schering Plough. AO is a member of a Roche advisory board. PW is a member of a Schering-Plough advisory board. WW receives lecture honoraria and consultancy honoraria from Roche. MW receives research support and honoraria for advisory board participation and lectures from Merck Serono, Roche and Schering Plough.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

References

- [1] Central Brain Tumour Registry of the United States. CBTRUS Statistical Report: Primary brain and central nervous system tumours diagnosed in the United States in 2004–2006. 2010.
- [2] Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med* 2005;352:987–96.
- [3] Stupp R, Hegi ME, Mason WP, van den Bent MJ, Taphoorn MJ, Janzer RC, et al. Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial. *Lancet Oncol* 2009;10:459–66.
- [4] Lamborn KR, Yung WK, Chang SM, Wen PY, Cloughesy TF, DeAngelis LM, et al. Progression-free survival: An important end point in evaluating therapy for recurrent high-grade gliomas. *Neuro Oncol* 2008;10:162–70.
- [5] Salmaggi A, Eoli M, Frigerio S, Silvani A, Gelati M, Corsini E, et al. Intracavitary VEGF, bFGF, IL-8, IL-12 levels in primary and recurrent malignant glioma. *J Neurooncol* 2003; 62:297–303.
- [6] Gutin PH, Iwamoto FM, Beal K, Mohile NA, Karimi S, Hou BL, et al. Safety and efficacy of bevacizumab with hypofractionated stereotactic irradiation for recurrent malignant gliomas. *Int J Radiat Oncol Biol Phys* 2009; 75:156–63.
- [7] Desjardins A, Reardon DA, Herndon JE 2nd, Marcello J, Quinn JA, Rich JN, et al. Bevacizumab plus irinotecan in recurrent WHO grade 3 malignant gliomas. *Clin Cancer Res* 2008;14:7068–73.
- [8] Vredenburgh JJ, Desjardins A, Herndon JE 2nd, Marcello J, Reardon DA, Quinn JA, et al. Bevacizumab plus irinotecan in recurrent glioblastoma multiforme. *J Clin Oncol* 2007; 25:4722–9.

- [9] Vredenburgh JJ, Desjardins A, Herndon JE 2nd, Dowell JM, Reardon DA, et al. Phase II trial of bevacizumab and irinotecan in recurrent malignant glioma. *Clin Cancer Res* 2007;13:1253–9.
- [10] Wick W, Weller M, van den BM, Stupp R. Bevacizumab and recurrent malignant gliomas: A European perspective. *J Clin Oncol* 2010;28:e188–9.
- [11] Friedman HS, Prados MD, Wen PY, Mikkelsen T, Schiff D, Abrey LE, et al. Bevacizumab alone and in combination with irinotecan in recurrent glioblastoma. *J Clin Oncol* 2009; 27:4733–40.
- [12] Kreisl TN, Kim L, Moore K, Duic P, Royce C, Stroud I, et al. Phase II trial of single-agent bevacizumab followed by bevacizumab plus irinotecan at tumour progression in recurrent glioblastoma. *J Clin Oncol* 2009;27:740–5.
- [13] Dhermain FG, Hau P, Lanfermann H, Jacobs AH, van den Bent MJ. Advanced MRI and PET imaging for assessment of treatment response in patients with gliomas. *Lancet Neurol* 2010;9:906–20.
- [14] Roth P, Wick W, Weller M. Steroids in neurooncology: Actions, indications, side-effects. *Curr Opin Neurol* 2010.
- [15] Wick W, Puduvalli VK, Chamberlain MC, van den Bent MJ, Carpentier AF, Cher LM, et al. Phase III study of enzastaurin compared with lomustine in the treatment of recurrent intracranial glioblastoma. *J Clin Oncol* 2010;28: 1168–74.
- [16] Wick A, Felsberg J, Steinbach JP, Herrlinger U, Platten M, Blaschke B, et al Efficacy and tolerability of temozolomide in an alternating weekly regimen in patients with recurrent glioma. *J Clin Oncol* 2007; 25:3357–61.